

Designing Optimal Binary Rating Systems

Nikhil Garg
Stanford University
nkgarg@stanford.edu

Ramesh Johari
Stanford University
rjohari@stanford.edu

October 30, 2018

Abstract

Modern online platforms rely on effective rating systems to learn about items. We consider the optimal design of rating systems that collect *binary feedback* after transactions. We make three contributions. First, we formalize the performance of a rating system as the speed with which it recovers the true underlying ranking on items (in a large deviations sense), accounting for both items’ underlying match rates and the platform’s preferences. Second, we provide an efficient algorithm to compute the binary feedback system that yields the highest such performance. Finally, we show how this theoretical perspective can be used to empirically design an implementable, approximately optimal rating system, and validate our approach using real-world experimental data collected on Amazon Mechanical Turk.

1 Introduction

Rating and ranking systems are everywhere, from online marketplaces (e.g., 5 star systems where buyers and sellers rate each other) to video platforms (e.g., thumbs up/down systems on YouTube and Netflix). A large literature on ratings systems has shown that they are uninformative in practice [3, 5, 12, 14, 19, 20, 27, 29, 30, 33]. One recurring pattern is that ratings tend to *binarize* – with most raters only using the extreme choices on the provided rating scale, and the vast majority of ratings receiving the best possible rating. For example, 75% of reviews on Airbnb receive a perfect rating of 5 stars [14]. Furthermore, several platforms have adopted a binary rating system, in which a user rates her experience as either positive or negative. Given the prevalence of binary feedback (either de facto or by design), in this work we investigate the optimal *design* of such binary rating systems so that the platform can learn as fast as possible about the items being rated.

The rating pipeline often works as follows: A buyer enters a platform and *matches* with an item (e.g. chooses a video to watch on Youtube, is paired with a driver on Uber, or selects a home on AirBnB). She has an experience (e.g. a view, a ride, or a stay). At the end of her experience, the platform asks her to *rate* her experience, i.e. it asks her a question. In a binary system, she either indicates that her experience was positive or negative. She then leaves. The platform uses the ratings it has received to score the quality of items, potentially showing such scores to future buyers.

By *designing* such a system, we mean: the platform can influence how the buyer rates – how likely she is to give a positive rating, conditional on the quality of her experience. It can do so by asking her different questions, e.g. “Was this experience above average” or “Was this experience the worst you’ve ever had?”. Different questions shift the probabilities at which items of various qualities receive positive ratings.

Our first question is: *what is the structure of optimal binary feedback?* A rating system in which every buyer gives positive ratings after each match, independent of item quality, will fail to learn anything about the items. Clearly, better items should be more likely to receive positive ratings than worse ones. But how much more likely? We answer this question.

Informally, suppose we have a set of items that match with buyers over time (at potentially differing rates), and we wish to rank the items by their true quality $\theta_i \in [0, 1]$. The platform cannot observe θ_i , however. Rather, in our model, after each match, an item with quality θ_i receives a positive rating with probability $\beta(\theta_i)$, and negative otherwise. In other words, the platform observes, for each item i , a sequence of ratings that are each $\text{Bernoulli}(\beta(\theta_i))$. Such ratings are the only knowledge the platform has about items. The platform ranks the items according to the percentage of its ratings (samples) that were positive. The function $\beta : [0, 1] \mapsto [0, 1]$ affects how quickly the platform learns the true ranking, and it prefers to maximize the learning rate. We show how to calculate an optimal β .

As an example, consider just three items with qualities $\theta_a > \theta_b > \theta_c$, and β such that $\beta(\theta_a) = 0.5$ and $\beta(\theta_c) = 0.1$, i.e. item a gets positive ratings after 50% of its matches, and item c after 10% of its matches. It is unclear what $\beta(\theta_b)$ should be. Trivially, $.1 < \beta(\theta_b) < .5$. Otherwise, even with infinitely many ratings the platform will mis-rank the items.

But can we be more precise? If $\beta(\theta_b) = .49$, it will take many ratings of both items a and b to learn that $\theta_a > \theta_b$, but only a few from c to learn that $\theta_c < \theta_b$. That may be good if the platform cares more about identifying the worst item, but not if it cares most about identifying the best. It may also be fine if items a and b match much more often with buyers than item c . Clearly, the optimal value for $\beta(\theta_b)$ is objective and context dependent. Of course, the problem becomes more challenging with more items i for which $\beta(\theta_i)$ must be chosen. Lastly, in this example, one might intuitively think $\beta(\theta_b) = 0.3$ is optimal by symmetry when the items matter equally and matching rates are identical. This guess is incorrect. The optimal is $\beta(\theta_b) \approx 0.28$, due to the nature of binomial variance.

In this work, we first formalize the above problem and show how to find an optimal $\beta(\theta)$, jointly for a whole set of items $[0, 1]$. β changes with the platform’s objective and the underlying matching rates of the items. Jumping ahead, Figure 1 shows optimal β in various settings under our model. For a platform that prioritizes accurately finding the worst sellers, for example, the top half of items should each get positive ratings at least 80 + % of the time; it is more important for the bottom half of items to be separated from one another, i.e. get positive ratings at differing percentages.

Once we have calculated the optimal rating function β (given context on the platform goals and matching rates), what should we do with it? Remember our original goal of *designing* a rating system.

Our second question is: *How does a platform build a rating system such that buyers behave near-optimally, i.e. according to a calculated β ?* The platform cannot directly control buyer rating behavior – it cannot force users to only give positive ratings to the best items. Rather, it has to ask *questions* such that, for each item θ , a fraction $\beta(\theta)$ of raters will give an item of quality θ a positive rating, and $1 - \beta(\theta)$ will give it a negative rating. For example, by asking the question, “Is this the best experience you’ve had on AirBnB,” the platform would be inducing rating behavior such that $\beta(\theta)$ is small for most θ . Most platforms today ask vague questions, such as thumbs up/down, that in practice lead to most items getting positive ratings most of the time. We show this is highly suboptimal if the platform wants to be able to rank its items quickly. We show how a platform would choose such questions using empirical rating behavior to build a near-optimal rating system.

Our main contributions and paper outline are:

1. **Rating system design as information maximization.** In Sections 3.1-3.2, we formulate

the design of rating systems as a problem of information maximization. In particular, a good rating system should recover the true ranking over items, and converge quickly in the number of ratings.

2. **Computing an optimal rating feedback function β .** In Section 3.3, we develop an efficient algorithm that calculates the optimal rating feedback function β , which depends on item search rates and the platform’s objective. We show how computing the optimal β enables quantitative insights and principled comparisons between designs.
3. **Real-world system design.** In Section 4, we show how a platform can use a simple experiment and existing data to empirically design a rating system that performs near-optimally, and to audit the current system’s performance. In Section 5, we demonstrate the value of this approach through an experiment on Mechanical Turk.

2 Related work

Much work, both empirical and model-based, discusses problems with and solutions for existing rating systems [2, 3, 5, 12, 14, 15, 19, 20, 22, 27, 29, 30, 33]. To our knowledge, we are the first to formalize a rating system design problem and then show how one can use empirical data to optimize such systems. In a related paper in submission [16], we test behavioral insights using a live trial on a large online labor platform and develop a related design problem for a multiple-choice system, which proves far less tractable.

Other works also tackle learning rates in online platforms, as we do [1, 4, 21, 23]. However, these works modify *how often buyers and items match* in order to learn about items, while we view the matching process as given and modify *the rating system* itself. The solutions are complementary.

The bandits literature also often seeks to rank items by quality based on a sequence of observations [24, 25, 28, 32]. Our problem is the *inverse* of the standard bandit setting – given an external arm-pulling policy, we design the feedback at each arm. (To be clear, the *rating* is not the same as a *reward*; buyers often give positive ratings after bad experiences.) Our specific theoretical framework is similar to that of Glynn and Juneja [17], who optimize a large deviations rate to derive an arm-pulling policy for best arm identification.

Also related is the literature surrounding the “twenty questions” interpretation of Shannon entropy [6, 8], which seeks a set of questions that can be used to identify a single item from its distribution. Dagan et al. [8] show how to almost match the performance of Huffman codes by asking only comparison or equality comparisons. Our work differs in two key respects: first, we seek to rank an entire set of items as opposed to identifying just a single item, with items receiving positive ratings given a question at differing probabilities; second, we consider non-adaptive policies (i.e. the platform cannot change its rating form in response to what it knows about an item already).

3 Model and optimization

In this section, we begin by formalizing the model, and then we show how to optimize the rating function in order to maximize the rate of learning. The focus is on finding an optimal $\beta : [0, 1] \mapsto [0, 1]$, i.e. a *function* that maps a item’s quality θ to the probability it should receive a positive rating. This section requires no data: we characterize the *optimal* rating system one could ever hope to build.

3.1 Model and problem specification

Our model is constructed to emphasize the learning rate of the platform through its rating system. Time is discrete ($k = 0, 1, 2, \dots$). Informally, the model is as outlined in the introduction: there is a set of items with quality $\theta \in [0, 1]$. Each time step, buyers enter and match with items, leaving a rating after the match according to $\beta(\theta)$. The platform keeps track of the ratings of each item and ranks the items by their rating scores. Formally, the key components are:

Items. The system consists of a set $[0, 1]$ of items, where each item is associated with a unique (but unknown) quality $\theta \in [0, 1]$; i.e., the system consists of a *continuum* of a unit mass of items whose qualities are unknown and uniform¹ in $[0, 1]$. Below, we *discretize* the continuous quality space $[0, 1]$ into M types, to calculate a stepwise increasing β . We'll make clear why we introduce a continuum but then discretize.

Matching with buyers. Items accumulate ratings over time by matching with buyers. In particular, we assume the existence of a nondecreasing *match function* $g(\theta)$, where an item θ receives $n_k(\theta) = \lfloor kg(\theta) \rfloor$ matches, and thus ratings, up to time k . In other words, item θ is matched approximately every $\frac{1}{g(\theta)}$ time steps. $g(\theta) \leq 1$ and bounded away from 0, i.e. $\exists c: g(\theta) > c > 0$. This accumulation captures the feature that better items may be more likely to match.

Ratings. The key quantity for our subsequent analysis is the *probability of a positive rating* for each θ , $\beta(\theta) \triangleq \Pr(\text{positive rating}|\theta)$. Let $y_\ell(\theta) \sim \text{Bernoulli}(\beta(\theta))$ be the rating an item of quality θ receives at the ℓ th time it matches.

Aggregating ratings and ranking sellers. These ratings are aggregated into a *reputation score*, $x_k(\theta)$, at each time k . The score is the fraction of positive ratings received up to time k : $x_k(\theta) \triangleq \frac{1}{n_k(\theta)} \sum_{\ell=0}^{n_k(\theta)} y_\ell(\theta)$ with $x_0(\theta) \triangleq 0$ for all θ . Thus, $x_k(\theta) \sim \frac{1}{n_k(\theta)} \text{Binomial}(\beta(\theta), n_k(\theta))$.

System state. The state of the system is given by a joint distribution $\mu_k(\Theta, X)$, which gives the *mass* of items of quality $\theta \in \Theta \subset [0, 1]$ with aggregate score $x_k(\theta) \in X \subset [0, 1]$ at time k . Because our model is a continuum, the evolution of the system state μ_k follows a deterministic dynamical system.

We have so far described these dynamics at the level of individual items; however, all such statements should be interpreted as describing the evolution of the joint distribution μ_k . The state update for μ_k is determined by the *mass* of items who match and the *distributions* of their ratings. A formal description of the system state evolution is in Appendix Section B.1.

Platform objective. We assume that the platform wants the ranking of items by observed reputation score to reflect the underlying true quality ranking as closely as possible. We primarily investigate how β influences the quality of the observed ranking.

Formally, given β and $\theta_1 > \theta_2$, define $P_k(\theta_1, \theta_2|\beta)$:


$$P_k(\theta_1, \theta_2|\beta) = \mu_k(x_k(\theta_1) > x_k(\theta_2)|\theta_1, \theta_2) - \mu_k(x_k(\theta_1) < x_k(\theta_2)|\theta_1, \theta_2) \quad (1)$$

This expression captures the accuracy of the ranking according to observed score. When $\theta_1 > \theta_2$ but $x_k(\theta_1) < x_k(\theta_2)$, the observed score ranking mistakenly swaps the ordering of items θ_1 and θ_2 . Thus, for a good rating system, $P_k(\theta_1, \theta_2|\beta)$ is large.

We consider the following objective:

$$W_k = \int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) P_k(\theta_1, \theta_2|\beta) d\theta_1 d\theta_2 \quad (2)$$

¹Any distribution can be handled by considering θ to be the *quantile* of the item rather than its absolute quality.



(a) Fix $g = 1$, with various objective function weights (b) Fix $w = 1$. Vary matching, $g = 1$, and $g = \frac{1+100}{11}$

Figure 1: Optimal β (with $M = 200$) with various objective weight functions w and matching rates g .

Weight function $w(\theta_1, \theta_2) > 0$ indicates how much the platform cares about not mistaking a quality θ_1 item with a quality θ_2 item. We consider scaled w such that $\int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) d\theta_1 d\theta_2 = 1$.

The central question we study is then the following: *What β yields the highest value of W_k ?* As noted in our introduction, the platform indirectly influences β through the design of its rating system. Our analysis of the optimal choice of β sets the benchmark the platform should try to achieve.

Discussion. Objective function. The specification (2) of the platform's objective is quite rich. It contains scaled versions of Kendall's τ (with $w(\theta_1, \theta_2) = 1$ for all θ_1, θ_2) and Spearman's ρ (with $w(\theta_1, \theta_2) = \theta_1 - \theta_2$) rank correlations as special cases. w allows the platform to encode, for example, that it cares more about correctly ranking just the very best, very worst, or items at both extremes.² Tarsitano [31] and da Costa and Roque [7] discuss other well-studied examples.

Relationship between model components. Qualitatively, β affects W_k as follows, as previewed in the introduction: when $\beta(\theta_1) \approx \beta(\theta_2)$, then $x_k(\theta_1) \approx x_k(\theta_2)$, and so $P_k(\theta_1, \theta_2 | \beta)$ is small (errors are common). A good design thus would have large $\beta(\theta_1) - \beta(\theta_2)$ for $\theta_1 > \theta_2$ where $w(\theta_1, \theta_2)$ is large.

Matching function g also affects P_k and thus W_k : when $g(\theta)$ is large, more ratings are sampled from item of quality θ , i.e. $n_k(\theta)$ is higher, and so $x_k(\theta)$ is more closely concentrated around its mean $\beta(\theta)$. Thus, $P_k(\theta, \theta' | \beta)$ increases (for all θ') with $g(\theta)$.

A good design of β thus considers both w and g .

Matching. As noted above, in our model we assume items receive an increasing number of ratings based on their true quality, through matching function $g(\theta)$. This is a reasonable approximation for our analysis, where we focus on the asymptotic rate of convergence of the ranking based on to the true ranking, as the number of ratings increases. In practice, items will be more likely to match when they have a higher *observed* aggregate score. Similarly, our model makes the stylized choice that all items have the same age. In reality, items have different ages in platforms.

3.2 Large deviations & discretization

Recall our central question: *What β yields the highest value of W_k ?* Here, we refine our objective W_k and constrain β to form a non-degenerate, feasible optimization task.

Large deviation rate function. As defined, W_k is not one objective: it has a different value per

²We use $\theta_1 \theta_2 (\theta_1 - \theta_2)$, $(1 - \theta_1)(1 - \theta_2)(\theta_1 - \theta_2)$, and $(\frac{1}{2} - \theta_1)^2 (\frac{1}{2} - \theta_2)^2 (\theta_1 - \theta_2)$ as examples.

time k , and no single choice of β can simultaneously optimize W_k for all k .³ Considering *asymptotic* performance is also insufficient: when β is strictly increasing in θ , $\lim_{k \rightarrow \infty} P_k(\theta_1, \theta_2 | \beta) = 1 \ \forall \theta_1, \theta_2$ by the law of large numbers. By bounded convergence, $\lim_{k \rightarrow \infty} W_k = 1$, and so *any* such β is asymptotically optimal.

For this reason, we consider maximization of the *rate* at which W_k converges, i.e., how *fast* the estimated item ranking converges to the true item ranking. We use a large deviations approach [9] to quantify this convergence rate. Formally, given a sequence $R_k \leq R$ with $\lim_{k \rightarrow \infty} R_k = R$, we refer to $-\lim_{k \rightarrow \infty} \frac{1}{k} \log(R - R_k) = c$ as the *large deviations rate* of convergence. If c exists then R_k approaches R exponentially fast, i.e., $R - R_k = e^{-kc+o(k)}$.

This discussion motivates the following optimization problem: *choose β to maximize the large deviations rate of convergence for W_k* , i.e., the rate at which the platform objective converges to its limit.

Discretizing β . Unfortunately, even this problem is degenerate if we consider continuous β : for any β that is not piecewise constant, the large deviations rate of convergence is zero, i.e., convergence of W_k to its limit is only polynomially fast, and characterizing the dependence of this convergence rate on β is intractable. Thus, the rate of convergence for W_k is not a satisfactory objective with continuous β .

We make progress by *discretizing* β ; in particular, we restrict attention to optimization over stepwise increasing β functions. Among stepwise increasing β , the large deviations rate of convergence to the limiting value of W_k can be shown to be nondegenerate, i.e. $\exists c, W > 0$ s.t. $W - W_k = e^{-kc+o(k)}$. (See Lemma C.4 in the Appendix for further discussion.)

Notationally, we will calculate an optimal stepwise increasing β with M levels, i.e. there are M intervals $S_i \subset [0, 1]$ and levels t_i such that when $\theta_1, \theta_2 \in S_i$, then $\beta(\theta_1) = \beta(\theta_2) \triangleq t_i$. The challenge is calculating an optimal $\mathbf{S}^* = \{S_i\}$ and $\mathbf{t}^* = \{t_i\}$.

The physical interpretation is that we group the items into M subsets (types) $S_i \subset [0, 1]$. When items θ_1, θ_2 are in the same subset, then their asymptotic reputation scores are the same, $\lim_k x_k(\theta_1) = \lim_k x_k(\theta_2) \triangleq t_i$. These items cannot be distinguished from one another even asymptotically.

Thus, though discretization allows us to define a large deviations rate for W_k , it comes at a cost: the limiting value of W_k is no longer one. Different discretization choices $\mathbf{S}$ result in different limiting values of W_k .

Note that, even for purely computational reasons, calculating β requires discretization.

Our optimization problem:

Within the class of stepwise increasing functions with M levels, find the β that is optimal, i.e. is

- (1) *Asymptotically optimal.* It yields the highest limiting value of W_k . AND
- (2) *Rate optimal.* It yields the fastest large deviations rate of convergence among *asymptotically optimal* β .

A remarkable result of our paper is a $\mathcal{O}\left(M \log^2 \frac{M}{\epsilon}\right)$ procedure to find an optimal β .

3.3 Solving the optimization problem

We now develop our procedure to find an optimal stepwise increasing β with M levels, i.e. M types of items.

³For example, consider a β such that the worst half of items never receive a positive rating and the rest always receive a positive rating. This function would perform comparatively well for a small number of ratings k , as it quickly distinguishes the best from the worst items. However, asymptotically, it would not be able to distinguish items within the same half. Some β' may lead to more mistakes initially but perform better at larger k .

Our first theorem below establishes that the problem decomposes into a two stage problem, the first for finding the optimal discretization intervals $\mathbf{S}^*$, and the second for the optimal levels $\mathbf{t}^*$ given $\mathbf{S}^*$.

Theorem 3.1. *The β defined by the following choices of $\mathbf{S}^*$ and $\mathbf{t}^*$ is optimal:*

$$\begin{aligned}\mathbf{S}^* &= \arg \max \sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) d(\theta_1, \theta_2) \\ \mathbf{t}^* &= \arg \max r(\mathbf{t}), \text{ where }^4 g_i \triangleq \inf_{\theta \in S_i} g(\theta) \text{ and}\end{aligned}$$

$$\begin{aligned}r(\mathbf{t}) &\triangleq - \lim_{k \rightarrow \infty} \frac{1}{k} \log W_k \\ &= \min_{0 \leq i \leq M-2} \inf_{a \in \mathbb{R}} \{g_{i+1} KL(a||t_{i+1}) + g_i KL(a||t_i)\}\end{aligned} \tag{3}$$

The proof is in the Appendix. The main hurdle is showing that the continuum of rates for $P_k(\theta_1, \theta_2)$ for each pair θ_1, θ_2 translates into a rate for W_k . This decomposition separates our two questions: $\mathbf{S}^*$ maximizes the limiting value of W_k *given any* $\mathbf{t}$, and depends only on w ; Then, $\mathbf{t}^*$ maximizes the rate at which the limiting value is reached, given g_i .

For Kendall's τ and Spearman's ρ , the optimal intervals are simply equispaced in $[0, 1]$, i.e. $S_i^* = [\frac{i}{M}, \frac{i+1}{M})$, because the entire item quality distribution is equally important. For other objective weight functions w , the difficulty of finding the optimal subsets $\mathbf{S}^*$ depends on the properties of w . Since $\mathbf{S}^*$ is trivial for Kendall's τ and Spearman's ρ – and w is just an analytic tool chosen by the platform to formalize its goals – we focus on finding the optimal levels $\mathbf{t}^*$ given g_i .

Discussion. One may naturally wonder why we introduced a continuum of quality $[0, 1]$ and then discretized into M subsets, instead of starting with M types. As established in Theorem 3.1, *how* we discretize (i.e. solving for $\mathbf{S}^*$) allows for optimization of different objective weight functions w ; it determines *which* items are most valuable to distinguish.

Suppose we started with a finite set of M items. Then the only remaining challenge is to equalize the rates at which we separate each item from its closest neighbor: the large deviations rate is unaffected by the weight function w in the objective, which is why w does not appear directly in the simplification of $r(\mathbf{t})$. In other words, *given* a discrete set of M items (equiv, given $\mathbf{S}^*$), calculating the optimal $\mathbf{t}$ is equivalent to solving a *maximin* problem for the rates at which each type is distinguished from each of the others. Thus, the algorithm discussed next also solves the *inverse* bandits problem in which there are M arms, we can choose the structure of the (binary) observations at each arm, and we wish to maximize the rate at which we correctly rank the arms, as measured by Kendall's τ or Spearman's ρ rank correlations. The solution corresponds to the Kendall's τ , i.e. $w = 1$, solution in Figure 1.

We further note that the choice of M is not consequential; in the Appendix Section B.4 we show that in an appropriate sense, a sequence of optimal β_M for each M converges as M gets large.

Algorithm to find the optimal levels We now describe how to find $\mathbf{t}^*$, the maximizer of (3).

The following lemma describes a system of equations to find the $\mathbf{t}^*$ that maximizes r . It states that $\mathbf{t}^*$ equalizes the rates at which each interval i is separated from its neighbors. The proof involves manipulation of (3) and convexity, and we provide it in the appendix.

⁴ $KL(a||b) = a \log \frac{b}{a} + (1-a) \log \frac{1-b}{1-a}$ is the Kullback-Leibler (KL) divergence between Bernoulli random variables with success probabilities a and b respectively.

Lemma 3.1. *The unique solution $\mathbf{t}^*$ to the following system of equations maximizes $r(\mathbf{t})$:*

$$\begin{aligned} r(t_0, t_1) &= r(t_1, t_2) = \dots = r(t_{M-2}, t_{M-1}) \\ t_0 &= 0, t_{M-1} = 1 \end{aligned} \tag{4}$$

$$\text{where } r(t_{i-1}, t_i) \triangleq -(g_{i-1} + g_i) \log \left[(1 - t_{i-1})^{\frac{g_{i-1}}{g_{i-1} + g_i}} (1 - t_i)^{\frac{g_i}{g_{i-1} + g_i}} + t_{i-1}^{\frac{g_{i-1}}{g_{i-1} + g_i}} t_i^{\frac{g_i}{g_{i-1} + g_i}} \right]$$

We do not know of any algorithm that efficiently and provably solves such convex equality systems in general. However, we leverage some structure in our setting to develop an algorithm, *NestedBisection*, with run-time and optimality guarantees. The efficiency of our algorithm results from the property that, given a rate, t_i is uniquely determined by the value of either of the adjacent levels t_{i-1}, t_{i+1} , reducing an exponentially large search space to an almost linear one. Physically, in other words, we only need to separate each type of item from its neighboring types.

Algorithm 1 contains pseudo-code, with a more detailed version available in Appendix Section B.2. Akin to branch and bound, the algorithm proceeds via bisection on the optimal value for the last unknown rating level, t_{M-2} . For each candidate value of t_{M-2} , the other levels can be found using a sequence of bisection subroutines. These candidate levels approximately obey all the equations in the system of equations (4) except the first one. The direction of the violation of the first equality reveals how to change the interval for the next outer bisection iteration.

Theorem 3.2. *NestedBisection finds an ϵ -optimal $\mathbf{t}$ in $\mathcal{O}(M \log^2 \frac{M}{\epsilon})$ operations, where ϵ -optimal means that $r(\mathbf{t})$ is within additive constant ϵ of optimal.*

We reserve the proof for the appendix. The main difficulty is finding a Lipschitz constant $\epsilon(\delta)$ for how much the rate (3) can change due to a shift of δ in one of the levels t_i . This requires lower bounding t_1 as a function of M . In practice, the algorithm runs instantaneously on a modern machine (e.g. for $M = 200$).

ALGORITHM 1: Nested Bisection

Data: Number of intervals M , match function g

Result: β levels, i.e. $\{t_0 \dots t_{M-1}\}$

Function *main* (M, δ, g)

```

    while Uncertainty region for  $t_{M-2}$  is bigger than error tolerance do
        Calculate  $r(t_{M-2}, 1)$ , the rate between current guess for  $t_{M-2}$ , and  $t_{M-1} = 1$ .
        Fixing  $t_{M-2}$ , find  $t_1 \dots t_{M-3}$  such that  $r(t_1, t_2) = r(t_2, t_3) = \dots = r(t_{M-3}, t_{M-2}) = r(t_{M-2}, 1)$ ,
        which can be done through a sequence of bisection routines.
        Calculate  $r(0, t_1)$ , the rate between current guess for  $t_1$ , and  $t_0 = 0$ .
        Compare  $r(t_{M-2}, 1)$  and  $r(0, t_1)$ , adjust uncertainty region for  $t_{M-2}$  accordingly.
    return  $\{t_i\}$ 

```

3.4 Visualization and discussion

Figure 1 shows how the optimal β (for $M = 200$) varies with weights w and matching rates g . As discussed informally above, higher relative weights in a region lead to a larger range of $\beta(\theta)$ in that region (e.g., w prioritizing the best items leads to a β shifted right from the others), making it easier to distinguish those items. Higher relative matching rates $g(\theta)$ have the opposite effect; when g is increasing, β shifts to better distinguish the worst items, as frequent sampling naturally increases accuracy for the best items. We formalize this shifting in Appendix Section B.3.

It's interesting to note that even the basic case, with $w = 1$ and $g = 1$, leads to a non-trivial β . One would expect that, with weight and matching functions that treat all items the same, that

β would be linear, i.e. $\beta(\theta) = \theta$. Instead, a third factor non-trivially impacts optimal design: binomial variance is highest near $\beta(\theta) = \frac{1}{2}$. Items who receive positive ratings at such frequency have high-variance scores, and thus the optimal β has a smaller mass of items with such scores.

4 Designing approximately optimal, implementable rating systems

We turn our attention to our second question: *How does a platform build and implement a real rating system such that buyers behave near-optimally, i.e. according to a calculated β ?* In this section, we give an example design procedure for how a platform would do so, and in the next section we validate our procedure through an experiment on Mechanical Turk.

Recall that $\beta(\theta)$ gives the probability at which an item of quality θ should receive a positive rating. However, the platform cannot force buyers to rate according to this function. Rather, it must ask *questions* of buyers that will induce a proportion $\beta(\theta)$ of them to give positive ratings for an item θ .

Throughout the section, we assume that an optimal $\beta(\theta)$ has been calculated (for some M , g , and w).

Resources available to platform. We suppose the platform has a set $\mathcal{Y}$ of possible binary questions that it could ask the buyer about the item, e.g., “Are you satisfied with your experience” or “Is this experience your best on our platform?”. Informally, at each rating opportunity (i.e., match made), the rater can be shown a single question $y \in \mathcal{Y}$. Let $\psi(\theta, y)$ be the probability an item quality θ would receive a positive rating when the rater is asked question $y \in \mathcal{Y}$.

We further suppose the platform has a set Θ of representative items for which it has access to item quality. Θ represents the level of granularity at which the platform can collect data about historical performance, or otherwise get expert labels. (We assume $M \gg |\Theta|$).

Using known set Θ , the platform can run an experiment to create an estimate $\hat{\psi}(\theta, y), \forall y \in \mathcal{Y}, \theta \in \Theta$.

Design heuristic. How can the platform build an effective rating system using these primitives and a calculated β ? We consider the following heuristic design: the platform randomly shows a question $y \in \mathcal{Y}$ to each buyer. The choice of the platform is a *distribution* $H(y)$ over $y \in \mathcal{Y}$; in other words, for the platform the design of the system amounts to choosing the frequency with which each question is shown. At each rating opportunity, y is chosen from $\mathcal{Y}$ according to H , independently across opportunities.

Clearly, the resulting probability that an item θ receives a positive rating is:


$$\tilde{\beta}(\theta) \triangleq \sum_{y \in \mathcal{Y}} \psi(\theta, y) H(y)$$

We would like to find a distribution H such that $\tilde{\beta}(\theta) = \beta(\theta)$ for all $\theta \in [0, 1]$, i.e., that the positive rating probability for each item is exactly the optimal value. However, there may not exist *any* set of questions $\mathcal{Y}$ with associated ψ and choice of H such that $\tilde{\beta} = \beta$.⁵

We propose the following heuristic to address this difficulty. Choose a probability distribution H to minimize the following L_1 distance:

$$\min_{H: \|H\|_1=1} \sum_{\theta \in \Theta} |\beta(\theta) - \sum_{y \in \mathcal{Y}} \hat{\psi}(\theta, y) H(y)| \quad (5)$$

⁵There are special cases where an exact solution exists. For example, let $\mathcal{Y} = [0, 1]$, and $\psi(\theta, y) = \mathbb{I}[\theta \geq y]$.



(a) Experimental $\hat{\psi}(\theta_i, y)$. Light blue lines are best 2 paragraphs, dark blue the worst. (b) β vs $\tilde{\beta}$ (using a calculated H) for $w = 1, g = 1$. (c) Simulated performance for $w = 1, g = 1$.

Figure 2: Experiment and simulation results

This heuristic uses the data available to the platform, $\hat{\psi}(\theta, y)$ for a set of items $\theta \in \Theta$, and designs H so at least these items receive ratings close to their optimal ratings $\beta(\theta_i)$. Then, as long as ψ is well-behaved, and Θ is representative of the full set, one can hope that $\tilde{\beta}(\theta) \approx \beta(\theta)$, for all $\theta \in [0, 1]$.

Discussion. *Real-world analogue and constraints.* A special case of this system is already in place on many on-line platforms, where a single question is shown to all users at all times (e.g. thumbs up/thumbs down for satisfaction). A static system can be designed by restricting H to only have mass at one question y . The procedure then selects the best static question to ask. More generally, any constraint (such as sparsity) can be included in the optimization (5).

Limitations. Our model does not allow for y to be chosen adaptively based on the platform’s current knowledge of the item. In practice, this may be a reasonable restriction for implementation purposes. Our model also restricts aggregation to be binary, i.e. the item receives a “win” for a positive answer and a “loss” for a negative answer, regardless of the question. In other words, the platform in our model does not use information on how “hard” a question y is.

5 Mechanical Turk experiment

In the previous sections, we showed how to characterize the optimal rating function β , and we described how to apply such a β to design a binary rating system using empirical data. In this section, we deploy an experiment on Amazon Mechanical Turk to apply these insights in practice. First, we collect data that can be used to create a reasonable real-world example of $\psi(\theta, y)$, as a proof of concept with which we can apply our optimization approach. Then, we use this model to demonstrate some key features of optimal and heuristic designs as computed via our methodology, and show that they perform well relative to natural benchmarks. Details of experimental design, simulation methodology, and results are in Appendix A.

Experiment description. We have a set of 10 English paragraphs of various (grammatical, writing) quality, along with *expert scores* θ . These paragraphs and expert scores are from a TOEFL book [10]; there were 5 unique possible experts ratings, i.e. $\theta \in \{.1, .3, .5, .7, .9\}$. For each possible rating, we have two paragraphs who received that score from experts.

We asked workers on Mechanical Turk to rate the writing quality of the paragraphs from a set of adjectives, $\mathcal{Y}$. Using this data, we estimate $\psi(\theta, y)$, i.e., the probability of positive rating when a question based on adjective $y \in \mathcal{Y}$ is shown for paragraph with quality $\theta \in \Theta$. (e.g., “Would you consider this paragraph of quality [adjective] or better?”) Figure 2a shows our estimated $\hat{\psi}$ for our 10 paragraphs.

Optimization. Next, we find the optimal β for various matching and weight functions using the methods from Section 3. In particular, we have β for all permutations of the cases $g = 1$, $g = \frac{1+10\theta}{11}$, and $w = 1$, $w = \theta_1\theta_2(\theta_1 - \theta_2)$, and $w = (1 - \theta_1)(1 - \theta_2)(\theta_1 - \theta_2)$. Recall that this step does not use data from the experiments.

Then, using $\hat{\psi}$ and calculated β s, we apply the heuristic from Section 4 to find the corresponding distributions H with which to sample the questions (adjectives) in $\mathcal{Y}$. Figure 2b shows the optimal β (with $g = 1$, $w = 1$), and estimated $\tilde{\beta}$ from the procedure.

Simulation. Finally, we study the performance of these designs via simulation in various settings. We simulate a system with 500 items and 100 buyers according to the model in Section 3.1, except that matching is *stochastic*: at each time, a random 100 items receive ratings, based on *observed* scores $x_k(\theta)$ rather than true quality θ . Furthermore, in some simulations, we have sellers *entering and exiting* the market with some probability at each time step. We measure the performance of all the designs. For comparison, we also simulate a *naive* $H = \frac{1}{|\mathcal{Y}|}$.

Note that our experiment only provided $\hat{\psi}$ associated with qualities $\theta \in \Theta$, and for our simulations we construct a full $\psi(\theta, y), \forall \theta \in [0, 1]$ from these points by averaging and interpolating (in order to model human behavior for the full system).

Further note that our *calibrated* simulations only provide rough evidence of the design approach working in real-life: Although we use real-world data, the simulations assume that our model reflects reality, except for where we deviate from the model as described above.

Results and discussion. Figure 2c shows the simulated performance (as measured by Kendall’s τ correlation) of the various designs over time, when $g = 1$. Further plots are in the Appendix Figure 6, showing performance under various weight functions w and matching functions g , and with items entering and exiting the market. We find that:

1. The optimal β for each setting outperforms other possible functions, as expected. The designs are robust to (some) assumptions in the model being broken, especially regarding matching.
2. The H produced through our procedure outperforms other possible designs, but is worse than the optimal system β one could design. In general, the simulated gap between an implemented system and optimal design β provides the platform quantitative insight on the system’s sub-optimality.
3. Comparing $\tilde{\beta}$ to β gives qualitative insight on *how* to improve one’s system. For example, in Figure 2b, $\tilde{\beta}$ is especially inaccurate for $\theta \in [0, .4]$. The platform must thus find better questions for items of such quality. $\hat{\psi}$ in Figure 2a corroborates: our question set fails to distinguish between two low quality paragraphs rated differently by experts (in dark blue and green, respectively).

We conclude by noting that a wide range of recent empirical work has documented that real-world rating systems experience inflation, with almost all items receiving positive ratings after almost every match [12, 13, 19, 30]. Our formulation helps understand how exactly – and whether – such inflation is suboptimal, along with providing guidance to platform designers. In particular, rating inflation can be interpreted as a current $\beta(\theta)$ that is very high for almost all item qualities θ . This system is well-performing if the platform objective is to just separate the worst items from the rest, or if high quality items receive many more ratings than low quality ones, but is clearly sub-optimal in other cases. Our paper provides a template for how a platform might address such a situation.

References

- [1] Daron Acemoglu, Ali Makhdoumi, Azarakhsh Malekian, and Asuman Ozdaglar. Fast and slow learning from reviews. Technical report, National Bureau of Economic Research, 2017.

- [2] Gary Bolton, Ben Greiner, and Axel Ockenfels. Engineering trust: reciprocity in the production of reputation information. *Management science*, 59(2):265–285, 2013.
- [3] Luís Cabral and Ali Hortaçsu. The Dynamics of Seller Reputation: Evidence from Ebay*. *The Journal of Industrial Economics*, 58(1):54–78, March 2010. ISSN 1467-6451. doi: 10.1111/j.1467-6451.2010.00405.x. URL <http://onlinelibrary.wiley.com/doi/10.1111/j.1467-6451.2010.00405.x/abstract>.
- [4] Yeon-Koo Che and Johannes Horner. Optimal design for social learning. 2015.
- [5] James Cook. Uber’s internal charts show how its driver-rating system actually works, February 2015. URL <http://www.businessinsider.com/leaked-charts-show-how-ubers-driver-rating-system-works-2015-2>.
- [6] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. John Wiley & Sons, November 2012. ISBN 978-1-118-58577-1.
- [7] J. Pinto da Costa and L. Roque. Limit distribution for the weighted rank correlation coefficient, rw. *REVSTAT-Statistical Journal*, 4(3), 2006.
- [8] Yuval Dagan, Yuval Filmus, Ariel Gabizon, and Shay Moran. Twenty (Simple) Questions. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2017, pages 9–21, New York, NY, USA, 2017. ACM. ISBN 978-1-4503-4528-6. doi: 10.1145/3055399.3055422. URL <http://doi.acm.org/10.1145/3055399.3055422>.
- [9] Amir Dembo and Ofer Zeitouni. *Large Deviations Techniques and Applications*, volume 38 of *Stochastic Modelling and Applied Probability*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2010. ISBN 978-3-642-03310-0 978-3-642-03311-7. URL <http://link.springer.com/10.1007/978-3-642-03311-7>. DOI: 10.1007/978-3-642-03311-7.
- [10] Educational Testing Service. TOEFL iBT Writing Sample Responses, 2005. URL http://toefl.uobabylon.edu.iq/papers/ibt_2014_12148630.pdf.
- [11] Paul Embrechts, Filip Lindskog, and Alexander Mcneil. Chapter 8 - Modelling Dependence with Copulas and Applications to Risk Management. In Svetlozar T. Rachev, editor, *Handbook of Heavy Tailed Distributions in Finance*, volume 1 of *Handbooks in Finance*, pages 329–384. North-Holland, Amsterdam, 2003. URL <https://www.sciencedirect.com/science/article/pii/B9780444508966500108>. DOI: 10.1016/B978-044450896-6.50010-8.
- [12] Apostolos Filippas, John J. Horton, and Joseph M. Golden. Reputation in the Long-Run. 2017.
- [13] A Fradkin, E Grewal, D Holtz, and M Pearson. Bias and reciprocity in online reviews: evidence from experiments on Airbnb. In *Proc. 16th ACM Conf. Electronic Commerce*, page 641, 2015.
- [14] Andrey Fradkin, Elena Grewal, and David Holtz. The Determinants of Online Review Informativeness: Evidence from Field Experiments on Airbnb. 2017. URL http://andreyfradkin.com/assets/reviews_paper.pdf.
- [15] Snehal Kumar (Neil) S. Gaikwad, Mark Whiting, Karolina Ziulkoski, Aipta Ballav, Aaron Gilbee, Senadhipathige S. Niranga, Vibhor Sehgal, Jasmine Lin, Leonardy Kristianto, Angela Richmond-Fuller, Jeff Regino, Durim Morina, Nalin Chhibber, Dinesh Majeti, Sachin Sharma, Kamila Mananova, Dinesh Dhakal, William Dai, Victoria Purynova, Samarth Sandeep, Varshine Chandrakanthan, Tejas Sarma, Adam Ginzberg, Sekandar Matin, Ahmed Nasser, Rohit Nistala, Alexander Stolzoff, Kristy Milland, Vinayak Mathur, Rajan Vaish, Michael S. Bernstein, Catherine Mullings, Shirish Goyal, Dilrukshi Gamage, Christopher Diemert, Mathias Burton, and Sharon Zhou. Boomerang: Rebounding the Consequences of Reputation Feedback on Crowdsourcing Platforms. pages 625–637. ACM Press, 2016. ISBN 978-1-4503-4189-9. doi: 10.1145/2984511.2984542. URL <http://dl.acm.org/citation.cfm?doid=2984511.2984542>.

- [16] Nikhil Garg and Ramesh Johari. Designing informative rating systems: Evidence from two experiments. 2018.
- [17] Peter Glynn and Sandeep Juneja. A large deviations perspective on ordinal optimization. In *Simulation Conference, 2004. Proceedings of the 2004 Winter*, volume 1. IEEE, 2004.
- [18] Fred Hicks, Lee Valentine, John Morrow, and Ian McDonald. Choosing Natural Adjective Ladders, 2000. URL http://www.mcdonald.me.uk/storytelling/lichert_article.htm.
- [19] John Horton and Joseph Golden. Reputation inflation: Evidence from an online labor market. *Work. Pap., NYU*, 2015.
- [20] Nan Hu, Paul A. Pavlou, and Jie (Jennifer) Zhang. Overcoming the J-Shaped Distribution of Product Reviews. SSRN Scholarly Paper ID 2369332, Social Science Research Network, Rochester, NY, October 2009. URL <https://papers.ssrn.com/abstract=2369332>.
- [21] Bar Ifrach, Costis Maglaras, Marco Scarsini, and Anna Zseleva. Bayesian Social Learning from Consumer Reviews. SSRN Scholarly Paper ID 2293158, Social Science Research Network, Rochester, NY, December 2017. URL <https://papers.ssrn.com/abstract=2293158>.
- [22] Nicole Immorlica, Brendan Lucier, Brian Rogers, and others. Emergence of cooperation in anonymous social networks through social capital. In *Proceedings of the 11th ACM Conference on Electronic Commerce*, 2010. URL <http://www.msr-waypoint.com/en-us/um/people/brlucier/misc/networkedPD-EC.pdf>.
- [23] Ramesh Johari, Vijay Kamble, and Yash Kanoria. Matching While Learning. *arXiv:1603.04549 [cs, stat]*, March 2016. URL <http://arxiv.org/abs/1603.04549>. arXiv: 1603.04549.
- [24] Sumeet Katariya, Branislav Kveton, Csaba Szepesvári, and Zheng Wen. DCM Bandits: Learning to Rank with Multiple Clicks. *arXiv:1602.03146 [cs, stat]*, February 2016. URL <http://arxiv.org/abs/1602.03146>. arXiv: 1602.03146.
- [25] Francis Maes, Louis Wehenkel, and Damien Ernst. Automatic Discovery of Ranking Formulas for Playing with Multi-armed Bandits. In *Recent Advances in Reinforcement Learning*, Lecture Notes in Computer Science, pages 5–17. Springer, Berlin, Heidelberg, September 2011. ISBN 978-3-642-29945-2 978-3-642-29946-9. doi: 10.1007/978-3-642-29946-9_5. URL https://link.springer.com/chapter/10.1007/978-3-642-29946-9_5.
- [26] Roger B. Nelsen. *An Introduction to Copulas*. Springer Science & Business Media, June 2007. ISBN 978-0-387-28678-5.
- [27] Chris Nosko and Steven Tadelis. The Limits of Reputation in Platform Markets: An Empirical Analysis and Field Experiment. Working Paper 20830, National Bureau of Economic Research, January 2015. URL <http://www.nber.org/papers/w20830>.
- [28] Filip Radlinski, Robert Kleinberg, and Thorsten Joachims. Learning diverse rankings with multi-armed bandits. In *Proceedings of the 25th international conference on Machine learning*, pages 784–791. ACM, 2008.
- [29] Shiva Rajaraman. Five Stars Dominate Ratings, September 2009. URL <https://youtube.googleblog.com/2009/09/five-stars-dominate-ratings.html>.
- [30] Steven Tadelis. Reputation and feedback systems in online platform markets. *Annual Review of Economics*, 8:321–340, 2016.
- [31] Agostino Tarsitano. Comparing the effectiveness of rank correlation statistics. *Working Papers, Università della Calabria, Dipartimento di Economia e Statistica*, pages 1–25, 2009.

- [32] Yisong Yue and Thorsten Joachims. Interactively optimizing information retrieval systems as a dueling bandits problem. In *Proceedings of the 26th Annual International Conference on Machine Learning*, pages 1201–1208. ACM, 2009.
- [33] Georgios Zervas, Davide Proserpio, and John Byers. A First Look at Online Reputation on Airbnb, Where Every Stay is Above Average. SSRN Scholarly Paper ID 2554500, Social Science Research Network, Rochester, NY, January 2015. URL <http://papers.ssrn.com/abstract=2554500>.

Appendix A Mechanical Turk experiment, simulations, and results

In this section, we expand upon the results discussed in Section 5. We design and run an experiment that a real platform may run to design a rating system. We follow the general framework in Section 4. We first run an experiment to estimate a $\psi(\theta, y)$, the probability at which each item with quality θ receives a positive answer under different questions y . Then, we design $H(y)$, using our optimal β for various settings (different objectives w and matching rates g). Then, we simulate several markets (using the various matching rates g) and measure the performance of the different rating system designs H , as measured by various objective functions (2).

A.1 Experiment description

We now describe our Mechanical Turk experiment. We ask subjects to rate the English proficiency of ten paragraphs. These paragraphs are modified TOEFL (Test of English as a Foreign Language) essays with known scores as determined by experts [10]. Subjects were given six answer choices, drawn randomly from the following list: *Abysmal*, *Awful*, *Bad*, *Poor*, *Mediocre*, *Fair*, *Good*, *Great*, *Excellent*, *Phenomenal*, following the recommendation of Hicks et al. [18]. *Poor* and *Good* are always chosen, and the other four are sampled uniformly at random for each worker. One paragraph is shown per page; returning to modify a previous answer is not allowed; and paragraphs are presented in a random order. This data is used to calibrate a model of ψ for optimization, i.e. to simulate a system with a set of questions $\mathcal{Y}$, where each question y corresponds to an adjective, “Would you characterize the performance of this item as [adjective] or better?”⁶

Different experiment trials are described below. Pilots were primarily used to garner feedback regarding the experiment from workers (fair pay, time needed to complete, website/UI comments, etc). All trials yield qualitatively similar results in terms of both paragraph ratings and feedback rating distributions for various scales.

Pilot 1 30 workers. Similar conditions as final experiment (6 words sampled for paragraph ratings, all uniformly at random, 5 point scale feedback rating), with identical question phrasing, “How does the following rate on English proficiency and argument coherence?”.

Pilot 2 30 workers. 7 words sampled for paragraph ratings, 6 point scale feedback rating, with the following question phrasing: “How does the following person rate on English proficiency and argument coherence?”.

Experiment 200 workers. 6 words sampled for paragraph ratings, with 2 fixed as described above, 5 point scale feedback rating. Question phrasing, “How does the following rate on English proficiency and argument coherence?”.

We use paragraphs modified from a set published by the Educational Testing Service [10]. There are 10 paragraphs, 5 each on 2 different topics. For each topic, the paragraphs have 5 distinct expert scores. Paragraphs are shortened to just a few sentences, and the top rated paragraphs are improved and the worst ones are made worse, preserving the ranking according to the expert scores.

Figure 3a shows time spent on each page of the experiment, Figure 3b shows the time spent per paragraph, and Figure 3c shows the cumulative density function for time spent by workers. The paragraphs are presented to workers in a random order. No workers are excluded in our data and all workers were paid \$1.00, including the ones that spent 2-3 seconds per page. 7/60 workers in the

⁶The data from the experiment is also used for a separate paper in submission, [16]. In that work, we analyze the full multi-option question directly, but the main focus is reporting the results of a separate, live trial on a large online labor platform.




Figure 3: Additional information for MTurk experiment




Figure 4: Paragraph rating distribution – for paragraph θ and rating word y , the empirical $\hat{\psi}(\theta, y)$ is shown. Colors encode the true quality as rated by experts (light blue is best quality, dark blue is worst).

pilots received a bonus of \$0.20 for providing feedback. The instructions advised workers to spend no more than a minute per question, though this was not enforced.

The instructions for the main experiment were as follows: “Please rate on English proficiency (grammar, spelling, sentence structure) and coherence of the argument, but not on whether you agree with the substance of the text.” No additional context was provided.

A.2 Calculating optimal β and H

Figure 4 shows the empirical $\hat{\psi}(\theta, y)$ as measured through our experiment. The colors encode the true quality as rated by experts (light blue is best quality, dark blue is worst); recall there are 10 paragraphs with 5 distinct expert ratings (paragraphs 0 and 5 are rated the best, paragraphs 4 and 9 are rated the worst).

With the β calculated and visualized using the methods in Section 3, we now find the optimal H for various settings using the methods in Section 4. We view our set of paragraphs as *representative items* Θ from a larger universe of paragraphs. In particular, we view our worst quality paragraphs as in the 10th percentile of paragraphs, and our best items as in the 90th percentile. In other words, from the empirical $\hat{\psi}$, we carry out the methods in Section 4 using a ψ s.t. $\psi(.1, y) = (\hat{\psi}(4, y) + \hat{\psi}(9, y))/2$ (and similarly for $\psi(.3, y), \psi(.5, y), \psi(.7, y), \psi(.9, y)$, where e.g. $\hat{\psi}(4, y)$ is the empirical rate at which paragraph 4 received a positive rating on question y).

Then, we solve the optimization problem for H stated in Section 4. From the above discussion,




Figure 5: Optimal $H(y)$ varying by $w(\theta_1, \theta_2)$ using Mechanical Turk data

we want to find an H such that the worst rated paragraphs in our experiment have a probability of receiving a positive rating that is approximately $\beta(.1)$.

Figure 5 shows the optimal H calculated for various platform settings. These distributions illustrate how often certain binary questions should be asked as it depends on the matching rates and platform objective. For example, as Figure 5a shows, when there is uniform matching and the platform cares about the entire ranking (i.e. has Kendall’s τ or Spearman’s ρ objective), it should ask most buyers to answer the question, “Would you rate this item as having ‘Fair’ quality or better?”.

Several qualitative insights can be drawn from the optimal H . Most importantly, note that the optimal designs vary significantly with the platform objective and matching rates. In other words, given the same empirical data $\hat{\psi}$, the platform’s design changes substantially based on its goals and how skewed matches are on the platform. Further, note that the differences in H follow from the differences in β that are illustrated in Figure 1: when the platform wants to accurately rank the best items, the questions that distinguish amongst the best (e.g., “Would you rate this item as having ‘Good’ quality?”) are drawn more often.

A.3 Simulation description

Using the above data and subsequent designs, we simulate markets with a binary rating system as described in Section 3.1. Our simulations have the following characteristics.

- 500 items. Items have i.i.d. quality in $[0, 1]$. For item with quality θ , we model buyer rating data using the ψ collected from the experiment as follows. In particular, we presume the items are convex combinations of the representative items in our experiment – items with quality $\theta \in [.1, .3]$ are assumed to have rating probabilities $\psi(\theta, y) = \alpha\psi(.1, y) + (1 - \alpha)\psi(.3, y)$, where $\alpha = (\theta - .1)/.2$. Similarly for θ in other intervals. This process yields the $\tilde{\beta}$ shown in Figure 2b.
- In some simulations, all items enter the market at time $k = 0$ and do not leave. In the others, with *entry and exit*, each item independently leaves the market with probability .02 at the end of each time period, and a new item with quality drawn i.i.d. from $[0, 1]$ enters.
- There are 100 buyers, each of which matches to an item independently. In other words, matching is independent across items, and items can match more than once per time period.

- Matching is random with probability as a function of an item’s *estimated* rank $\hat{\theta}$ according to score, rather than actual rank. In other words, the optimal systems were designed assuming item θ would match at rate $g(\theta)$; instead it matches according to $g(\hat{\theta})$, where $\hat{\theta}$ is the item’s rank according to score. We use both $g = 1$ and *linear* search, $g(\hat{\theta}) = \frac{1+10\hat{\theta}}{11}$.
- $\mathcal{Y}$ is the set of 9 adjectives from our MTurk experiments.
- We test several possible H : *naive* with $H(y) = \frac{1}{|\mathcal{Y}|}$, and then the various optimal H calculated for the different sections, illustrated in Figure 5.

A.3.1 Simulation results

Figure 6 contains plots from a simulated system that has binary ratings. Figures 6a, 6b are with uniform search ($g = 1$), Figures 6c and 6d plot the objective prioritizing the worst items, and Figures 6e and 6f are with linearly increasing search. For each setting, we include both plots with and without birth/death.

Together, the results suggest that the asymptotic and rate-wise optimality of our calculated β hold even under deviations of the model, and that the real-world design approach outlined in Section 4 would provide substantial information benefits to platforms.

Several specific qualitative insights can be drawn from the figures, alongside those discussed in the main text.

1. From all the plots with uniform search, the H designed using our methods for the given setting outperforms other H designs, as expected, and the optimal β (for the given setting) significantly outperforms other designs both *asymptotically* and *rate-wise*.
2. Qualitatively, again with uniform search, heterogeneous item age also does not affect the results. In fact, it seems as if the optimal β and best possible H (given the data) as calculated from our methods outperforms other designs both *asymptotically* and *rate-wise*. Note that this is true even though items entering and leaving the market means that the system may not enter the asymptotics under which our theoretical results hold.
3. Figures 6c and 6d show the same system parameters as Figures 6a, 6b, i.e. uniform search. However, while 6a, 6b show Kendall’s τ correlation over time, 6c and 6d show the objective prioritizing bottom items ($w = (1 - \theta_1)(1 - \theta_2)(\theta_1 - \theta_2)$). Note that the β calculated for the actual objective outperforms that calculated for Kendall’s τ , including asymptotically.


Similarly, complementing the fact that H design changes significantly with the weight function, these plots show the value of designing while taking into account one’s true objective value – the different designs perform differently. Mis-specifying one’s objective (e.g. designing to differentiate the best items when one truly cares about the worst items) leads to a large gap in performance (e.g. see the gap between the dark green and red lines in 6c and 6d).

Note that comparing the performance of β for the misspecified objective and H for the true objective is not a fair comparison: the former differentiates between all items (though potentially not in a rate-optimal way), while H is constrained by reality, i.e. ψ and $\mathcal{Y}$.


4. Now, consider Figures 6e and 6f, which plot the system with linearly increasing search. Note that, contrary to expectation, the optimal β for *uniform* search outperforms the β for the actual system simulated, with linear search! This pattern is especially true for small time k and with item birth/death.

This inversion can be explained as follows. *Uniformization* occurs with heterogeneous age and matching according to observed quality: new items of high type are likely to be mis-ranked lower, while new items of low type are more likely to be mis-ranked higher. (We note that this may not matter in practice, where the search function itself is fit through data, which already captures this effect.) These errors are prominent at low time k and with item birth/death, i.e., in the latter our system never reaches the asymptotics at which the linear β is the optimal design.


This pattern can be seen more clearly by comparing the two β curves in Figures 6e, without item birth/death. At small k , when errors are common and so search is more effectively uniform, the β for uniform matching performs the best. However, as such errors subside over time, the performance of the β for linear search catches up and eventually surpasses that of uniform optimal β .




(a) Uniform matching, no birth/death




(b) Uniform matching, 2% probability of death per time




(c) Uniform matching, no birth/death, worst items weighted objective



(d) Uniform matching, 2% birth/death, worst items weighted objective



(e) Linear matching, no birth/death



(f) Linear matching, 2% probability of death per time

Figure 6: Simulations from data from Mechanical Turk experiment – Binary rating system

Appendix B Supplementary theoretical information and results

We now give some additional detail and develop additional results. Section B.1 contains the formal specification and update of our deterministic dynamical system. Section B.2 gives our algorithm, Nested Bisection, in far more detailed pseudo-code. Section B.3 formalizes our earlier qualitative discussion on how matching rates affects the function β . Section B.4 includes a convergence result for functions β_M as M increases. Finally, Section B.5 contains simple results on how one can learn $\psi(\theta, y)$ through experiments, even if one does not have a reference set of items Θ with known quality before one begins experiments.

B.1 Formal specification of system state update

Recall that $\mu_k(\Theta, X)$ is the mass of items with true quality $\theta \in \Theta \subseteq [0, 1]$ and a reputation score $x \in X \subseteq [0, 1]$ at time k . Let $E_k = \{\theta : n_k(\theta) = n_{k-1}(\theta) + 1\}$. These are the items who receive an additional rating at time k ; for all $\theta \in E_k^c$, $n_k(\theta) = n_{k-1}(\theta)$. Our system is completely deterministic, and evolves according to the distributions of the individual seller dynamics.

For each $\theta \in E_k, x, x'$, define $\omega(\theta, x, x')$ as follows:

$$\omega(\theta, x, x') = \beta(\theta)\mathbb{I}\{n_k(\theta)x - n_{k-1}(\theta)x' = 1\} + (1 - \beta(\theta))\mathbb{I}\{n_k(\theta)x - n_{k-1}(\theta)x' = 0\}.$$

Then ω gives the probability of transition from x' to x when an item receives a rating. We then have:

$$\mu_{k+1}(\Theta, X) = \int_{E_k} \int_{x'=0}^1 \int_{x \in X} \omega(\theta, x, x') dx \mu_k(dx', d\theta) + \int_{E_k^c} \int_{x \in X} \mu_k(dx, d\theta).$$

It is straightforward but tedious to check that the preceding dynamics are well defined, given our primitives.

B.2 Detailed algorithm

Here, we present the Nested Bisection algorithm, which is described at a high level and summarized in pseudo-code in the main text, in more detail.

ALGORITHM 2: Nested Bisection given in more detail

Data: Set size M , grid width δ , match function g */* Assume $\delta \ll \min_i t_i - t_{i-1}$ */*

Result: β_M levels $\{t_0 \dots t_{M-1}\}$

Function *main* (M, δ, g)

```

     $t_0 = 0, t_{M-1} = 1$ 
     $\ell = 1 - \frac{1}{M-1}, u = 1 - \delta$ 
    while  $u - \ell > \delta/2$  do
         $j_{M-2} = \frac{r+\ell}{2}$ 
         $\text{rate}_{\text{last}} = -g_{M-2} \log(t_{M-2})$ 
         $\{j_1 \dots j_{M-3}\} = \text{CalculateOtherLevels}(j_{M-2}, \text{rate}_{\text{last}}, g)$ 
         $\text{rate}_{\text{first}} = -g_1 \log(1 - t_1)$ 
        if  $\text{rate}_{\text{first}} < \text{rate}_{\text{last}}$  then  $\ell = j_{M-2}$ 
        else  $u = j_{M-2}$ 
     $\{t_1 \dots t_{M-2}\} = \text{CalculateOtherLevels}(u, g)$ 
     $t_{M-2} = u$ 
    return  $\{t_i\}$ 

```

Function *PairwiseRate* ($t_{m-1}, t_m, g_m, g_{m-1}$)

```

    return  $-(g_{m-1} + g_m) \log \left[ (1 - t_{m-1})^{\frac{g_{m-1}}{g_{m-1} + g_m}} (1 - t_m)^{\frac{g_m}{g_{m-1} + g_m}} + t_{m-1}^{\frac{g_{m-1}}{g_{m-1} + g_m}} t_m^{\frac{g_m}{g_{m-1} + g_m}} \right]$ 

```

Function *CalculateOtherLevels* ($j_{M-2}, \text{rate}_{\text{target}}, g$)

```

    /* Given target rate from current guess  $j_{M-2}$ , sequentially fix other levels. */
    foreach  $m \in M - 3 \dots 1$  do
         $j_m = \text{BisectNextLevel}(j_{m+1}, \text{rate}_{\text{target}}, g_m, g_{m+1})$ 
    return  $\{j_1 \dots j_{M-3}\}$ 

```

Function *BisectNextLevel* ($j_m, \text{rate}_{\text{target}}, g_{m-1}, g_m$)

```

     $\ell = 0, r = j_m - \delta$ 
    while  $r - \ell > \delta/2$  do
         $j_{m-1} = \frac{r+\ell}{2}$ 
         $\text{rate}_m = \text{PairwiseRate}(j_{m-1}, j_m, g_{m-1}, g_m)$ 
        if  $\text{rate}_m \leq \text{rate}_{\text{target}}$  then  $r = j_{m-1}$ 
        else  $\ell = j_{m-1}$ 
    return  $r$ 

```

B.3 Formalization of effect of matching rates shifting

Matching concentrating at the top items moves mass of $\beta(\theta)$ away from high θ , and subsequently mass of $H(y)$ away from the questions that help distinguish the top items, as observed in Figures 1b and 5b above. Informally, this occurs because when matching concentrates, top items are accumulating many ratings more ratings comparatively, and so the amount of information needed per rating is comparatively less. We formalize this intuition in Lemma B.1 below.

The lemma states that if matching rates shift such that there is an index k above which matching rates increase and below which they decrease, then correspondingly the levels of β , (i.e. t_i) become closer together above k .

Lemma B.1. *Suppose $k, g, \tilde{g}$ such that $\forall j \in \{k+1 \dots M-1\}, g_j \geq \tilde{g}_j$, and $\forall j \in \{0 \dots k-1\}, g_j \leq \tilde{g}_j$, and $g_k = \tilde{g}_k$. Then, $t_k^* \geq \tilde{t}_k^*$.*

Proof. This proof is similar to that of Lemma 3.1, except that with the matching function changing the overall rate function can either increase, decrease, or stay the same. Suppose the overall rate function decreased or stayed the same when the matching function changed from $\tilde{g}$ to g . Then $g_{M-2} > \tilde{g}_{M-2}$ and the target rate is no larger, and so $t_{M-2}^* > \tilde{t}_{M-2}^*$. Then, $t_{M-3}^* > \tilde{t}_{M-3}^*$ (a smaller width is needed because the matching rates are higher and the rate is no larger, and the next value also increased). This shifting continues until $t_{k+1}^* > \tilde{t}_{k+1}^*$. Then, $t_k^* > \tilde{t}_k^*$.

Suppose instead that the overall rate function increased when the matching function changed from $\tilde{g}$ to g . Then $g_1 < \tilde{g}_1$ and the target rate is larger, and so $t_1^* > \tilde{t}_1^*$. Then, $t_2^* > \tilde{t}_2^*$ (a larger width is needed and the previous value also increased). This shifting continues until $t_{k-1}^* > \tilde{t}_{k-1}^*$. Then, $t_k^* > \tilde{t}_k^*$. $\square$

B.4 Limit of β as $M \rightarrow \infty$

Let β_M^w denote the optimal β with M intervals for weight function w , with intervals $\{S_i^{wM}\} = \{[s_i^{wM}, s_{i+1}^{wM})\}$ and levels $\mathbf{t}^{wM}$. Let $q_{wM}(\theta) = i/M$ when $\theta \in [s_i^{wM}, s_{i+1}^{wM})$, i.e. the quantile of interval item of type θ is in.

Then, we have the following convergence result for β_M .

Theorem B.1. *Let g be uniform. Suppose w such that q_{wM} converges uniformly. Then, $\forall C \in \mathbb{N}, \exists \beta^w$ s.t. $\beta_{C2N+1}^w \rightarrow \beta^w$ uniformly as $N \rightarrow \infty$.*

The proof is technical and is below. We leverage the fact that, for g uniform, the levels of β_{2M} can be analytically written as a function of the levels of β_M . We believe (numerically observe) that this theorem holds for the entire sequence as opposed to the each such subsequence, and for general matching functions g . However, our proof technique does not carry over, and the proof would leverage more global properties of the optimal β_M .

Furthermore, the condition on w is light. For example, it holds for Kendall's τ , Spearman's ρ , and all other examples mentioned in this work.

This convergence result suggests that the choice of M when calculating a asymptotic and rate optimal β is not consequential. As M increases, the limiting value of W_k increases to 1 (i.e. the asymptotic value increases), but the optimal rate decreases to 0. As discussed above, with strictly increasing and continuous β , the asymptotic value is 1 but the large deviations rate does not exist, i.e. convergence is polynomial.

This result could potentially be strengthened as follows: first, show convergence on the entire sequence as opposed to these exponential subsequences, as conjectured; second, show desirable properties of the limiting function itself. It is conceivable but not necessarily true that the limiting

function is “better” than other strictly continuous increasing functions in some rate sense, even though the comparison through large deviations rate is degenerate.

B.5 Learning $\psi(\theta, y)$ through experiments

Now, we show how a platform would run an experiment to decide to learn $\psi(\theta, y)$. In particular, one potential issue is that the platform does not have any items with known quality that it can use as representative items in its optimization. In this case, we show that it can use ratings within the experiment itself to identify these representative items. The results essentially follow from the law of large numbers.

We assume that $|\Theta| = L$ representative items $i \in \{1 \dots L\}$ are in the experiment, and each are matched N times. The experiment proceeds as follows: every time an item is matched, show the buyer a random question from $\mathcal{Y}$. For each word $y \in \mathcal{Y}$, track the empirical $\hat{\psi}(i, y)$, the proportion of times a positive response was given to question y . Alternatively, if $\mathcal{Y}$ is totally ordered (i.e. a positive rating for a given y also implies positive ratings would be given to all “easier” y'), and can be phrased as a multiple choice question, data collection can be faster: e.g., as we do in our experiments: $\mathcal{Y}$ consists of a set of totally ordered adjectives that can describe the item; the rater is asked to pick an adjective out of the set; this is interpreted as the item receiving a positive answer to the questions induced by the chosen answer and all worse adjectives, and a negative answer to all better adjectives.

First, suppose the platform approximately knows the quality θ_i of each item i , and θ_i are evenly distributed in $[0, 1]$. Suppose the items are ordered by index, i.e. $\theta_1 < \theta_2 < \dots < \theta_L$. Then let $\hat{\psi}(\theta, y) = \hat{\psi}(i, y)$ when $\theta \in [\theta_{i-1}, \theta_i]$. Call this procedure *KnownTypeExperiment*.

Lemma B.2. *Suppose $\psi(\theta, y)$ is Lipschitz continuous in θ . With *KnownTypeExperiment*, $\hat{\psi}(i, y) \rightarrow \psi(\theta_i, y) \forall y$ uniformly as $N \rightarrow \infty$. As $L \rightarrow \infty$, $\hat{\psi}(\theta, y) \rightarrow \psi(\theta, y) \forall \theta$ uniformly.*

Proof. The proof follows directly from the Strong Law of Large Numbers. As $N \rightarrow \infty$, $\forall i$, $\hat{\psi}(i, x) \rightarrow \psi(\theta_i, x)$ uniformly. Now, let $L \rightarrow \infty$. $\forall \epsilon, \exists L' \text{ s.t. } \forall L > L', \forall \theta, \exists i \text{ s.t. } |\theta - \theta_i| < \epsilon$. $\psi(\theta, x)$ is Lipschitz in θ by assumption, and so $\hat{\psi}(\theta, x) \rightarrow \psi(\theta, x)$ uniformly. $\square$

We now relax the assumption that the platform has an existing set of items with known qualities. Suppose instead the platform has many items L of unknown quality who are expected to match N times each over the experiment time period. For each item, the platform would again ask questions from $\mathcal{Y}$, drawn according to any distribution (with positive mass on each question). Then generate $\hat{\psi}(\theta, y)$ as follows: first, rank the items according to their ratings during the experiment itself. Then, for each y , $\hat{\psi}(\theta, y)$ is the empirical performance of the θ th percentile item in the ranking, i.e. $\hat{\psi}(\theta, y) = \hat{\psi}(\theta_i, y)$ for $\theta \in [\frac{i-1}{L}, \frac{i}{L}]$. Call this procedure *UnknownTypeExperiment*.

Lemma B.3. *Suppose $\psi(\theta, y)$ is Lipschitz continuous in θ . With *UnknownTypeExperiment*, $\hat{\psi}(\theta, y) \rightarrow \psi(\theta, y) \forall y, \theta$ uniformly as $L, N \rightarrow \infty$.*

Proof. Fix L . Denote each item in the experiment as $i \in \{1 \dots L\}$ (with true quality $\theta_i \neq \theta_j$), and each item has N samples. Without loss of generality, assume the items are indexed according to their rank on the average of their scores on the samples, defined as the percentage of positive ratings received. $i = 1$ is then the worst item, and $i = L$ is the best item according to scores in the experiment.

For $\psi(\theta, x)$ increasing in θ , as $N \rightarrow \infty$, $Pr(\theta_i > \theta_j | i < j) \rightarrow 0$ almost surely by SLLN, and for a fixed L , $\{\theta_i\}$ this convergence is uniform. Furthermore, by SLLN, $\hat{\psi}(i, x) \rightarrow \psi(\theta_i, x)$ as $N \rightarrow \infty$. Recall $\hat{\psi}(\theta, x) = \hat{\psi}(i, x)$ for $\theta \in [\frac{i-1}{L}, \frac{i}{L}]$.

Now, let $L \rightarrow \infty$. $\forall \epsilon, \exists L'$ s.t. $\forall L > L', \forall \theta, \exists i$ s.t. $|\theta - \theta_i| < \epsilon$. $\psi(\theta, x)$ is Lipschitz in θ by assumption, and so $\hat{\psi}(\theta, x) \rightarrow \psi(\theta, x)$ uniformly. $\square$

Appendix C Proofs

In this Appendix section, we prove our results.

Sections C.1-C.3 develop rate functions for P_k and W_k . While rates for P_k follow immediately from large deviation results, the rate function for W_k requires more effort as the quantity is an integral over a continuum of (θ_1, θ_2) , each of which has a rate corresponding to that of $P_k(\theta_1, \theta_2)$.

Then in Section C.4 we prove Theorem 3.1 and Lemma 3.1.

Section C.5 then contains additional necessary lemmas required for the proof of the algorithm and convergence result, Theorem B.1. The main difficulty for the former is showing a Lipschitz constant in the resulting rate if a level t_i is shifted, which requires lower and upper bounds for t_1 and t_{M-2} , respectively. For the former, we need to relate the solutions of the sequence of optimization problems used to find β_M as M increases. It turns out that both properties follow by relating the levels of β_M to those of β_{2M-1} .

These additional lemmas are used to prove the algorithm approximation bound (Theorem 3.2) and the convergence result (Theorem B.1) in Section C.6 and C.7, respectively.

Finally in Section C.8 we prove the comments we make in the main text about Kendall's τ and Spearman's ρ rank correlations belonging in our class of objective functions, with asymptotic values of W_k maximized when $\mathbf{s}$ is equispaced in $[0, 1]$.

C.1 Rate functions for $P_k(\theta_1, \theta_2)$

Lemma C.1.

$$\lim_{k \rightarrow \infty} -\frac{1}{k} \log [\mu((x_k(\theta_1) - x_k(\theta_2)) \leq 0 | \theta_1, \theta_2)] = \inf_{a \in \mathbb{R}} \{g(\theta_1)I(a|\theta_1) + g(\theta_2)I(a|\theta_2)\}$$

where $I(a|\ell) = \sup_z \{za - \Lambda(z|\theta)\}$, and $\Lambda(z|\theta)$ is the log moment generating function of a single sample from $x(\theta_1)$ and $g(\theta)$ is the sampling rate.

Proof. $\lim_{k \rightarrow \infty} -\frac{1}{k} \log [\mu((x_k(\theta_1) - x_k(\theta_2)) \leq 0 | \theta_1, \theta_2)]$

$$= \lim_{k \rightarrow \infty} -\frac{1}{k} \log \left[\int_{a \in \mathbb{R}} \mu((x_k(\theta_1) = a | \theta_1) \mu(x_k(\theta_2) \geq a | \theta_2) da \right] \quad (6)$$

$$= \lim_{k \rightarrow \infty} -\frac{1}{k} \log \left[\int_{a \in \mathbb{R}} e^{-kg(\theta_1)I(a|\theta_1)} e^{-kg(\theta_2)I(a|\theta_2)} da \right] \quad (7)$$

$$= \inf_{a \in \mathbb{R}} \{g(\theta_1)I(a|\theta_1) + g(\theta_2)I(a|\theta_2)\} \quad \text{Laplace principle} \quad (8)$$

$\square$

Where (7) is a basic result from large deviations, where $kg(\theta_i)$ is the number of samples item of quality θ_i has received.

Note that this lemma also appears in Glynn and Juneja [17], which uses the Gartner-Ellis Theorem in the proof. Our proof is conceptually similar but instead uses Laplace's principle.

Recall that $\text{KL}(a||b) = a \log \frac{b}{a} + (1-a) \log \frac{1-b}{1-a}$ is the Kullback-Leibler (KL) divergence between Bernoulli random variables with success probabilities a and b respectively. It is well known that for a Bernoulli random variable with success probability t ,

$$I(a|t) = \text{KL}(a||t)$$

Then, we have

Lemma C.2. *Let $\theta_1 > \theta_2$ and $I(a|\theta) = KL(a||\beta(\theta))$. Further, Let $\bar{P}_k(\theta_1, \theta_2) = 1 - P_k(\theta_1, \theta_2)$. Then,*

$$-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{P}_k(\theta_1, \theta_2) = \inf_{a \in \mathbb{R}} \{g(\theta_1)I(a|\theta_1) + g(\theta_2)I(a|\theta_2)\}, \quad (9)$$

Proof. Follows directly from Lemma C.1.

$$\begin{aligned} & -\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{P}_k(\theta_1, \theta_2 | \beta) \\ &= \lim_{k \rightarrow \infty} -\frac{1}{k} \log [1 + \mu_k(x_k(\theta_1) - x_k(\theta_2) < 0 | \theta_1, \theta_2) - \mu_k(x_k(\theta_1) - x_k(\theta_2) > 0 | \theta_1, \theta_2)] \\ &= \lim_{k \rightarrow \infty} -\frac{1}{k} \log [2\mu_k(x_k(\theta_1) - x_k(\theta_2) < 0 | \theta_1, \theta_2) + \mu_k(x_k(\theta_1) - x_k(\theta_2) = 0 | \theta_1, \theta_2)] \\ &= \lim_{k \rightarrow \infty} -\frac{1}{k} \log [\mu_k(x_k(\theta_1) - x_k(\theta_2) \leq 0 | \theta_1, \theta_2)] \\ &= \inf_{a \in \mathbb{R}} \{g(\theta_1)I(a|\theta_1) + g(\theta_2)I(a|\theta_2)\} \end{aligned} \quad \text{Lemma C.1}$$

□

C.2 Laplace's principle with sequence of rate functions

In order to derive a rate function for $\bar{W}_k = (\lim_k W_k) - W_k$, we need to be able to relate its rate to that of $\bar{P}_k(\theta_1, \theta_2)$. The following theorem, related to Laplace's principle for large deviations allows us to do so.

Theorem C.1. *Suppose that X is compact with finite Lebesgue measure $\mu(X) < \infty$. Suppose that $\varphi(x)$ has an essential infimum $\underline{\varphi}$ on X , that $\varphi_n(x)$ has an essential infimum $\underline{\varphi}_n$, that both φ and all φ_n are nonnegative, and that $\varphi_n \rightarrow \varphi$ uniformly:*

$$\lim_{n \rightarrow \infty} \sup_{x \in X} |\varphi_n(x) - \varphi(x)| = 0.$$

Then:

$$\lim_{n \rightarrow \infty} \frac{1}{n} \log \int_X e^{-n\varphi_n(x)} dx = -\underline{\varphi}. \quad (10)$$

Proof. First, we note that for all x and n , $e^{-n\varphi_n(x)} \leq e^{-n\underline{\varphi}_n}$. Therefore, letting $(*)$ denote the LHS of (10), we have:

$$(*) \leq \lim_{n \rightarrow \infty} \frac{1}{n} \log \int_X e^{-n\underline{\varphi}_n} dx = -\underline{\varphi},$$

where the last limit follows from the fact that φ_n converges uniformly to φ , so that $\underline{\varphi}_n \rightarrow \underline{\varphi}$.

Next, for $\epsilon > 0$ let $A_n(\epsilon) = \{x : \varphi_n(x) \leq \underline{\varphi}_n + \epsilon\}$ and let $A(\epsilon) = \{x : \varphi(x) \leq \underline{\varphi} + \epsilon\}$. It follows (again by uniform convergence) that for all sufficiently large n , $A(\epsilon/2) \subseteq A_n(\epsilon)$, so that $\mu(A(\epsilon/2)) \leq \mu(A_n(\epsilon))$ for all sufficiently large n . Further, $\mu(A(\epsilon/2)) > 0$, since $\underline{\varphi}$ is the essential infimum.

Since:

$$\int_X e^{-n\varphi_n(x)} dx \geq \mu(A_n(\epsilon)) e^{-n(\underline{\varphi}_n + \epsilon)},$$

it follows that:

$$(*) \geq -\underline{\varphi} - \epsilon + \lim_{n \rightarrow \infty} \frac{1}{n} \log \mu(A_n(\epsilon)).$$

To complete the proof, observe that since $\mu(A_n(\epsilon))$ is bounded below by a positive constant for all sufficiently large n , the last limit is zero. Therefore:

$$(*) \geq -\underline{\varphi} - \epsilon.$$

Since ϵ was arbitrary, this completes the proof. $\square$

Remark C.1. Let $X = [0, 1] \times [0, 1]$, $\varphi_n(\theta_1, \theta_2) = -\frac{1}{n} \log \bar{P}_n(\theta_1, \theta_2)$. Then, all the conditions for Theorem C.1 are met.

C.3 Rate function for W_k

Our next lemma shows that we can obtain a nontrivial large deviations rate for W_k when β is a step-wise increasing function.

Recall $W_k = \int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) P_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2)$.

Let $\bar{P}_k(\theta_1, \theta_2) = 1 - P_k(\theta_1, \theta_2)$.

Further, let $\bar{W}_k = (\lim_k W_k) - W_k = \int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) \bar{P}_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2)$. (recall we assumed w integrates to 1 without loss of generality).

Lemma C.3. Suppose β is piecewise constant with M levels $\{t_i\}$. Let $g_i \triangleq \inf_{\theta \in S_i} g(\theta) = g(s_i)$. Then,

$$-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{W}_k = \min_{0 \leq i \leq M-2} \inf_{a \in \mathbb{R}} \{g_{i+1} I(a | t_{i+1}) + g_i I(a | t_i)\} \triangleq r, \quad (11)$$

where $I(a | t) = KL(a || t)$ as defined in Lemma C.2.

Proof. When β is step-wise increasing with M levels $\{t_i\}$, then

$$\bar{W}_k = \sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) \bar{P}_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2)$$

as $\bar{P}_k(\theta_1, \theta_2) = 0$ when $\beta(\theta_1) = \beta(\theta_2)$.

$$-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{W}_k$$

$$= -\lim_{k \rightarrow \infty} \frac{1}{k} \log \int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) \bar{P}_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2) \quad (12)$$

$$= -\lim_{k \rightarrow \infty} \frac{1}{k} \log \sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) \bar{P}_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2) \quad (13)$$

$$= -\max_{0 \leq i < j < M} \left(\lim_{k \rightarrow \infty} \frac{1}{k} \left[\log \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) \bar{P}_k(\theta_j, \theta_i | \beta) d(\theta_1, \theta_2) \right] \right) \quad (14)$$

$$= -\max_{0 \leq i < j < M} \sup_{\theta_1 \in S_j, \theta_2 \in S_i} \left(\lim_{k \rightarrow \infty} \frac{1}{k} [\log w(\theta_1, \theta_2) \bar{P}_k(\theta_j, \theta_i | \beta)] \right) \quad (15)$$

$$= -\max_{0 \leq i < j < M} \sup_{\theta_1 \in S_j, \theta_2 \in S_i} \left(\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{P}_k(\theta_j, \theta_i | \beta) \right) \quad (16)$$

$$= \min_{0 \leq i < j < M} \inf_{\theta_1 \in S_j, \theta_2 \in S_i} \left(-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{P}_k(\theta_j, \theta_i | \beta) \right) \quad (17)$$

$$= \min_{0 \leq i < j < M} \inf_{\theta_1 \in S_j, \theta_2 \in S_i} \inf_{a \in \mathbb{R}} \{g(\theta_1)I(a|t_j) + g(\theta_2)I(a|t_i)\} \quad (18)$$

$$= \min_{0 \leq i < j < M} \inf_{a \in \mathbb{R}} \{g_j I(a|t_j) + g_i I(a|t_i)\} \quad (19)$$

$$= \min_{0 \leq i < M-1} \inf_{a \in \mathbb{R}} \{g_{i+1} I(a|t_{i+1}) + g_i I(a|t_i)\} \quad (20)$$

The last line follows from adjacent t_i, t_{i+1} dominating the rate due to monotonicity properties.

Line (15) follows from Theorem C.1.

Line (14) follows from: $\forall a_i^\epsilon \geq 0$, $\limsup_{\epsilon \rightarrow 0} \left[\epsilon \log \left(\sum_i^N a_i^\epsilon \right) \right] = \max_i^N \limsup_{\epsilon \rightarrow 0} \epsilon \log(a_i^\epsilon)$, which is a finite case version (with fewer assumptions) of Theorem C.1. See, e.g., Lemma 1.2.15 in [9] for a proof of this property. $\square$

Lemma C.4. $\beta(\theta)$ is piecewise constant $\iff \exists c(\beta) > 0$ s.t. $-\lim_{k \rightarrow \infty} \frac{1}{k} \log(\bar{W}_k) = c(\beta)$.

Proof. $\implies$ follows directly from Lemma C.3: $\inf_{a \in \mathbb{R}} \{g_{i+1} I(a|t_{i+1}) + g_i I(a|t_i)\} > 0$ when $t_i \neq t_{i+1}$, which holds when β is piece-wise constant with the appropriate number of levels.

$\impliedby$ Consider β that is not piece-wise constant. Recall that we further assume that β is non-decreasing, and discontinuous only on a measure 0 set. Following algebra steps similar to those in Lemma C.3, but for general β :

$$-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{W}_k = -\lim_{k \rightarrow \infty} \frac{1}{k} \log \int_{\theta_1 > \theta_2} w(\theta_1, \theta_2) \bar{P}_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2) \quad (21)$$

$$= \inf_{\theta_1 > \theta_2} \left(-\lim_{k \rightarrow \infty} \frac{1}{k} \log \bar{P}_k(\theta_j, \theta_i | \beta) \right) \quad (22)$$

$$= 0 \quad (23)$$

Where the last line follows from β continuous at some θ_1 , and so $\lim_{\theta_2 \rightarrow \theta_1} \bar{P}_k(\theta_1, \theta_2 | \beta) = 1$.

Intuitively, what goes wrong with continuous β is that $\bar{P}_k(\theta_1, \theta_2 | \beta)$ does not converge uniformly:

$$\forall \epsilon, k, \exists \theta_2 \neq \theta_1 \quad \bar{P}_k(\theta_1, \theta_2) > \epsilon$$

i.e. close by items are very hard to distinguish from one another. Then, because the large deviations rate of $\overline{W}_k$ is dominated by the worst rates under the integral, we don't get a positive rate. $\square$

C.4 Proofs of Lemma 3.1 and Theorem 3.1

Remark C.2. *The KL divergence for two Bernoulli random variables is continuous and strictly convex, with minima at $a = b$, when $a, b \in (0, 1)$. Note that $\inf_a \{g_i \text{KL}(a||t_i) + g(i+1) \text{KL}(a||t_{i+1})\}$, for all feasible g , is also continuous and strictly convex in t_i, t_{i+1} , with minima at $t_i = t_{i+1}$.*

One consequence of the above fact is that fixing either t_i or t_{i+1} and moving the other farther away monotonically increases KL, while moving it closer decreases KL.

C.4.1 Proof of Theorem 3.1

Proof. We use the same notation as the proof for Lemma C.3.

Part 1.

$$\lim_{k \rightarrow \infty} W_k = \lim_{k \rightarrow \infty} \sum_{0 \leq i < j < M} \left[\int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) P_k(\theta_1, \theta_2 | \beta) d(\theta_1, \theta_2) \right] \quad (24)$$

$$= \sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) d(\theta_1, \theta_2) \quad (25)$$

(25) follows from bounded convergence and $P_k(\theta_1, \theta_2 | \beta) \rightarrow 1$ for $\theta_1 \in S_j, \theta_2 \notin S_j$. Thus choosing $\mathbf{s}$ to maximize (25) maximizes the asymptotic value of W_k .

Part 2. Follows directly from Lemma C.3. $\square$

C.4.2 Proof of Lemma 3.1

Proof. Recall $r(\mathbf{t}) \triangleq -\lim_{k \rightarrow \infty} \frac{1}{k} \log W_k = \min_{0 \leq i \leq M-2} \inf_{a \in \mathbb{R}} \{g_{i+1} \text{KL}(a||t_{i+1}) + g_i \text{KL}(a||t_i)\}$.

We show the following:

$r(\mathbf{t}) =$

$$\min \left(\log(1 - t_1)^{-g_1}, \right. \\ \left. \log \left[(1 - t_{i-1})^{\frac{g_{i-1}}{g_{i-1} + g_i}} (1 - t_i)^{\frac{g_i}{g_{i-1} + g_i}} + t_{i-1}^{\frac{g_{i-1}}{g_{i-1} + g_i}} t_i^{\frac{g_i}{g_{i-1} + g_i}} \right]^{-g_{i-1} - g_i} \text{ for } 1 < i < M - 1, \right. \\ \left. \log(t_{M-2})^{-g_{M-2}} \right)$$

and $\mathbf{t}^*$ maximizes $r_w(\mathbf{t}) \iff$ all the terms inside the minimization $r_w(\mathbf{t}^*)$ are equal. Further, the optimal levels $\mathbf{t}^*$ are unique. The result immediately follows, that $\{t_i\}$ is the unique solution that equalizes the rates inside the minimization, by noting that the optimal r has $t_0 = 0, t_{M-1} = 1$.

We first prove the alternative form for r . Note that $\{g_{i-1} \text{KL}(a||t_{i-1}) + g_i \text{KL}(a||t_i)\}$ is convex in a , and so we can find an analytic form for the infimum over a .

Let $a_i = \arg \inf_{a \in [t_{i-1}, t_i]} \{g_{i-1} \text{KL}(a||t_{i-1}) + g_i \text{KL}(a||t_i)\}$

$$\implies \nabla_{a_i} [g_{i-1} \text{KL}(a_i||t_{i-1}) + g_i \text{KL}(a_i||t_i)] = 0$$

$$\implies \nabla_{a_i} \left[g_{i-1} \left(a_i \log \frac{a_i}{t_{i-1}} + (1 - a_i) \log \frac{1 - a_i}{1 - t_{i-1}} \right) + g_i \left(a_i \log \frac{a_i}{t_i} + (1 - a_i) \log \frac{1 - a_i}{1 - t_i} \right) \right] = 0$$

$$\implies g_{i-1} \left(\log \frac{a_i}{t_{i-1}} - \log \frac{1 - a_i}{1 - t_{i-1}} \right) + g_i \left(\log \frac{a_i}{t_i} - \log \frac{1 - a_i}{1 - t_i} \right) = 0$$

$$\implies \log \left(\frac{a_i}{1 - a_i} \right)^{g_{i-1} + g_i} = \log \left(\frac{t_{i-1}}{1 - t_{i-1}} \right)^{g_{i-1}} + \log \left(\frac{t_i}{1 - t_i} \right)^{g_i}$$

$$\implies \frac{a_i}{1 - a_i} = \left[\left(\frac{t_{i-1}}{1 - t_{i-1}} \right)^{g_{i-1}} \left(\frac{t_i}{1 - t_i} \right)^{g_i} \right]^{\frac{1}{g_{i-1} + g_i}}$$

$$\implies a_i = \frac{c}{1 + c}, \text{ where } c = \left[\left(\frac{t_{i-1}}{1 - t_{i-1}} \right)^{g_{i-1}} \left(\frac{t_i}{1 - t_i} \right)^{g_i} \right]^{\frac{1}{g_{i-1} + g_i}}$$

Then,

$$g_{i-1} \text{KL}(a_i||t_{i-1}) + g_i \text{KL}(a_i||t_i)$$

$$\begin{aligned} &= g_{i-1} a \log \frac{a}{t_{i-1}} + g_i a \log \frac{a}{t_i} + g_{i-1} (1 - a) \log \frac{1 - a}{1 - t_{i-1}} + g_i (1 - a) \log \frac{1 - a}{1 - t_i} \\ &= a \left[(g_{i-1} + g_i) \log \frac{a}{1 - a} + g_{i-1} \log \frac{1 - t_{i-1}}{t_{i-1}} + g_i \log \frac{1 - t_i}{t_i} \right] + \log(1 - a)^{g_{i-1} + g_i} - \log(1 - t_{i-1})^{g_{i-1}} (1 - t_i)^{g_i} \\ &= (g_{i-1} + g_i) \log(1 - a) - \log(1 - t_{i-1})^{g_{i-1}} (1 - t_i)^{g_i} \end{aligned} \quad (26)$$

$$\begin{aligned} &= -(g_{i-1} + g_i) \log \left[1 + \left[\left(\frac{t_{i-1}}{1 - t_{i-1}} \right)^{g_{i-1}} \left(\frac{t_i}{1 - t_i} \right)^{g_i} \right]^{\frac{1}{g_{i-1} + g_i}} \right] (1 - t_{i-1})^{\frac{g_{i-1}}{g_{i-1} + g_i}} (1 - t_i)^{\frac{g_i}{g_{i-1} + g_i}} \\ &= -(g_{i-1} + g_i) \log \left[(1 - t_{i-1})^{\frac{g_{i-1}}{g_{i-1} + g_i}} (1 - t_i)^{\frac{g_i}{g_{i-1} + g_i}} + t_{i-1}^{\frac{g_{i-1}}{g_{i-1} + g_i}} t_i^{\frac{g_i}{g_{i-1} + g_i}} \right] \end{aligned} \quad (27)$$

Where line (26) uses $\frac{a}{1-a} = c$ and $(g_{i-1} + g_i) \log c = \log \left[\left(\frac{t_{i-1}}{1 - t_{i-1}} \right)^{g_{i-1}} \left(\frac{t_i}{1 - t_i} \right)^{g_i} \right]$. Note that the first and last rates emerge, respectively, by plugging in $t_0 = 0, t_M = 1$, which holds trivially at the optimum from monotonicity.

We note that a similar derivation, of the large deviation rate for two binomial distributions with different probability of successes and match rates, appears in Glynn and Juneja [17]. In that work, the authors seek to optimize the g in order to identify the single best item out of a set of possible items, and a concave program emerges. In this work, because we optimize the probability of successes and care about retrieving a ranking of the items, no such concave or convex program emerges.

Now we show that $\mathbf{t}^*$ maximizes $r_w(\mathbf{t}) \iff$ all the terms inside the minimization $r_w(\mathbf{t})$ are equal.

equalizes $\implies$ optimal. Let $r(i)$ be the i th term in the minimization, starting at $i = 1$. Note that (holding the other fixed) increasing t_i increases the i th term monotonically and decreases the $(i + 1)$ th term monotonically. Suppose β s.t. $r(i) = r(j) \forall i, j$. To increase the minimization term, one must increase $r(i)$, $\forall i$. To increase $r(1)$, t_1 must increase, regardless of what the other levels are. Then, to increase $r(2)$, t_2 must increase ... to increase $r(M - 2)$, t_{M-2} must increase. However, to increase $r(M - 1)$, t_{M-2} must decrease, and we have a contradiction. Thus, one cannot increase all terms simultaneously.

equalizes $\Leftarrow$ optimal. Suppose $\mathbf{t}$ maximizes $r(\mathbf{t})$ but the terms inside the minimization are not equal. Then $\exists i$ s.t. $r(i) = \min_j r(j)$ and either $r(i) \neq r(i-1)$ or $r(i) \neq r(i+1)$. $r(i)$ can be increased without lowering the overall rate. This method can be repeated $\forall i : r(i) = \min_j r(j)$ and so $\mathbf{t}$ would not be optimal, a contradiction.

Uniqueness follows from the overall rate unique determining t_1, t_{M-2} and so iteratively uniquely determining the rest. $\square$

C.5 Additional necessary lemmas

Now, we begin the set-up that will lead to a proof for Theorem 3.2. It turns out that proving the theorem requires, in the process, essentially proving our convergence result with $M \rightarrow \infty$, Theorem B.1. For Theorem 3.2, we need a lower bound for t_1 as a function of M . This seems hard to do in general. Luckily, in our case, there is a property for how $\mathbf{t}^*$ changes when M is doubled. Using this property, we can derive that $t_1^* \geq \mathcal{O}(M^{-3})$.

Recall that step-wise increasing β with M intervals $S_i = [s_i, s_{i+1})$ has levels $\{t_i\}_{i=0}^{M-1}$, where $t_0 = 1, t_{M-1} = 1$, and $s_0 \triangleq 0, s_M \triangleq 1$.

Furthermore, we use the following notation for the large deviation rate

$$r_i = -(g_{i-1} + g_i) \log \left[(1 - t_{i-1})^{\frac{g_{i-1}}{g_{i-1}+g_i}} (1 - t_i)^{\frac{g_i}{g_{i-1}+g_i}} + t_{i-1}^{\frac{g_{i-1}}{g_{i-1}+g_i}} t_i^{\frac{g_i}{g_{i-1}+g_i}} \right] \quad (28)$$

for $i \in \{1 \dots M-1\}$, which implies $r_1 = -g_1 \log(1 - t_1)$ and $r_{M-1} = -g_{M-2} \log(t_{M-2})$.

We further use r^{M-1} to be the rate achieved by the optimal β_M with M intervals.

Lemma C.5. Suppose g uniform, i.e. $g_i = 1, \forall i$ and that β_M has values $\{t_i\}_{i=0}^{M-1}$. Then β_{2M-1} has values $\{t'_i\}_{i=0}^{2M-2}$, where $t'_{2i} = t_i, \forall i \in \{0 \dots M-1\}$, $t'_1 = \frac{1}{2}(1 - \sqrt{1 - t_1})$ and $t'_{2M-3} = \frac{1}{2}(1 + \sqrt{t_{M-2}})$.

Proof. We first set the values $t'_{2i} = t_i$ and then optimally choose the remaining values t'_k, k odd. Then, we show that the resulting large deviation rates between all adjacent pairs are equal. Then, by the proof of Lemma 3.1, which showed that equalizing the rates between adjacent intervals is a sufficient condition for optimality, β_{2M-1} has the levels $\{t'_i\}_{i=0}^{2M-2}$.

Let r' denote rates between adjacent t' as r does for t . Supposing $t'_2 = t_1$, we find t'_1 such that $r'_1 = r'_2$ and $t'_1 < t'_2$.

$$\begin{aligned} -\log(1 - t'_1) &= -2 \log \left[\sqrt{(1 - t'_1)(1 - t'_2)} + \sqrt{t'_1 t'_2} \right] \\ \implies 1 - t'_1 &= (1 - t'_1)(1 - t'_2) + t'_1 t'_2 + 2\sqrt{(1 - t'_1)(1 - t'_2)t'_1 t'_2} \\ \implies t'_1 &= \frac{1}{2} \left(1 - \sqrt{1 - t'_2} \right) = \frac{1}{2} (1 - \sqrt{1 - t_1}) \end{aligned}$$

Similarly, $r'_{2M-3} = r'_{2M-2}$ when $t'_{2M-3} = \frac{1}{2}(1 + \sqrt{t_{M-2}})$. It follows that $r'_1 = r'_2 = r'_{2M-3} = r'_{2M-2}$ by choosing such t'_1, t'_{2M-3} .

Next, we find $t'_k \in (t'_{k-1}, t'_{k+1})$ for $k \in \{3, 5, \dots, 2M-5\}$ such that the rates $r'_k = r'_{k+1}$.

$$\begin{aligned} -2 \log \left[\sqrt{(1-t'_k)(1-t'_{k-1})} + \sqrt{t'_k t'_{k-1}} \right] &= -2 \log \left[\sqrt{(1-t'_k)(1-t'_{k+1})} + \sqrt{t'_k t'_{k+1}} \right] \\ \implies t'_k &= \frac{c}{1+c}, \text{ where } c = \left[\frac{\sqrt{1-t'_{k+1}} - \sqrt{1-t'_{k-1}}}{\sqrt{t'_{k-1}} - \sqrt{t'_{k+1}}} \right]^2 \end{aligned}$$

Now, we show that $r'_k = r'_j, \forall j, k$ by showing that the difference between each rate r_i and its analogous rate r'_{2i} is constant. $r_k = r_j, \forall j, k$ by assumption and so $r'_k = r'_j, \forall j, k$ follows.

$r_{M-1} = -\log t_{M-2}$ and $r'_{2M-2} = -\log \frac{1}{2}(1 + \sqrt{t_{M-2}})$. Thus if $r_i = -\log x$ for some x , then $r_{2i} = -\log \frac{1}{2}(1 + \sqrt{x})$ would imply that all the rates are equal. Thus, it is sufficient to show that

$$\left[\sqrt{(1-t'_{2i-1})(1-t'_{2i})} + \sqrt{t'_{2i-1} t'_{2i}} \right]^2 = \frac{1}{2} \left[1 + \sqrt{(1-t_{i-1})(1-t_i)} + \sqrt{t_{i-1} t_i} \right] \quad (29)$$

$$\equiv \left[\sqrt{\left(1 - \frac{c}{1+c}\right)(1-t_i)} + \sqrt{\frac{c}{1+c} t_i} \right]^2 = \frac{1}{2} \left[1 + \sqrt{(1-t_{i-1})(1-t_i)} + \sqrt{t_{i-1} t_i} \right] \quad (30)$$

$$\text{where } c = \left[\frac{\sqrt{1-t_i} - \sqrt{1-t_{i-1}}}{\sqrt{t_{i-1}} - \sqrt{t_i}} \right]^2$$

The proof for (30) is algebraically tedious and is shown in Remark C.3 below.

Then, by the proof of Lemma 3.1, which shows that equalizing the rates inside the minimization terms implies an optimal $\{t_i\}$, β_{2M-1} has the levels $\{t'_i\}_{i=0}^{2M-2}$. □

Remark C.3.

$$\begin{aligned} \left[\sqrt{\left(1 - \frac{c}{1+c}\right)(1-t_i)} + \sqrt{\frac{c}{1+c} t_i} \right]^2 &= \frac{1}{2} \left[1 + \sqrt{(1-t_{i-1})(1-t_i)} + \sqrt{t_{i-1} t_i} \right] \\ \text{where } c &= \left[\frac{\sqrt{1-t_i} - \sqrt{1-t_{i-1}}}{\sqrt{t_{i-1}} - \sqrt{t_i}} \right]^2 \end{aligned}$$

Proof. Let $x = \sqrt{t_i}, y = \sqrt{1-t_i}, z = \sqrt{t_{i-1}}$, and $w = \sqrt{1-t_{i-1}}$. Note that $x > z, w > y, y = 1 - x^2, w = 1 - z^2$. Then,

$$\frac{c}{c+1} = \frac{(y-w)^2}{2-2xz-2yw}, \text{ and } \frac{1}{c+1} = \frac{(x-z)^2}{2-2xz-2yw}$$

(To show the above two equalities, factor out $\frac{1}{(x-z)^2}$ from numerator and denominator, and substitute $y = 1 - x^2, w = 1 - z^2$).

Now, the left hand side:

$$\begin{aligned}
& \left[\sqrt{\left(1 - \frac{c}{1+c}\right)(1-t_i)} + \sqrt{\frac{c}{1+c}t_i} \right]^2 \\
&= \frac{1}{2-2xz-2yw} \left[\sqrt{(x-z)^2y^2} + \sqrt{(y-w)^2x^2} \right]^2 \\
&= \frac{(x-z)^2y^2 + (y-w)^2x^2 + 2xy(x-z)(w-y)}{2-2xz-2yw} \quad \sqrt{(y-w)^2} = w-y, \sqrt{(x-z)^2} = x-z \\
&= \frac{z^2y^2 + w^2x^2 - 2wxyz}{2-2xz-2yw}
\end{aligned}$$

The right hand side:

$$\begin{aligned}
& \frac{1}{2} \left[1 + \sqrt{(1-t_{i-1})(1-t_i)} + \sqrt{t_{i-1}t_i} \right] \\
&= \frac{1}{2} [1 + (wy + xz)]
\end{aligned}$$

Multiplying both sides by $2 - 2xz - 2yw$, we have:

$$\begin{aligned}
& \left[\sqrt{\left(1 - \frac{c}{1+c}\right)(1-t_i)} + \sqrt{\frac{c}{1+c}t_i} \right]^2 = \frac{1}{2} \left[1 + \sqrt{(1-t_{i-1})(1-t_i)} + \sqrt{t_{i-1}t_i} \right] \\
&\quad \equiv z^2y^2 + w^2x^2 - 2wxyz = 1 - (wy + xz)^2 \\
&\quad \equiv z^2(1-x^2) + (1-z^2)x^2 - 2wxyz = 1 - w^2y^2 - x^2z^2 - 2wxyz \\
&\quad \equiv z^2 - 2x^2z^2 + x^2 = 1 - (1-z^2)(1-x^2) - x^2z^2 \\
&\quad \equiv 0 = 0
\end{aligned}$$

□

Corollary C.1. Suppose g uniform, i.e. $g_i = 1, \forall i$. $\forall \epsilon > 0, \exists M$ s.t. $\forall M' \geq M, r^{M'} < \epsilon$.

Proof. Let $M = 2^N, M' = 2^{N+1} - 1$, for some N . We show that $r^{M'} \leq \frac{1}{2}r^M$. The corollary follows by noting that $r^{K'} < r^K \forall K' > K$ and that $r^K < \infty, \forall K$.

$$\begin{aligned}
r^M - r^{M'} &= -\log t_{M-2}^M + \log t_{M'-2}^{M'} \\
&= -\log t_{M-2}^M + \log \left[\frac{1}{2} + \frac{1}{2} \sqrt{t_{M-2}^M} \right] \quad \text{Lemma C.5} \\
&= \log \left[\frac{1}{2} \frac{1}{t_{M-2}^M} + \frac{1}{2} \frac{1}{\sqrt{t_{M-2}^M}} \right] \\
&\geq -\frac{1}{2} \log t_{M-2}^M \quad \sqrt{t_{M-2}^M} \geq t_{M-2}^M \\
&\implies r^{M'} \leq \frac{1}{2} r^M
\end{aligned}$$

□

Corollary C.2. Suppose g uniform, i.e. $g_i = 1, \forall i$. $\forall \delta > 0, \exists N$ s.t. $\forall M \geq N, \max_k t_k^M - t_{k-1}^M < \delta$.

Proof. This corollary follows directly from Corollary C.1. If the rates are upper bounded, then so are the level differences.

We first find where the rate is minimized given a width between levels of δ

$$\begin{aligned} x_m &= \arg \min_x -2 \log \left[\sqrt{(1-x-\delta)(1-x)} + \sqrt{x(x+\delta)} \right] \\ &= \frac{1}{2} - \frac{1}{2}\delta \end{aligned}$$

Then given an upper bound of ϵ on the rate, there is a bound on δ determined by the largest possible difference at levels symmetric around $\frac{1}{2}$.

$$\begin{aligned} r^L &= -2 \log \left[2 \sqrt{\left(\frac{1}{2} - \delta\right)\left(\frac{1}{2} + \delta\right)} \right] \\ &= -\log [1 - 4\delta^2] \\ &\geq \epsilon \text{ when } \delta > \frac{1}{2} \sqrt{1 - e^{-\epsilon}} \end{aligned}$$

□

Lemma C.6. Suppose g is non-decreasing in θ . Then, $t_{M-2} \geq 1 - \frac{1}{M-1}$.

Proof. Note that, with uniform matching, $\forall x \in (0, 1], y \in [0, 1-x]$ the rate with values $t_{i-1} = y, t_i = y+x$ is no more than the last with $t_{M-2} = 1-x$. With width x , in other words, the extreme points have a larger rates than the middle points. For $i \notin \{1, M-1\}$:

$$\begin{aligned} r_i &= \inf_a \{g_{i-1} \text{KL}(a||t_{i-1}) + g_i \text{KL}(a||t_i)\} \\ &= \inf_a \{\text{KL}(a||y) + \text{KL}(a||y+x)\} && \text{uniform matching} \\ &= -2 \log \left[(1-y)^{\frac{1}{2}}(1-y-x)^{\frac{1}{2}} + y^{\frac{1}{2}}(y+x)^{\frac{1}{2}} \right] \\ &= -\log \left[(1-y)(1-y-x) + y(y+x) + 2[(1-y)(1-y-x)y(y+x)]^{1/2} \right] \\ &\leq -\log(1-x) \end{aligned} \tag{31}$$

Where line (31) follows from line (27).

By the proof of Lemma 3.1, the optimal levels equalize the rates between each level. Then, when g is non-decreasing, $g_{M-2} \geq g_\ell, \forall \ell \in \{1 \dots M-3\}$. Then, at the same level differences, the rate corresponding to the last level is no smaller. Thus, to equalize the rates, the last width must be no larger than any other width. Thus, $t_{M-2} \geq 1 - \frac{1}{L}$. □

Lemma C.7. With uniform matching ($g_i = 1$), $r^{2^{N+1}-1} \geq \frac{1}{5}r^{2^N}$.

Proof. Let $K = 2^N, K' = 2^{N+1} - 1$. Note that $t_{K-1}^K \geq \frac{1}{2}$ by Lemma C.6.

$$\begin{aligned}
r^K - r^{K'} &= -\log t_{K-2}^K + \log t_{K'-2}^{K'} \\
&= -\log t_{K-2}^K + \log \left[\frac{1}{2} + \frac{1}{2} \sqrt{t_{K-2}^K} \right] && \text{Lemma C.5} \\
&= \log \left[\frac{1}{2} \frac{1}{t_{K-2}^K} + \frac{1}{2} \frac{1}{\sqrt{t_{K-2}^K}} \right] \\
&\leq \log (t_{K-2}^K)^{-\frac{4}{5}} && \frac{1}{2} (t_{K-2}^K)^{-1} + \frac{1}{2} (t_{K-2}^K)^{-\frac{1}{2}} \leq (t_{K-2}^K)^{-\frac{4}{5}} \text{ when } t_{K-2}^K \in \left[\frac{1}{2}, 1 \right] \\
\implies r^{K'} &\geq \frac{1}{5} r^K
\end{aligned}$$

□

Lemma C.8. *With uniform matching ($g_i = 1$), $\exists C > 0$ s.t. $\forall M, t_1^M \geq CM^{-3}$.*

Proof. By Lemma C.7, $\exists C_2 > 0$ s.t. $r^M \geq C_2 5^{-\lceil \log_2 M \rceil}$. Then

$$\begin{aligned}
-\log(1 - t_1^M) &= r^M \\
&\geq C_2 5^{-\lceil \log_2 M \rceil} \\
\implies t_1^M &\geq 1 - \exp \left[-C_2 5^{-\lceil \log_2 M \rceil} \right] \\
&\geq 1 - \exp \left[-C_3 M^{-\frac{1}{\log_5 2}} \right] \\
&\geq \frac{e-1}{e} C_3 M^{-\frac{1}{\log_5 2}} && e^{-x} \leq 1 - \frac{e-1}{e} x \text{ for } x \in [0, 1] \\
\implies \exists C > 0 \text{ s.t. } t_1^M &\geq CM^{-3}
\end{aligned}$$

□

Corollary C.3. *With monotonically non-decreasing g , $\exists C > 0$ s.t. $\forall M, t_1^M \geq CM^{-3}$.*

Proof. The result follows from noting that t_1^M with uniform matching lower bounds the first value with any other monotonically non-decreasing g , which is a direct application of Lemma B.1 – scale g such that $g_1 = 1$. Then, $g_j \geq 1, j > 1$ and $g_0 \leq 1$. Then, the condition of the lemma holds. □

Lemma C.9. *The run-time of `NestedBisection` is $O(M \log^2 \frac{1}{\delta})$, where δ is the bisection grid width and M is the number of intervals.*

Proof. The outer bisection, in *main*, runs at most $\log_2 \frac{2}{\delta} + 1$ iterations. Each outer iteration calls `BisectNextLevel` $M - 3$ times, and the inner bisection in each call runs for at most $\log_2 \frac{2}{\delta}$ iterations. Thus the run-time of algorithm is $O(M \log^2 \frac{1}{\delta})$. □

C.6 Proof for Theorem 3.2

Finally, we are ready to prove Theorem 3.2. It follows from formalizing the relationship between δ , the bisection grid width, and ϵ , the additive approximation error in the rate function.

Proof. Recall M is the number of intervals (levels) in β . We use j, t, t^* to denote the levels in a certain iteration, the returned levels, and the optimal levels, respectively. We use $r(\cdot)$ to denote the individual rates between returned levels, i.e. $r(1) = -g_1 \log(1 - t_1)$, $r(m) = \{g_{m-1} \text{KL}(a_m || t_{m-1}) + g_m \text{KL}(a_m || t_m)\}$, $m \in \{2 \dots M-2\}$, $r(M-1) = -g_{M-2} \log(t_{M-2})$, and use r^* to denote the optimal rate.

By Lemma C.6, $t_{M-2}^* \geq 1 - \frac{1}{M-1}$. By assumption, $t_{M-2}^* < 1 - \delta$. Thus, $t_{M-2}^* \in [1 - \frac{1}{M-1}, 1 - \delta]$, the starting interval for the outer bisection.

First, suppose the outer bisection terminates such that $t_{M-2} \leq t_{M-2}^* + \delta$. We prove that this case always occurs below.

In this case, $r^* - r(M-1)$ is at most $-g_{M-2} \log(t_{M-2}^*) + g_{M-2} \log(t_{M-2}^* + \delta) = g_{M-2} \log\left(\frac{t_{M-2}^* + \delta}{t_{M-2}^*}\right)$. For all $m \in \{M-2 \dots 2\}$, in the final *CalculateOtherLevels* call the algorithm will use bisection to match the corresponding rate with this last rate, $r(M-1) = -g_{M-2} \log(t_{M-2})$, setting t_{m-2} to the smallest value such that $r(m) \leq r(M-1)$ (i.e. the right end of the final interval is chosen).

Then, $\forall m \in \{M-2 \dots 2\}$, $r(m) \in [r(M-1) - \epsilon(\delta), r(M-1)]$, where $\epsilon(\delta)$ is an upper bound on the change in the rate functions with a shift of δ in one of the parameters.

For now, assume $r(1) = -g_1 \log(t_1) \geq r(M-1)$. We prove that this occurs below. Then,

$$\begin{aligned} r(m) &\geq r(M-1) - \epsilon(\delta) \\ &\geq -g_{M-2} \log(t_{M-2}^* + \delta) - \epsilon(\delta) \end{aligned} \quad \forall m \in \{1 \dots M\}$$

Now we characterize $\epsilon(\delta)$ in the region $[t_1^* + \delta, t_{M-2}^* + \delta]$. In particular, we want to bound the rate loss from the other levels $r(m), m > 1$ after the $g_{M-2} \log\left(\frac{t_{M-2}^* + \delta}{t_{M-2}^*}\right)$ loss in $r(M-1)$. Note that the only source of error is a level shifting right by δ . $r_j(\cdot)$ denotes individual rates between levels j in an intermediary iteration. Let a_i' be the minimum point inside the rate infimum after the shift by δ .

$$\begin{aligned} \epsilon(\delta) &= \sup_{t_{i-1}, t_i} [g_{i-1} \text{KL}(a_i || t_{i-1}) + g_i \text{KL}(a_i || t_i) - g_{i-1} \text{KL}(a_i' || t_{i-1} + \delta) - g_i \text{KL}(a_i' || t_i)] \\ &\leq \sup_{t_{i-1}, t_i} [g_{i-1} \text{KL}(a_i' || t_{i-1}) + g_i \text{KL}(a_i' || t_i) - g_{i-1} \text{KL}(a_i' || t_{i-1} + \delta) + g_i \text{KL}(a_i' || t_i)] \quad a_i \text{ is inf point} \\ &= \sup_{t_{i-1}, t_i} g_{i-1} \left[a_i' \log \frac{t_{i-1} + \delta}{t_{i-1}} + (1 - a_i') \log \frac{1 - t_{i-1} - \delta}{1 - t_{i-1}} \right] \\ &\leq \sup_{t_{i-1}, t_i} g_{i-1} \left[a_i' \log \frac{t_{i-1} + \delta}{t_{i-1}} \right] \quad \text{2nd term negative} \\ &\leq g_{M-2} \left[\log \frac{t_1^* + \delta}{t_1^*} \right] \quad t_j \geq t_1^*, g_j \leq g_{M-2} \\ \implies r(m) &\geq r^* - g_{M-2} \log\left(\frac{t_{M-2}^* + \delta}{t_{M-2}^*}\right) - g_{M-2} \left[\log \frac{t_1^* + \delta}{t_1^*} \right] \\ &\geq r^* - g_{M-2} \frac{\delta}{t_{M-2}^*} - g_{M-2} \frac{\delta}{t_1^*} \quad \log(1+x) \leq x \\ &\geq r^* - \delta g_{M-2} \left[\frac{M-1}{M-2} + \frac{1}{t_1^*} \right] \quad t_{M-2}^* \geq 1 - \frac{1}{M-1} \end{aligned}$$

By Corollary C.3, $\exists C > 0$ s.t. $t_1^* \geq CM^{-3} \implies r(m) \geq r^* - \delta g_{M-2} \left[\frac{M-1}{M-2} + CM^3 \right]$. Then, let $\delta = \frac{\epsilon}{g_{M-2} \left[\frac{M-1}{M-2} + CM^3 \right]}$. Supposing the algorithm terminates in such an iteration, it finds an ϵ -optimal

β in time $O\left(M \log^2 \frac{g_{M-2} \left[\frac{M-1}{M-2} + CM^3\right]}{\epsilon}\right) = O\left(M \log^2 \frac{M}{\epsilon}\right)$.

Next, we show that the algorithm only terminates the outer bisection when $u \leq t_{M-2}^* + \delta$. The claim follows from $\ell \leq t_{M-2}^*$ being an algorithm invariant. The initial $\ell = 1 - \frac{1}{M-1} \leq t_{M-2}^*$ by Lemma C.6. ℓ can only be set to be $> t_{M-2}^*$ if in the current iteration, $j_{M-2} > t_{M-2}^*$ and $r_j(1) < r_j(M-1)$. However, if $j_{M-2} \geq t_{M-2}^*$, then $r_j(1) \geq r_j(M-1)$ ($j_m \geq t_m^* \forall m$), following from a shifting argument like that given in Lemma 3.1 and that the inner bisection is such that $r_j(m) \leq r_j(M-1), m \in \{2 \dots M-2\}$, i.e. all the values $t_m > t_m^*$. Thus, $\ell \leq t_{M-2}^*$ is an algorithm invariant and $u > t_{M-2}^* + \delta \implies u - \ell > \delta$.

Finally, we show that $r(1) \geq r(M-1)$ at the returned $\{t_i\}$. By assumption, in the initial iteration, $u \geq t_{M-2}^*$, and recall that the returned $\{t_i\}$ such that $t_{M-2} = u$ from the final iteration. As shown in the previous paragraph, $j_{M-2} \geq t_{M-2}^* \implies r_j(1) \geq r_j(M-1)$. Thus, if the algorithm terminates in the first iteration, then $r(1) \geq r(M-1)$. In any subsequent iteration, u is changed only if $r_j(1) \geq r_j(M-1)$ at its new value. Thus, $r_j(1) \geq r_j(M-1)$ is an algorithm invariant, and $r(1) \geq r(M-1)$.

The algorithm terminates in finite time. Thus, it terminates when $t_{M-2} = u \leq t_{M-2}^* + \delta$ and finds a (ϵ, M, g) -optimal β in time $O\left(M \log^2 \frac{M}{\epsilon}\right)$. □

In Theorem 3.2, there is an guarantee of an additive error away from the optimal rate. To instead have a multiplicative error bound for uniform matching, one can use the lower bound on the optimal rate from Lemma C.7, $\exists C > 0$ s.t. $r^* \geq CM^{-3}$. Then, for uniform matching, the algorithm returns a $(1 - \epsilon)$ multiplicative approximation in time $O\left(M \log^2 \frac{M}{\epsilon}\right)$.

C.7 Proof of Theorem B.1

Let β_M^w denote the optimal β with M intervals for weight function w , with intervals $\mathbf{s}^{wM}$ and levels $\mathbf{t}^{wM}$. Let $q_{wM}(\theta) = i/M$ when $\theta \in [s_i^{wM}, s_{i+1}^{wM})$, i.e. the *quantile* of interval item of type θ is in. Then we have the following convergence result for β_M .

Theorem B.1. *Let g be uniform. Suppose w such that q_{wM} converges uniformly. Then, $\forall C \in \mathbb{N}, \exists \beta^w$ s.t. $\beta_{C2N+1}^w \rightarrow \beta^w$ uniformly as $N \rightarrow \infty$.*

Proof. Note that the condition on q implies that $\exists \overline{M}$ s.t. $\forall M > \overline{M}, \forall \theta, \exists x_\theta$ such that $\theta \in [s_{\lfloor x_\theta M \rfloor}^M, s_{\lceil x_\theta M \rceil}^M)$.

Let $M' = 2M - 1, M'' = 4M - 3, M^q = 2^q M - 2^q + 1$. $\theta \in [s_{\lfloor x_\theta M \rfloor}^M, s_{\lceil x_\theta M \rceil}^M) \implies \beta_M(\theta) = t_{\lfloor x_\theta M \rfloor}^M \in [t_{\lfloor x_\theta M \rfloor - 1}^M, t_{\lfloor x_\theta M \rfloor + 1}^M]$. Then,

$$\begin{aligned} \beta_{M'}(\theta) &= t_{\lfloor x_\theta M' \rfloor}^{M'} \\ &= t_{\lfloor x_\theta (2M-1) \rfloor}^{M'} \\ &\in [t_{2\lfloor x_\theta M \rfloor - 2}^{M'}, t_{2\lfloor x_\theta M \rfloor + 2}^{M'}] \\ &\subset [t_{\lfloor x_\theta M \rfloor - 1}^M, t_{\lfloor x_\theta M \rfloor + 1}^M] \end{aligned} \quad \text{Lemma C.5}$$

And, for general q ,

$$\begin{aligned}
\beta_{M^q}(\theta) &= t_{\lfloor x_\theta(2^q M - 2^q + 1) \rfloor}^{M^q} \\
&\in \left[t_{\lfloor x_\theta 2^q M \rfloor - 2^q}^{M^q}, t_{\lfloor x_\theta(2^q M) \rfloor + 1}^{M^q} \right] \\
&\subset \left[t_{2^q \lfloor x_\theta M \rfloor - 2^q}^{M^q}, t_{2^q \lfloor x_\theta M \rfloor + 1}^{M^q} \right] \\
&\subset \left[t_{\lfloor x_\theta M \rfloor - 1}^M, t_{\lfloor x_\theta M \rfloor + 1}^M \right] \quad \text{Lemma C.5}
\end{aligned}$$

Then, $\forall N' > 1, \theta$: $\beta_{2^{N'} M - 2^{N'} + 1}(\theta) \in \left[t_{\lfloor x_\theta M \rfloor - 1}^M, t_{\lfloor x_\theta M \rfloor + 1}^M \right]$ and

$$|\beta_{2^{N'} M - 2^{N'} + 1}(\theta) - \beta_M(\theta)| \leq t_{\lfloor x_\theta M \rfloor + 1}^M - t_{\lfloor x_\theta M \rfloor - 1}^M$$

By Corollary C.2, $\forall \delta > 0, \exists K$ s.t. $\forall K' > K, t_{\lfloor x_\theta K' \rfloor + 1}^{K'} - t_{\lfloor x_\theta K' \rfloor - 1}^{K'} < 2\delta$.

By the Cauchy criterion, $\exists \beta$ s.t. $\beta_{(C-1)2^N + 1} \rightarrow \beta$ uniformly.

By change of variables, $\exists \beta$ s.t. $\beta_{C2^N + 1} \rightarrow \beta$ uniformly. $\square$

Corollary C.4. *For Kendall's tau and Spearman's rho correlation measures, $\exists \beta$ s.t. $\beta_{2^N} \rightarrow \beta$ uniformly as $N \rightarrow \infty$.*

Proof. For Kendall's tau and Spearman's rho, $\{s_i\}$ is spaced such that $\forall i, j, s_i - s_{i-1} = s_j - s_{j-1}$. Thus, $x_\theta = \theta$ meets the criterion. $\square$

C.8 Kendall's tau and Spearman's rho related proofs

Definition C.1 (see e.g. Embrechts et al. [11], Nelsen [26]). *The population version of Kendall-tau correlation between item true quality and rating scores is proportional to*

$$W_k^\tau \triangleq 2 \int_{\theta_1 > \theta_2} P_k(\theta_1, \theta_2) d\theta_1 d\theta_2$$

Similarly, given items with qualities $\theta_1, \theta_2, \theta_3$, the population version of Spearman's rho correlation between item true quality and rating scores is

$$W_k^\rho \triangleq 6 \int_{\theta_1 > \theta_2, \theta_3} P_k(\theta_1, \theta_3) d\theta_1 d\theta_2 d\theta_3$$

Lemma C.10. *Spearman's ρ can also be written as being proportional to $\int_{\theta_1 > \theta_2} (\theta_1 - \theta_2) P_k(\theta_1, \theta_2) d\theta_1 d\theta_2$, i.e. with $w(\theta_1, \theta_2) = (\theta_1 - \theta_2)$.*

Proof. Recall $P_k(\theta_1, \theta_3) = \Pr((\theta_1 - \theta_2)(x_1^k - x_3^k) > 0)$

$$\begin{aligned}
&= \int_{\theta_1 > \theta_2, \theta_3} \Pr(x_1^k - x_3^k > 0) d\theta_1 d\theta_2 d\theta_3 + \int_{\theta_1 < \theta_2, \theta_3} \Pr(x_1^k - x_3^k < 0) d\theta_1 d\theta_2 d\theta_3 \\
&= \int_{\theta_1, \theta_3} \Pr(x_1^k - x_3^k > 0) \left[\int_{\theta_2=0}^{\theta_1} d\theta_2 \right] d\theta_1 d\theta_3 + \int_{\theta_1, \theta_3} \Pr(x_1^k - x_3^k < 0) \left[\int_{\theta_2=\theta_1}^1 d\theta_2 \right] d\theta_1 d\theta_3 \\
&= \int_{\theta_1, \theta_3} \left[\Pr(x_1^k - x_3^k > 0) \theta_1 + \Pr(x_1^k - x_3^k < 0) (1 - \theta_1) \right] d\theta_1 d\theta_3 \\
&= \int_{\theta_1, \theta_3} \left[\Pr(x_1^k - x_3^k < 0) + \theta_1 \left[\Pr(x_1^k - x_3^k > 0) - \Pr(x_1^k - x_3^k < 0) \right] \right] d\theta_1 d\theta_3
\end{aligned}$$

Similarly,

$$\begin{aligned}
Pr((\theta_1 - \theta_2)(x_1^k - x_3^k) < 0) &= \\
&= \int_{\theta_1, \theta_3} \left[Pr(x_1^k - x_3^k > 0) + \theta_1 \left[Pr(x_1^k - x_3^k < 0) - Pr(x_1^k - x_3^k > 0) \right] \right] d\theta_1 d\theta_3 \\
&= \int_{\theta_1, \theta_3} \left[Pr(x_3^k - x_1^k > 0) + \theta_3 \left[Pr(x_3^k - x_1^k < 0) - Pr(x_3^k - x_1^k > 0) \right] \right] d\theta_1 d\theta_3
\end{aligned}$$

Where the second equality follows from θ_1, θ_3 interchangeable. Then

$$\begin{aligned}
W_k^\rho &= 3 \int_{\theta_1, \theta_2} (\theta_1 - \theta_2) P_k(\theta_1, \theta_2) d\theta_1 d\theta_2 \\
&= \int_{\theta_1 > \theta_2} 6(\theta_1 - \theta_2) P_k(\theta_1, \theta_2) d\theta_1 d\theta_2
\end{aligned}$$

□

Note that Spearman's ρ is similar to Kendall's τ with an additional weighting for how far apart the two values that are flipped are.

Lemma C.11. *When w is constant, i.e. for Kendall's τ rank correlation, the intervals $\mathbf{s}$ that maximize (25),*

$$\sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) d(\theta_1, \theta_2) = \sum_{0 \leq i < j < M} (s_{i+1} - s_i)(s_{j+1} - s_j) \quad (32)$$

, are $\{s_i = \frac{i}{M}\}_{i=0}^M$.

Lemma C.12. *When w is $(\theta_1 - \theta_2)$, i.e. for Spearman's ρ rank correlation, the intervals $\mathbf{s}$ that maximize (25),*

$$\sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) d(\theta_1, \theta_2) \quad (33)$$

are $\{s_i = \frac{i}{M}\}_{i=0}^M$, i.e. the same as those for Kendall's τ .

Proof.

$$\begin{aligned}
\sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} w(\theta_1, \theta_2) d(\theta_1, \theta_2) &= \sum_{0 \leq i < j < M} \int_{\theta_2 \in S_i, \theta_1 \in S_j} (\theta_1 - \theta_2) d(\theta_1, \theta_2) \\
&= \sum_{0 < i < j \leq M} \left(\frac{s_j + s_{j-1}}{2} - \frac{s_i + s_{i-1}}{2} \right) (s_i - s_{i-1})(s_j - s_{j-1})
\end{aligned}$$

Finding an asymptotically optimal $\{s_i\}$ then is a constrained third order polynomial maximization problem with M variables. The maximum is achieved at $\{s_i = \frac{i}{M}\}_{i=0}^M$, as for Kendall's tau correlation. □