

# MIP\* = RE

Zhengfeng Ji<sup>\*1</sup>, Anand Natarajan<sup>†2,3</sup>, Thomas Vidick<sup>‡3</sup>, John Wright<sup>§2,3,4</sup>, and Henry Yuen<sup>¶5</sup>

<sup>1</sup>*Centre for Quantum Software and Information, University of Technology Sydney*

<sup>2</sup>*Institute for Quantum Information and Matter, California Institute of Technology*

<sup>3</sup>*Department of Computing and Mathematical Sciences, California Institute of Technology*

<sup>4</sup>*Department of Computer Science, University of Texas at Austin*

<sup>5</sup>*Department of Computer Science and Department of Mathematics, University of Toronto*

November 7, 2022

## Abstract

We show that the class MIP\* of languages that can be decided by a classical verifier interacting with multiple all-powerful quantum provers sharing entanglement is equal to the class RE of recursively enumerable languages. Our proof builds upon the quantum low-degree test of (Natarajan and Vidick, FOCS 2018) and the classical low-individual degree test of (Ji, et al., 2020) by integrating recent developments from (Natarajan and Wright, FOCS 2019) and combining them with the recursive compression framework of (Fitzsimons et al., STOC 2019).

An immediate byproduct of our result is that there is an efficient reduction from the Halting Problem to the problem of deciding whether a two-player nonlocal game has entangled value 1 or at most  $\frac{1}{2}$ . Using a known connection, undecidability of the entangled value implies a negative answer to Tsirelson's problem: we show, by providing an explicit example, that the closure  $C_{qa}$  of the set of quantum tensor product correlations is strictly included in the set  $C_{qc}$  of quantum commuting correlations. Following work of (Fritz, Rev. Math. Phys. 2012) and (Junge et al., J. Math. Phys. 2011) our results provide a refutation of Connes' embedding conjecture from the theory of von Neumann algebras.

---

\*Email: zhengfeng.ji@uts.edu.au

†Email: anandn@caltech.edu

‡Email: vidick@caltech.edu

§Email: wright@cs.utexas.edu

¶Email: hyuen@cs.toronto.edu

# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Interactive proof systems . . . . .	6
1.2	Statement of result . . . . .	10
1.3	Consequences . . . . .	12
1.4	Open questions . . . . .	14
<b>2</b>	<b>Proof Overview</b>	<b>17</b>
<b>3</b>	<b>Preliminaries</b>	<b>24</b>
3.1	Turing machines . . . . .	24
3.2	Linear spaces . . . . .	27
3.3	Finite fields . . . . .	29
3.3.1	Subfields and bases . . . . .	29
3.3.2	Bit string representations . . . . .	32
3.4	Polynomials and the low-degree code . . . . .	34
3.5	Linear spaces and registers . . . . .	35
3.6	Measurements and observables . . . . .	36
3.7	Generalized Pauli observables . . . . .	37
<b>4</b>	<b>Conditionally Linear Functions, Distributions, and Samplers</b>	<b>41</b>
4.1	Conditionally linear functions and distributions . . . . .	41
4.2	Conditionally linear samplers . . . . .	47
<b>5</b>	<b>Nonlocal Games and MIP*</b>	<b>50</b>
5.1	Games and strategies . . . . .	50
5.2	Distance measures . . . . .	52
5.3	The class MIP* . . . . .	56
5.4	Normal form verifiers . . . . .	57
<b>6</b>	<b>Types</b>	<b>59</b>
6.1	Typed samplers, deciders, and verifiers . . . . .	59
6.2	Graph distributions . . . . .	61
6.3	Detyping typed verifiers . . . . .	63
<b>7</b>	<b>Classical and Quantum Low-degree Tests</b>	<b>66</b>
7.1	The classical low-degree test . . . . .	66
7.1.1	The game . . . . .	66
7.1.2	Complexity of the classical low-degree test. . . . .	71
7.2	The Magic Square game . . . . .	72
7.3	The Pauli basis test . . . . .	77
7.3.1	The game . . . . .	77
7.3.2	Canonical parameters and complexity of the Pauli basis test . . . . .	85

<b>8</b>	<b>Introspection Games</b>	<b>88</b>
8.1	Overview . . . . .	88
8.2	The introspective verifier . . . . .	89
8.2.1	Complexity of the introspective verifier . . . . .	95
8.2.2	Introspection theorem . . . . .	96
8.3	Completeness of the introspective verifier . . . . .	97
8.3.1	Preliminary lemmas . . . . .	97
8.3.2	Completeness of the introspective verifier . . . . .	98
8.4	Soundness of the introspective verifier . . . . .	101
8.4.1	The Pauli twirl . . . . .	101
8.4.2	Preliminary lemmas . . . . .	107
8.4.3	Soundness proof . . . . .	109
<b>9</b>	<b>Oracularization</b>	<b>118</b>
9.1	Overview . . . . .	118
9.2	Oracularizing normal form verifiers . . . . .	118
9.3	Completeness and complexity of the oracularized verifier . . . . .	119
9.4	Soundness of the oracularized verifier . . . . .	121
<b>10</b>	<b>Answer Reduction</b>	<b>123</b>
10.1	Circuit preliminaries . . . . .	124
10.2	A Cook-Levin theorem for bounded deciders . . . . .	125
10.3	A succinct 5SAT description for deciders . . . . .	132
10.4	A PCP for normal form deciders . . . . .	137
10.4.1	Preliminaries . . . . .	137
10.4.2	The PCP . . . . .	138
10.5	A normal form verifier for the PCP . . . . .	143
10.5.1	Parameters and notation . . . . .	144
10.5.2	The answer-reduced verifier . . . . .	144
10.5.3	Main theorem for answer reduction . . . . .	146
10.6	Completeness of the answer-reduced verifier . . . . .	148
10.7	Soundness of the answer-reduced verifier . . . . .	149
<b>11</b>	<b>Parallel Repetition</b>	<b>158</b>
11.1	The anchoring transformation . . . . .	159
11.2	The parallel repetition transformation . . . . .	160
<b>12</b>	<b>Gap-Preserving Compression</b>	<b>164</b>
12.1	Proof of Theorem 12.1 . . . . .	164
12.2	An MIP* protocol for the Halting problem . . . . .	169
12.3	An explicit separation . . . . .	174
<b>A</b>	<b>Analysis of the Pauli basis test</b>	<b>176</b>
A.1	Preliminaries . . . . .	176
A.2	Strategies . . . . .	178
A.3	Expanding the Hilbert space and defining commuting observables . . . . .	182

A.4	Combining the $X$ and $Z$ measurements . . . . .	187
A.5	Applying the classical low-degree test . . . . .	195
A.6	Pulling the $X$ and $Z$ measurements apart . . . . .	206

# 1 Introduction

For integer  $n, k \geq 2$  define the quantum (spatial) correlation set  $C_{qs}(n, k)$  as the subset of  $\mathbb{R}^{n^2k^2}$  that contains all tuples  $(p_{abxy})$  representing families of bipartite distributions that can be locally generated in non-relativistic quantum mechanics. Formally,  $(p_{abxy}) \in C_{qs}(n, k)$  if and only if there exist separable Hilbert spaces  $\mathcal{H}_A$  and  $\mathcal{H}_B$ , for every  $x \in \{1, \dots, n\}$  (resp.  $y \in \{1, \dots, n\}$ ), a collection of projections  $\{A_a^x\}_{a \in \{1, \dots, k\}}$  on  $\mathcal{H}_A$  (resp.  $\{B_b^y\}_{b \in \{1, \dots, k\}}$  on  $\mathcal{H}_B$ ) that sum to identity, and a state (unit vector)  $\psi \in \mathcal{H}_A \otimes \mathcal{H}_B$  such that

$$\forall x, y \in \{1, 2, \dots, n\}, \quad \forall a, b \in \{1, 2, \dots, k\}, \quad p_{abxy} = \psi^*(A_a^x \otimes B_b^y)\psi. \quad (1)$$

Note that due to the normalization conditions on  $\psi$  and on  $\{A_a^x\}$  and  $\{B_b^y\}$ , for each  $x, y$ ,  $(p_{abxy})$  is a probability distribution on  $\{1, 2, \dots, k\}^2$ . By taking direct sums it is easy to see that the set  $C_{qs}(n, k)$  is convex. Let  $C_{qa}(n, k)$  denote its closure (it is known that  $C_{qs}(n, k) \neq C_{qa}(n, k)$ , see [Slo19a]).

Our main result is that the family of sets  $\{C_{qa}(n, k)\}_{n, k \in \mathbb{N}}$  is extraordinarily complex, in the following computational sense. For any  $0 < \varepsilon < 1$  define the  $\varepsilon$ -weak membership problem for  $C_{qa}$  as the problem of deciding, given  $n, k \in \mathbb{N}$  and a point  $p = (p_{abxy}) \in \mathbb{R}^{n^2k^2}$ , whether  $p$  lies in  $C_{qa}(n, k)$  or is  $\varepsilon$ -far from it in  $\ell_1$  distance, promised that one is the case. Then we show that for any given  $0 < \varepsilon < 1$  the  $\varepsilon$ -weak membership problem for  $C_{qa}$  cannot be solved by a Turing machine that halts with the correct answer on every input.

We show this by directly reducing the Halting problem to the weak membership problem for  $C_{qa}$ : we show that for all  $0 < \varepsilon < 1$  and any Turing machine  $\mathcal{M}$  one can efficiently compute integers  $n, k \in \mathbb{N}$  and a linear functional  $\ell_{\mathcal{M}}$  on  $\mathbb{R}^{n^2k^2}$  such that, whenever  $\mathcal{M}$  halts it holds that

$$\sup_{p \in C_{qa}(n, k)} |\ell_{\mathcal{M}}(p)| = 1, \quad (2)$$

whereas if  $\mathcal{M}$  does not halt then

$$\sup_{p \in C_{qa}(n, k)} |\ell_{\mathcal{M}}(p)| \leq 1 - \varepsilon. \quad (3)$$

By standard results in convex optimization, this implies the aforementioned claim on the undecidability of the  $\varepsilon$ -weak membership problem for  $C_{qa}$  (for any  $0 < \varepsilon < 1$ ).

Our result has interesting consequences for long-standing conjectures in quantum information theory and the theory of von Neumann algebras. Through a connection that follows from the work of Navascues, Pironio, and Acin [NPA08] the undecidability result implies a negative answer to Tsirelson's problem [Tsi06]. Let  $C_{qc}(n, k)$  denote the set of quantum commuting correlations, which is the set of tuples  $(p_{abxy})$  arising from operators  $\{A_a^x\}$  and  $\{B_b^y\}$  acting on a single Hilbert space  $\mathcal{H}$  and a state  $\psi \in \mathcal{H}$  such that

$$\forall x, y \in \{1, \dots, n\}, \quad \forall a, b \in \{1, \dots, k\}, \quad p_{abxy} = \psi^*(A_a^x B_b^y)\psi \quad \text{and} \quad [A_a^x, B_b^y] = 0. \quad (4)$$

Then Tsirelson's problem asks if, for all  $n, k$ , the sets  $C_{qa}(n, k)$  and  $C_{qc}(n, k)$  are equal. Using results from [NPA08] we give integer  $n, k$  and an explicit linear function  $\ell$  on  $\mathbb{R}^{n^2k^2}$  such that

$$\sup_{p \in C_{qc}(n, k)} |\ell(p)| = 1, \quad \text{but} \quad \sup_{p \in C_{qa}(n, k)} |\ell(p)| \leq \frac{1}{2},$$

which implies that  $C_{qa}(n, k) \neq C_{qc}(n, k)$ . By an implication of Fritz [Fri12] and Junge et al. [JNP<sup>+</sup>11] we further obtain that Connes' Embedding Conjecture [Con76] is false; in other words, there exist type II<sub>1</sub>

von Neumann factors that do not embed in an ultrapower of the hyperfinite  $\text{II}_1$  factor. We explain these connections in more detail in Section 1.3 below.

Our approach to constructing such linear functionals on correlation sets goes through the theory of interactive proofs from complexity theory. To explain this connection we first review the concept of interactive proofs. The reader familiar with interactive proofs may skip the next section to arrive directly at a formal statement of our main complexity-theoretic result in Section 1.2.

## 1.1 Interactive proof systems

An *interactive proof system* is an abstraction that generalizes the familiar notion of *proof*. Intuitively, given a formal statement  $z$  (for example, “this graph admits a proper 3-coloring”), a proof  $\pi$  for  $z$  is information that enables one to check the validity of  $z$  more efficiently than without access to the proof (in this example,  $\pi$  could be an explicit assignment of colors to each vertex of the graph).

Complexity theory formalizes the notion of proof in a way that emphasizes the role played by the verification procedure. To explain this, first recall that in complexity theory a *language*  $L$  is a subset of  $\{0, 1\}^*$ , the set of all bit strings of any length, that intuitively represents all problem instances to which the answer should be “yes”. For example, the language  $L = \text{3-COLORING}$  contains all strings  $z$  such that  $z$  is the description (according to some pre-specified encoding scheme) of a 3-colorable graph  $G$ . We say that a language  $L$  admits efficiently verifiable proofs if there exists an algorithm  $V$  (formally, a polynomial-time Turing machine) that satisfies the following two properties: (i) for any  $z \in L$  there is a string  $\pi$  such that  $V(z, \pi)$  returns 1 (we say that  $V$  “accepts”), and (ii) for any  $z \notin L$  there is no string  $\pi$  such that  $V(z, \pi)$  accepts. Property (i) is generally referred to as the *completeness* property, and (ii) is the *soundness*. The set of all languages  $L$  with both these completeness and soundness properties is denoted by the complexity class NP.

Research in complexity and cryptography in the 1980s and 1990s led to a significant generalization of the notion of “efficiently verifiable proof”. The first modification is to allow *randomized* verification procedures by relaxing (i) and (ii) to *high probability* statements: every  $z \in L$  should have a proof  $\pi$  that is accepted *with probability at least  $c$*  (the completeness parameter), and for no  $z \notin L$  should there be a proof  $\pi$  that is accepted *with probability larger than  $s$*  (the soundness parameter). A common setting is to take  $c = \frac{2}{3}$  and  $s = \frac{1}{3}$ ; standard amplification techniques reveal that the exact values do not significantly affect the class of languages that admit such proofs, provided that they are chosen within reasonable bounds.

The second modification is to allow *interactive* verification. Informally, this means that instead of receiving a proof string  $\pi$  in its entirety and making a decision based on it, the verification algorithm (called the “verifier”) instead communicates with another algorithm called a “prover”, and based on the communication decides whether  $z \in L$ . There are no restrictions on the computational power of the prover, whereas the verifier is required to run in polynomial time.<sup>1</sup>

To understand how randomization and interaction can help for proof checking, consider the following example of an interactive proof for the language GRAPH NON-ISOMORPHISM, which contains all pairs of graphs  $(G_0, G_1)$  such that  $G_0$  and  $G_1$  are *not* isomorphic.<sup>2</sup> It is not known if GRAPH NON-ISOMORPHISM  $\in$  NP, because it is not clear how to give an efficiently verifiable proof string that two graphs  $G_0$  and  $G_1$  are

---

<sup>1</sup>The reader may find the following mental model useful: in an interactive proof, an all-powerful prover is trying to convince a skeptical, but computationally limited, verifier that a string  $z$  (known to both) lies in the set  $L$ , even when it may be that in fact  $z \notin L$ . By interactively interrogating the prover, the verifier can reject false claims, i.e. determine with high statistical confidence whether  $z \in L$  or not. Importantly, the verifier is allowed to probabilistically and adaptively choose its messages to the prover.

<sup>2</sup>Here and in the rest of the section, we implicitly assume that graphs and tuples of graphs have a canonical encoding as binary strings.

not isomorphic. (A proof of isomorphism is, of course, trivial: given a bijection from the vertices of  $G_0$  to those of  $G_1$  it is straightforward to verify that the bijection induces an isomorphism.) However, consider the following *randomized, interactive* verification procedure. Suppose the input to the verifier and prover is a pair of  $n$ -vertex graphs  $(G_0, G_1)$  (if the graphs do not have the same number of vertices, they are trivially non-isomorphic and the verification procedure can automatically accept). The verifier first selects a uniformly random  $b \in \{0, 1\}$  and a uniformly random permutation  $\sigma$  of  $\{1, \dots, n\}$  and sends the graph  $H = \sigma(G_b)$  to the prover. The prover is then supposed to respond with a bit  $b' \in \{0, 1\}$ ; if  $b' = b$  the verifier accepts and if  $b' \neq b$  it rejects.

Clearly, if  $G_0$  and  $G_1$  are not isomorphic then there exists a prover strategy to compute  $b$  from  $H$  with probability 1: using its unlimited computational power, the prover can determine whether  $H$  is isomorphic to  $G_0$  or to  $G_1$ . However, if  $G_0$  and  $G_1$  are isomorphic then the distribution of  $H$  is uniform over the isomorphism class of  $G_0$ , which is the same as the isomorphism class of  $G_1$ , and the prover (despite having unlimited computational power) cannot distinguish between whether the verifier generated  $H$  using  $G_0$  or  $G_1$ . Thus the probability that *any* prover can correctly guess  $b' = b$  is exactly  $\frac{1}{2}$ . As a result, we have shown that the graph non-isomorphism problem has an interactive proof system with completeness  $c = 1$  and soundness  $s = \frac{1}{2}$ . Note how little “information” is communicated by the prover: a single bit! The extreme succinctness of the “proof” comes from the fact that whether  $G_0$  is isomorphic to  $G_1$  determines whether a prover can reliably compute, given the data available to it (which is  $G_0, G_1$ , and  $H$ ), the correct bit  $b$ .

We denote by IP the class of languages that admit randomized interactive proof systems such as the one just described. The class IP is easily seen to contain NP, but it is thought to be a much larger class: one of the famous results of complexity theory is that IP is exactly the same as PSPACE [LFKN90, Sha90], the class of languages decidable by Turing machines using polynomial space.<sup>3</sup> Thus a polynomial-time verifier, when augmented with the ability to interrogate an all-powerful prover and use randomization, can solve computational problems that are (believed to be) vastly more difficult than those that can be checked using static, deterministic proofs (i.e. NP problems).

**Multiprover interactive proofs.** We now discuss a generalization of interactive proofs called *multiprover interactive proofs*. Here, a polynomial-time verifier can interact with *two* (or more) provers to decide whether a given instance  $z$  is in a language  $L$  or not. In this setting, after the verifier and all the provers receive the common input  $z$ , the provers are not allowed to communicate with each other, and the verifier “cross-interrogates” the provers in order to decide if  $z \in L$ . The provers may coordinate a joint strategy ahead of time, but once the protocol begins the provers can only interact with the verifier. As we will see, the extra condition that the provers cannot communicate with each other is a powerful constraint that can be leveraged by the verifier.

Consider the computational problem of deciding membership in a *promise* language called GAP-MAXCUT. A promise language  $L$  is specified by two disjoint subsets  $L_{yes}, L_{no} \subseteq \{0, 1\}^*$ , and the task is to decide whether a given instance  $z$  is in  $L_{yes}$  or  $L_{no}$ , promised that  $z \in L_{yes} \cup L_{no}$ . In a proof system for a promise language, the completeness case consists of accepting with probability at least  $c$  if  $z \in L_{yes}$ , and the soundness case consists of accepting with probability at most  $s$  if  $z \in L_{no}$ . If  $z \notin L_{yes} \cup L_{no}$ , then there are no constraints on the behavior of the verifier.

The promise language GAP-MAXCUT is defined as follows: GAP-MAXCUT<sub>yes</sub> (resp. GAP-MAXCUT<sub>no</sub>)

---

<sup>3</sup>The reason PSPACE is considered a “difficult” class of problems is because many computational problems believed to require super-polynomial or exponential time (such as 3-COLORING or deciding whether a quantified Boolean formula is true) can be solved using a polynomial amount of space.

is the set of all graphs  $G$  with a *cut* (i.e. a bipartition of the vertices) such that at least 90% of edges cross the cut (resp. at most 60% of edges cross the cut).<sup>4</sup> For simplicity, we also assume that all graphs in  $\text{GAP-MAXCUT}_{yes} \cup \text{GAP-MAXCUT}_{no}$  are regular, i.e. the degree is a constant across all vertices in the graph.

The GAP-MAXCUT problem clearly lies in NP, since given a candidate cut it is easy to count the number of edges that cross it and verify that it is at least 90% of the total number of edges. Observe that the length of the proof and the time required to verify it are linear in the size of the graph (the number of vertices and edges). *Finding* the proof is of course much harder, but we are only concerned with the complexity of the verification procedure.

Now consider the following simple *two-prover interactive proof system* for GAP-MAXCUT. Given a graph  $G$ , the verification procedure first samples a uniformly random edge  $e = \{u, v\}$  in  $G$ . It then sends a uniformly random  $x \in \{u, v\}$  to the first prover, and a uniformly random  $y \in \{u, v\}$  to the second prover. Each prover sees its respective question only and is expected to respond with a single bit,  $a, b \in \{0, 1\}$  respectively. The verification procedure accepts if and only if  $a = b$  if  $x = y$ , and  $a \neq b$  if  $x \neq y$ .

We claim that the verification procedure described in the preceding paragraph is a *multiprover interactive proof system* for the language GAP-MAXCUT, with completeness  $c = 0.95$  and soundness  $s = 0.9$ , in the following sense. First, whenever  $G \in \text{GAP-MAXCUT}_{yes}$  then there is a successful strategy for the provers: specifically, the provers can fix an optimal bipartition and consistently answer “0” when asked about a vertex from one side of the partition, and “1” when asked about a vertex from the other side; assuming there exists a cut that is crossed by at least 90% of the edges, this strategy succeeds with probability at least  $\frac{1}{2} + \frac{1}{2}0.9$ , where the first factor  $\frac{1}{2}$  arises from the case when both provers are sent the same vertex, in which case they always succeed.

Conversely, suppose given a strategy for the provers that is accepted with probability  $p = \frac{1}{2} + \frac{1}{2}(1 - \delta)$  when the verification procedure is executed on a (regular)  $n$ -vertex graph  $G$ . We then claim that  $G$  has a cut crossed by at least a  $1 - 2\delta$  fraction of all edges. To show this, we leverage the non-communication assumption on the provers. Since either prover’s question is always a single vertex, their strategy can be represented by a function from the vertices of  $G$  to answers in  $\{0, 1\}$ . Any such function specifies a bipartition of  $G$ . While the provers’ bipartitions need not be identical, the fact that they succeed with high probability, for the case when they are sent the same vertex, implies that they must be consistent with high probability. Finally, the fact that they also succeed with high probability when sent opposite endpoints of a randomly chosen edge implies that either prover’s bipartition must be cut by a large number of edges. Taking the contrapositive establishes the soundness property.

We denote by MIP the class of languages that have multiprover interactive proof systems such as the one described in the preceding paragraph. Note that, in comparison to the NP verification procedure for GAP-MAXCUT considered earlier, the interactive, two-prover verification is much more efficient in terms of the effort required for the verifier. Assuming the graph is provided in a convenient format,<sup>5</sup> it is possible to sample a random edge and verify the provers’ answers in time and space that scales *logarithmically* with the size of the graph. This exponential improvement in the efficiency of the verification procedure serves as the starting point for another celebrated result from complexity theory: MIP is exactly the same as the class NEXP [BFL91], which are problems that admit *exponential-time checkable proofs*.<sup>6</sup> The class NEXP

---

<sup>4</sup>The specific numbers 90% and 60% are not too important; the only thing that really matters is that the first one is strictly less than 100% and the second strictly larger than 50%, as otherwise the problem becomes much easier.

<sup>5</sup>For example, the graph can be specified via a circuit that takes as input an edge index — using some arbitrary ordering — and returns labels for the two endpoints of the edge.

<sup>6</sup>An example of such a problem is the language SUCCINCT-3-COLORING, which contains descriptions of polynomial-size circuits  $C$  that specify a 3-colorable graph  $G_C$  on *exponentially many* vertices.



contains PSPACE, but is believed to be much larger; this suggests that the ability to interrogate more than one prover enables a polynomial-time verifier to verify much more complex statements.

**Nonlocal games.** In this paper we will only be concerned with multiprover interactive proof systems that consist of a single round of communication with two provers: the verifier first sends its questions to each of the provers, the provers respond with their answers, and the verifier decides whether to accept or reject. The class of problems that admit such interactive proofs is denoted  $\text{MIP}(2, 1)$ , and it is known that  $\text{MIP} = \text{MIP}(2, 1)$  [FL92]. Such proof systems have a convenient reformulation using the language of *nonlocal games*, that we now explain.

In a nonlocal game, we say that a verifier interacts with multiple non-communicating *players* (instead of provers — there is no formal difference between the two terms). An  $n$ -question,  $k$ -answer nonlocal game  $\mathfrak{G}$  is specified by two procedures: a *question sampling* procedure that samples a pair of questions  $(x, y) \in \{1, \dots, n\}^2$  for the players according to a distribution  $\mu$  (known to the verifier and the players), and a *decision* procedure that takes as input the players' questions and their respective answers  $a, b \in \{1, \dots, k\}$  and evaluates a predicate  $D(x, y, a, b) \in \{0, 1\}$  to determine the verifier's acceptance or rejection. In classical complexity theory, the main quantity associated with a nonlocal game  $\mathfrak{G}$  is its *classical value*, which is the maximum success probability that two cooperating but non-communicating players have in the game. Formally, the classical value is defined as

$$\text{val}(\mathfrak{G}) = \sup_{p \in C_c(n, k)} \sum_{x, y} \mu(x, y) \sum_{a, b} D(x, y, a, b) p_{abxy}, \quad (5)$$

where the set  $C_c(n, k)$  is the set of *classical correlations*, which are tuples  $(p_{abxy})$  such that there exists a set  $\Lambda$  with probability measure  $\nu$  and for every  $\lambda \in \Lambda$  functions  $A^\lambda, B^\lambda : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, k\}$  such that

$$\forall x, y \in \{1, 2, \dots, n\}, \quad \forall a, b \in \{1, 2, \dots, k\}, \quad p_{abxy} = \Pr_{\lambda \sim \nu} (A^\lambda(x) = a \wedge B^\lambda(y) = b).$$

This definition captures the intuitive notion that a classical strategy for the players is specified by (i) a distribution  $\nu$  on  $\Lambda$  that represents some probabilistic information shared by the players that is independent of the verifier's questions, and (ii) two functions  $A^\lambda, B^\lambda$  that represent each players' "local strategy" for answering given their shared randomness  $\lambda$  and question  $x$  or  $y$ .<sup>7</sup> Note that due to the shared randomness  $\lambda$ , the set  $C_c(n, k)$  is a (closed) convex subset of  $[0, 1]^{n^2 k^2}$ .

To make the connection with interactive proof systems, observe that the assertion that  $L \in \text{MIP}(2, 1)$  precisely amounts to the specification of an efficient mapping<sup>8</sup> from problem instances  $z$  to games  $\mathfrak{G}_z$  such that whenever  $z \in L$  then  $\text{val}(\mathfrak{G}_z) \geq \frac{2}{3}$ , whereas if  $z \notin L$  then  $\text{val}(\mathfrak{G}_z) \leq \frac{1}{3}$ . Thus the complexity of the optimization problem (5) captures the complexity of the decision problem  $L$ . The aforementioned characterization of MIP as the class NEXP by [BFL91] shows that in general this optimization problem is very difficult: it is as hard as deciding any language in NEXP.

<sup>7</sup>For the functional analyst we briefly note that if we define a tensor

$$L = \sum_{x, y, a, b} \mu(x, y) D(x, y, a, b) e_{xa} \otimes e_{yb} \in \mathbb{R}^{nk} \otimes \mathbb{R}^{nk}$$

then  $\text{val}(\mathfrak{G}) = \|L\|_{\ell_1^{\text{tr}}(\ell_\infty)^k \otimes_\epsilon \ell_1^{\text{tr}}(\ell_\infty)^k}$ , with  $\otimes_\epsilon$  denoting the injective tensor norm of Banach spaces. (For more connections between interactive proofs, nonlocal games and tensor norms we refer to the survey [PV16].)

<sup>8</sup>Here by "efficient" we mean that there should be a polynomial-time Turing machine that on input  $z$  returns (i) a polynomial-size randomized circuit that samples from  $\mu$ , and (ii) a polynomial-size circuit that evaluates the predicate  $D$ .

## 1.2 Statement of result

We now introduce the main complexity class that is the focus of this paper:  $\text{MIP}^*$ , the “entangled-prover” analogue of the class  $\text{MIP}$  considered earlier. Informally the class  $\text{MIP}^*$ , first introduced in [CHTW04], contains all languages that can be decided by a classical polynomial-time verifier interacting with multiple *quantum* provers sharing *entanglement*. We focus on the class  $\text{MIP}^*(2, 1)$ , which corresponds to the setting of one-round protocols with two provers. Equivalently, a language  $L$  is in  $\text{MIP}^*(2, 1)$  if and only if there is an efficient mapping from instances  $z \in \{0, 1\}^*$  to nonlocal games  $\mathfrak{G}_z$  such that if  $z \in L$ , then  $\text{val}^*(\mathfrak{G}_z) \geq 2/3$  and otherwise  $\text{val}^*(\mathfrak{G}_z) \leq 1/3$ . Here, for an  $n$ -question,  $k$ -answer game  $\mathfrak{G}$ , we let  $\text{val}^*(\mathfrak{G})$  denote its *entangled value*, which is defined as

$$\text{val}^*(\mathfrak{G}) = \sup_{p \in C_q(n,k)} \sum_{x,y} \mu(x,y) \sum_{a,b} D(x,y,a,b) p_{abxy}, \quad (6)$$

where the set  $C_q(n,k)$  is the set of all finite-dimensional quantum correlations, i.e. correlations of the form (1) where  $\mathcal{H}$  is restricted to be finite-dimensional. Although the sets  $C_q(n,k)$  and  $C_{qs}(n,k)$  in general are distinct [CS18], it is an easy exercise to verify that they have the same closure  $C_{qa}(n,k)$ , and therefore the supremum in (6) can equivalently be taken over  $C_q(n,k)$ ,  $C_{qs}(n,k)$  or  $C_{qa}(n,k)$ . We use  $C_q$  for convenience in the analysis.

Since  $C_c(n,k) \subseteq C_q(n,k)$  for all  $n,k$ , we have that  $\text{val}(\mathfrak{G}) \leq \text{val}^*(\mathfrak{G})$ ; in other words, quantum spatial strategies can perform at least as well as classical strategies in a nonlocal game.

The consideration of quantum strategies and the set  $C_{qs}(n,k)$  for the definition of  $\text{MIP}^*$  is motivated by a long line of works in the foundations of quantum mechanics around the topic of *Bell inequalities*, that are linear functionals which separate the sets  $C_c(n,k)$  and  $C_{qs}(n,k)$ . The simplest such functional is the *CHSH inequality* [CHSH69], that shows  $C_c(2,2) \subsetneq C_{qs}(2,2)$ . The CHSH inequality can be reformulated as a game  $\mathfrak{G}$  such that  $\text{val}^*(\mathfrak{G}) > \text{val}(\mathfrak{G})$ . This game is very simple: it is defined by setting  $\mu(x,y) = \frac{1}{4}$  for all  $x,y \in \{0,1\}$  and  $D(x,y,a,b) = 1$  if and only if  $a \oplus b = x \wedge y$ . It can be shown that  $\text{val}(\mathfrak{G}) = \frac{3}{4}$  and  $\text{val}^*(\mathfrak{G}) = \frac{1}{2} + \frac{1}{2\sqrt{2}} > \frac{3}{4}$ . The study of Bell inequalities is a large area of research not only in foundations, where they are a tool to study the nonlocal properties of entanglement, but also in quantum cryptography, where they form the basis for cryptographic protocols for e.g. quantum key distribution [Eke91].

The introduction of entanglement in the setting of interactive proofs has interesting consequences for complexity theory; indeed it is not *a priori* clear how the class  $\text{MIP}^*$  compares to  $\text{MIP}$ . Take a language  $L \in \text{MIP}(2, 1)$ , and let  $z$  be an instance. Then the associated game  $\mathfrak{G}_z$  is such that  $\text{val}(\mathfrak{G}_z) \geq \frac{2}{3}$  if  $z \in L$ , and  $\text{val}(\mathfrak{G}_z) \leq \frac{1}{3}$  otherwise. The fact that in general  $\text{val}^*(\mathfrak{G}_z) \geq \text{val}(\mathfrak{G}_z)$  (and that as demonstrated by the CHSH game inequality can be strict) cuts both ways. On the one hand, the soundness property can be affected, so that instances  $z \notin L$  could have  $\text{val}^*(\mathfrak{G}_z) = 1$ , meaning that we would not be able to establish that  $L \in \text{MIP}^*$ . On the other hand, a language  $L \in \text{MIP}^*(2, 1)$  may not necessarily be in  $\text{MIP}$ , because for  $z \in L$  the fact that  $\text{val}^*(\mathfrak{G}_z) \geq \frac{2}{3}$  does not automatically imply  $\text{val}(\mathfrak{G}_z) > \frac{1}{3}$  (in other words, the game  $\mathfrak{G}_z$  may require the players to use a quantum strategy in order to win with probability greater than  $1/3$ ). Just as the complexity of the class  $\text{MIP}$  is characterized by the complexity of approximating the classical value of nonlocal games (the optimization problem in (5)), the complexity of  $\text{MIP}^*$  is intimately related to the complexity of approximating the entangled value of games (the optimization problem in (6)).

In [IV12] the first non-trivial lower bound on  $\text{MIP}^*$  was shown, establishing that  $\text{MIP} = \text{NEXP} \subseteq \text{MIP}^*$ . (Earlier results [KKM<sup>+</sup>11, IKM09] gave more limited hardness results, for approximating the entangled value up to inverse polynomial precision.) This was proved by arguing that for the specific games constructed by [BFL91] that show  $\text{NEXP} \subseteq \text{MIP}$ , the classical and entangled values are approximately the

same. In other words, the classical soundness and completeness properties of the proof system of [BFL91] are maintained in the presence of shared entanglement between the provers. Following [IV12] a sequence of works [Vid16, Ji16, NV18b, Ji17, NV18a, FJVY19] established progressively stronger lower bounds on the complexity of approximating the entangled value of nonlocal games, culminating in [NW19] which showed that approximating the entangled value is at least as hard as NEXP, the collection of languages decidable in non-deterministic *doubly exponential* time. This proves that  $\text{NEXP} \subseteq \text{MIP}^*$ , and since it is known that  $\text{NEXP} \not\subseteq \text{NEEXP}$  it follows that  $\text{MIP} \neq \text{MIP}^*$ .

In contrast to these increasingly strong lower bounds the only upper bound known on  $\text{MIP}^*$  is the trivial inclusion  $\text{MIP}^* \subseteq \text{RE}$ , the class of recursively enumerable languages, i.e. languages  $L$  such that there exists a Turing machine  $\mathcal{M}$  such that  $x \in L$  if and only if  $\mathcal{M}$  halts and accepts on input  $x$ . This inclusion follows since the supremum in (6) can be approximated from below by performing an exhaustive search in increasing dimension and with increasing accuracy. We note that, in addition to containing all decidable languages, this class also contains undecidable problems such as the Halting problem, which is to decide whether a given Turing machine eventually halts.

Our main result is a proof of the reverse inclusion:  $\text{RE} \subseteq \text{MIP}^*$ . Combined with the preceding observation it follows that

$$\text{MIP}^* = \text{RE},$$

which is a full characterization of the power of entangled-prover interactive proofs. In particular for any  $0 < \varepsilon < 1$ , it is an undecidable problem to determine whether a given nonlocal game has entangled value 1 or at most  $1 - \varepsilon$  (promised that one is the case).

**Proof summary.** The proof of the inclusion  $\text{RE} \subseteq \text{MIP}^*$  is obtained by designing an entangled-prover interactive proof for the Halting problem, which is complete for the class RE. Specifically, we design an efficient transformation that maps any Turing machine  $\mathcal{M}$  to a nonlocal game  $\mathfrak{G}_{\mathcal{M}}$  such that, if  $\mathcal{M}$  halts (when run on an empty input tape) then there is a quantum strategy for the provers that succeeds with probability 1 in  $\mathfrak{G}_{\mathcal{M}}$  (i.e.  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) = 1$ ), whereas if  $\mathcal{M}$  does not halt then no quantum strategy can succeed with probability larger than  $\frac{1}{2}$  in the game (i.e.  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) \leq \frac{1}{2}$ ). Furthermore, the game  $\mathfrak{G}_{\mathcal{M}}$  has the property that whenever  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) = 1$  then this fact is witnessed by a *synchronous* strategy, i.e. a strategy where the players always give the same answer when simultaneously asked the same question. Synchronous strategies, or correlations, were first introduced in [PSS<sup>+</sup>16] and have played an important role in approaches to CEP based on quantum information and the study of nonlocal games; see e.g. [KPS18] and references therein. In the paper we use a terminology of *projective, consistent and commuting* (PCC) strategies (see Definition 5.11 in Section 5.1), a notion which implies the notion of being synchronous.

A very rough sketch of this construction is as follows (we give a detailed overview in Section 2). Given an infinite family of games  $\{\mathfrak{G}_n\}_{n \in \mathbb{N}}$ , we say that the family is *uniformly generated* if there is a polynomial-time Turing machine that on input  $n$  returns a description of the game  $\mathfrak{G}_n$ . Given a game  $\mathfrak{G}$  and  $p \in [0, 1]$  let  $\mathcal{E}(\mathfrak{G}, p)$  denote the minimum local dimension of an entangled state shared by the players in order for them to succeed in  $\mathfrak{G}$  with probability at least  $p$ .

We proceed in two steps. First, we design a *compression* procedure for a specific class of nonlocal games that we call *normal form*. Given as input a uniformly generated family  $\{\mathfrak{G}_n\}_{n \in \mathbb{N}}$  of normal form games, the compression procedure returns another uniformly generated family  $\{\mathfrak{G}'_n\}_{n \in \mathbb{N}}$  of normal form games with the following properties: (i) for all  $n$ , if  $\text{val}^*(\mathfrak{G}_{2^n}) = 1$  then  $\text{val}^*(\mathfrak{G}'_n) = 1$ , and (ii) for all  $n$ , if  $\text{val}^*(\mathfrak{G}_{2^n}) \leq \frac{1}{2}$  then  $\text{val}^*(\mathfrak{G}'_n) \leq \frac{1}{2}$  and moreover

$$\mathcal{E}(\mathfrak{G}'_n, \frac{1}{2}) \geq \max \left\{ \mathcal{E}(\mathfrak{G}_{2^n}, \frac{1}{2}), 2^{2^{\Omega(n)}} \right\}.$$

The construction of this compression procedure is our main contribution. Informally, it combines the recursive compression technique developed in [Ji17, FJVV19] with the so-called “introspection” technique of [NW19] that was used to prove  $\text{NEEXP} \subseteq \text{MIP}^*$ . The introspection technique itself relies heavily on the quantum low individual degree test of [NV18a, JNV<sup>+</sup>20] to robustly self-test certain distributions that arise from constructions of *classical* probabilistically checkable proofs. The quantum low-degree test and the introspection technique allow us to avoid the shrinking gap limitation of the results from [FJVV19].

In the second step, we use the compression procedure in an iterated fashion to construct an interactive proof system for the Halting problem. Fix a Turing machine  $\mathcal{M}$  and consider the following family of nonlocal games  $\{\mathfrak{G}_{\mathcal{M},n}^{(0)}\}_{n \in \mathbb{N}}$ : for all  $n \in \mathbb{N}$ , if  $\mathcal{M}$  halts in at most  $n$  steps (when run on an empty input tape), then  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(0)}) = 1$ , and otherwise  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(0)}) \leq \frac{1}{2}$ .<sup>9</sup>

Constructing such a family of games is trivial; furthermore, they can be made in the “normal form” required by the compression procedure. However, consider applying the compression procedure to obtain a family of normal form games  $\{\mathfrak{G}_{\mathcal{M},n}^{(1)}\}_{n \in \mathbb{N}}$ . Then for all  $n \in \mathbb{N}$ , it holds that if  $\mathcal{M}$  halts in at most  $2^n$  steps then  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(1)}) = 1$ , and otherwise  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(1)}) \leq \frac{1}{2}$ , and furthermore any strategy that achieves a value of at least  $\frac{1}{2}$  requires an entangled state of dimension at least  $2^{2^{\Omega(n)}}$ .

Intuitively, one would expect that iterating this procedure and “taking the limit” gives a family of games  $\{\mathfrak{G}_{\mathcal{M},n}^{(\infty)}\}_{n \in \mathbb{N}}$  such that if  $\mathcal{M}$  halts then  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(\infty)}) = 1$  for all  $n \in \mathbb{N}$ , whereas if  $\mathcal{M}$  does not halt then no finite-dimensional strategy can succeed with probability larger than  $\frac{1}{2}$  in  $\mathfrak{G}_{\mathcal{M},n}^{(\infty)}$ , for all  $n \in \mathbb{N}$ ; in particular  $\text{val}^*(\mathfrak{G}_{\mathcal{M},n}^{(\infty)}) \leq \frac{1}{2}$ . Formally, we do not take such a limit but instead define directly the family of games  $\{\mathfrak{G}_{\mathcal{M},n}^{(\infty)}\}_{n \in \mathbb{N}}$  as a *fixed point* of the Turing machine that implements the compression procedure. The game  $\mathfrak{G}_{\mathcal{M}}$  can then be taken as  $\mathfrak{G}_{\mathcal{M},1}^{(\infty)}$ . We describe this in more detail in Section 2.

### 1.3 Consequences

Our result is motivated by a connection with Tsirelson’s problem from quantum information theory, itself related to Connes’ Embedding Conjecture in the theory of von Neumann algebras [Con76]. In a celebrated sequence of papers, Tsirelson [Tsi93] initiated the systematic study of quantum correlation sets. Recall the definition of the set of quantum spatial correlations

$$C_{qs}(n, k) = \{ (p_{abxy}) \mid p_{abxy} = \langle \psi | A_a^x \otimes B_b^y | \psi \rangle, |\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B, \forall xy, \{A_a^x\}_a, \{B_b^y\}_b \text{ POVM} \}, \quad (7)$$

where here  $|\psi\rangle$  ranges over all unit norm vectors  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  with  $\mathcal{H}_A$  and  $\mathcal{H}_B$  arbitrary (separable) Hilbert spaces, and a POVM is defined as a collection of positive semidefinite operators that sum to identity. (From now on we use the Dirac ket notation  $|\psi\rangle$  for states.) Recall the closure  $C_{qa}(n, k)$  of  $C_{qs}(n, k)$ .

Tsirelson observed that there is a natural alternative definition to the quantum spatial correlation set, called the *quantum commuting correlation set* and defined as

$$C_{qc}(n, k) = \{ (p_{abxy}) \mid p_{abxy} = \langle \psi | A_a^x B_b^y | \psi \rangle \}, \quad (8)$$

where  $|\psi\rangle \in \mathcal{H}$  is a quantum state,  $\{A_a^x\}$  and  $\{B_b^y\}$  are POVMs for all  $x, y$ , and  $[A_a^x, B_b^y] = 0$  for all  $a, b, x, y$ . Note the key difference with spatial correlations is that in (8) all operators act on the same (separable) Hilbert space. The requirement that operators associated with different inputs (questions)  $x, y$  commute

---

<sup>9</sup>There is nothing special about the choice of  $\frac{1}{2}$ ; this can be set to any constant that is less than 1.

is arguably a minimal requirement within the context of quantum mechanics for there to not exist any causal connection between outputs (answers)  $a, b$  obtained in response to the respective input.

The set  $C_{qc}(n, k)$  is closed and convex, and it is easy to see that  $C_{qa}(n, k) \subseteq C_{qc}(n, k)$  for all  $n, k \geq 1$ . When Tsirelson initially introduced these sets he claimed that equality holds. However, it was later pointed out that this is not obviously true. The question of equality between  $C_{qc}$  and  $C_{qa}$  (for all  $n, k$ ) is now known as *Tsirelson's problem* [Tsi06]. Let  $C_q(n, k)$  denote the same as  $C_{qs}(n, k)$  except that both  $\mathcal{H}_A$  and  $\mathcal{H}_B$  in (7) are restricted to finite-dimensional spaces. Then more generally one can consider the following chain of inclusions

$$C_q(n, k) \subseteq C_{qs}(n, k) \subseteq C_{qa}(n, k) \subseteq C_{qc}(n, k), \quad (9)$$

for all  $n, k \in \mathbb{N}$ , and ask which (if any) of these inclusions are strict. We let  $C_q, C_{qs}, C_{qa}, C_{qc}$  denote the union of  $C_q(n, k), C_{qs}(n, k), C_{qa}(n, k), C_{qc}(n, k)$ , respectively, over all integers  $n, k \in \mathbb{N}$ . In a breakthrough work, Slofstra established the first separation between these four correlation sets by proving that  $C_{qs} \neq C_{qc}$  [Slo19b]; he later proved the stronger statement that  $C_{qs} \neq C_{qa}$  [Slo19a]. As a consequence of the technique used to demonstrate the separation Slofstra also obtains the complexity-theoretic statement that the problem of determining whether an element  $p$  lies in  $C_{qc}$ , even promised that if it does, then it also lies in  $C_{qa}$ , is undecidable. Interestingly, this is shown by reduction from the *complement* of the halting problem; for our result we reduce from the halting problem (see Section 1.4 for further discussion of this point). Since his work, simpler proofs of Slofstra's results have been found [DPP19, MR18, Col19]. In [CS18], Coladangelo and Stark showed that  $C_q \neq C_{qs}$  by exhibiting a 5-input, 3-output correlation that can be attained using infinite-dimensional spatial strategies (i.e. infinite-dimensional Hilbert spaces, a state and POVMs satisfying (7)) but cannot be attained via finite-dimensional strategies.

As already noted in [FNT14] (and further elaborated on by [FJVY19]), the undecidability of  $\text{MIP}^*(2, 1)$  implies the separation  $C_{qa} \neq C_{qc}$ .<sup>10</sup> This follows from the observation that if  $C_{qa} = C_{qc}$ , then there exists an algorithm that can correctly determine if a nonlocal game  $\mathfrak{G}$  satisfies  $\text{val}^*(\mathfrak{G}) = 1$  or  $\text{val}^*(\mathfrak{G}) \leq \frac{1}{2}$  and always halts: this algorithm interleaves a hierarchy of semidefinite programs providing outer approximations to the set  $C_{qc}$  [NPA08, DLTW08] with a simple exhaustive search procedure providing inner approximations to  $C_q$ . Our result that  $\text{RE} \subseteq \text{MIP}^*(2, 1)$  implies that no such algorithm exists, thus resolving Tsirelson's problem in the negative.

We furthermore exhibit an explicit nonlocal game  $\mathfrak{G}$  such that  $\text{val}^*(\mathfrak{G}) < \text{val}^{\text{co}}(\mathfrak{G}) = 1$ , where  $\text{val}^{\text{co}}(\mathfrak{G})$  is defined as  $\text{val}^*(\mathfrak{G})$  except that the supremum is taken over the set  $C_{qc}(n, k)$  in (8). This in turn yields an explicit correlation that is in the set  $C_{qc}$  but not in  $C_{qa}$ . This game closely resembles the game  $\mathfrak{G}_{\mathcal{M}}$  described in the sketch of the proof that  $\text{RE} \subseteq \text{MIP}^*$ , where  $\mathcal{M}$  is the Turing machine that runs the hierarchy of semidefinite programs on the game  $\mathfrak{G}_{\mathcal{M}}$  and halts if it certifies that  $\text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}}) < 1$ . It is in principle possible to determine an upper bound on the parameters  $n, k$  for our separating correlation from the proof. While we do not provide such a bound, there is no step in the proof that requires it to be astronomical; e.g. we believe (without proof) that  $10^{20}$  is a clear upper bound.

**Connes' Embedding Conjecture.** Connes' Embedding Conjecture (CEC) [Con76] is a conjecture in the theory of von Neumann algebras. Briefly, CEC posits that every type  $\text{II}_1$  von Neumann factor embeds into an ultrapower of the hyperfinite  $\text{II}_1$  factor. We refer to [Oza13] for a precise formulation of the conjecture and connections to other conjectures in operator algebras, such as Kirchberg's QWEP conjecture. In independent work Fritz [Fri12] and Junge et al. [JNP<sup>+</sup>11] showed that a positive answer to CEC would imply

<sup>10</sup>Technically [FNT14] make the observation for the commuting-prover analogue  $\text{MIP}^{\text{co}}(2, 1)$ , discussed further in Section 1.4, but the reasoning is the same.

a positive resolution of Tsirelson’s problem, i.e. that  $C_{qa}(n, k) = C_{qc}(n, k)$  for all  $n, k$ . (This was later promoted to an equivalence by Ozawa [Oza13].) Since our result disproves this equality for some  $n, k$  it also implies that CEC does not hold. In work that appeared subsequently to the first announcement of our results, Goldbrink and Hart [GH16, GH20] show using arguments from continuous logic that the uncomputability of approximating  $\text{val}^*(\mathfrak{G})$  refutes the CEC. Interestingly, their argument uses general elementary considerations from logic and completely bypasses Tsirelson’s problem and its equivalence with Kirchberg’s QWEP conjecture. We note that using the constructive aspect of our result it may be possible to give an explicit description of a factor that does not embed into an ultrapower of the hyperfinite  $\text{II}_1$  factor, but we do not give such a construction.

**Entanglement tests.** As a step towards showing our result for any integer  $n \geq 1$  we construct a game  $\mathfrak{G}_n$ , with question and answer length polynomial in the size of the smallest Turing machine  $\mathcal{M}_n$  that halts (on the empty tape) in exactly  $n$  steps (i.e. the *Kolmogorov complexity* of  $n$ ), such that  $\text{val}^*(\mathfrak{G}_n) = 1$  yet any quantum strategy that succeeds in  $\mathfrak{G}_n$  with probability larger than  $\frac{1}{2}$  must use an entangled state whose Schmidt rank is at least  $2^{\Omega(n)}$ . This is by far the most efficient entanglement test that we are aware of.

**Prover and round reduction for  $\text{MIP}^*$  protocols.** Let  $\text{MIP}^*(k, r)$  denote the collection of languages decidable by  $\text{MIP}^*$  protocols with  $k \geq 2$  provers and  $r$  rounds. Prior to our work it was known how to perform *round reduction* for  $\text{MIP}^*$  protocols, at the cost of adding provers; it was shown by [Ji17, FJVY19] that  $\text{MIP}^*(k, r) \subseteq \text{MIP}^*(k + 15, 1)$  for all  $k, r$ . However, it was an open question whether the complexity of the class  $\text{MIP}^*$  increases if we add more provers. Our main complexity-theoretic result implies that  $\text{MIP}^* = \text{MIP}^*(2, 1)$ . This follows from the following chain of inclusions: for all polynomially-bounded functions  $k, r$ ,

$$\text{MIP}^*(2, 1) \subseteq \text{MIP}^*(k, r) \subseteq \text{RE} \subseteq \text{MIP}^*(2, 1).$$

The first inclusion follows since the verifier in an  $\text{MIP}^*$  protocol can always ignore extra provers and rounds; the second inclusion follows from a simple exhaustive-search procedure that enumerates over strategies for a given  $\text{MIP}^*(k, r)$  protocol; the third result is proven in this paper.<sup>11</sup>

However, this method of reducing provers and rounds in a given  $\text{MIP}^*$  protocol is indirect; it involves first converting a given  $\text{MIP}^*$  protocol into a Turing machine that accepts if and only if the  $\text{MIP}^*$  protocol has value larger than  $\frac{1}{2}$ , and then constructing an  $\text{MIP}^*(2, 1)$  protocol to decide whether the Turing machine halts. In particular this transformation does not generally preserve the complexity of the provers and verifier in the original protocols. We leave it as an open question to find a more direct method for reducing the number of provers in an  $\text{MIP}^*$  protocol.

## 1.4 Open questions

We mention several questions left open by our work.

**Explicit constructions of counter-examples to Connes’ Embedding Conjecture.** We provide an explicit counter-example to Tsirelson’s problem in the form of a game whose entangled value differs from its commuting-operator value. Through the aforementioned connection with Connes’ embedding conjecture [Fri12, JNP<sup>+</sup>11, Oza13], the counter-example may lead to the construction of interesting objects in

---

<sup>11</sup>In fact, we note that the second term  $\text{MIP}^*(k, r)$  can be replaced by  $\text{QMIP}^*(k, r)$ , which is the analogous class with a quantum verifier and quantum messages, since the first inclusion is trivial and the second remains true. As a result, we obtain that  $\text{QMIP}^* = \text{MIP}^*(2, 1)$  as well.

other areas of mathematics. A first question is whether it can lead to an explicit description of a type  $\Pi_1$  factor that does not satisfy the Connes embedding property. Such a construction could be obtained along the lines of [KPS18], using the fact that our game  $\mathfrak{G}$  such that  $\text{val}^*(\mathfrak{G}) < \text{val}^{\text{co}}(\mathfrak{G}) = 1$  has the property of being *synchronous*, i.e. perfect strategies in the game are required to return the same answer when both parties are provided the same question.

Going further, one may ask if the example can eventually lead to a construction of a group that is not sofic, or even not hyperlinear (see e.g. [CLP15] for the connection).

**The complexity of variants of  $\text{MIP}^*$ .** Our result characterizes the complexity class  $\text{MIP}^*$  as the set of recursively enumerable languages. One can also consider the complexity class  $\text{MIP}^{\text{co}}$ , which stands for *multiprover interactive proofs in the commuting-operator model*. For the sake of the discussion we consider only two-prover one-round protocols; a language  $L$  is in  $\text{MIP}^{\text{co}}(2, 1)$  if there exists an efficient reduction that maps  $z \in \{0, 1\}^*$  to a nonlocal game  $\mathfrak{G}_z$  such that if  $z \in L$  then  $\text{val}^{\text{co}}(\mathfrak{G}_z) \geq \frac{2}{3}$ , and otherwise  $\text{val}^{\text{co}}(\mathfrak{G}_z) \leq \frac{1}{3}$ .

The semidefinite programming hierarchy of [NPA08, DLTW08] can be used to show that  $\text{MIP}^{\text{co}}(2, 1)$  is contained in the complement of RE, denoted as  $\text{coRE}$ : to certify that  $z \notin L$  it suffices to run the hierarchy until it obtains a certificate that  $\text{val}^{\text{co}}(\mathfrak{G}_z) < \frac{2}{3}$ . Since it is known that  $\text{RE} \neq \text{coRE}$ ,<sup>12</sup> this implies that  $\text{MIP}^*(2, 1) \neq \text{MIP}^{\text{co}}(2, 1)$ .

It is thus plausible that  $\text{MIP}^{\text{co}} = \text{coRE}$ ,<sup>13</sup> which would provide a very pleasing “dual” complexity characterization to  $\text{MIP}^* = \text{RE}$ . One possible route to proving this would be to adapt our gap-preserving compression framework to the commuting-operator setting by showing that each of the steps (question reduction, answer reduction, and parallel repetition) remain sound against commuting-operator strategies. Using the connection established in [FNT14], this would imply that the operator norm over the maximal  $C^*$  algebra  $C^*(F_2 * F_2)$ , where  $F_2$  is the free group on two elements, is uncomputable.

Another interesting open question concerns the *zero gap* variants of  $\text{MIP}^*$  and  $\text{MIP}^{\text{co}}$ , which we denote by  $\text{MIP}_0^*$  and  $\text{MIP}_0^{\text{co}}$ , respectively. These classes capture the complexity of deciding whether a nonlocal game  $\mathfrak{G}$  has entangled value (or commuting-operator value respectively) *exactly* equal to 1. In [Slo19a], Slofstra shows that there is an efficient reduction from Turing machines  $\mathcal{M}$  to nonlocal games  $\mathfrak{G}_{\mathcal{M}}$  such that  $\mathcal{M}$  does not halt if and only if  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) = \text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}}) = 1$ . This implies that  $\text{coRE} = \text{MIP}_0^{\text{co}}(2, 1)$  and furthermore  $\text{coRE} \subseteq \text{MIP}_0^*$ . However, since  $\text{RE} \subseteq \text{MIP}^*(2, 1) \subseteq \text{MIP}_0^*(2, 1)$ , this implies that  $\text{MIP}_0^*(2, 1)$  is *strictly* bigger than both RE and  $\text{coRE}$ . In fact, it was shown by Mousavi, et al. [MNY20] that the class  $\text{MIP}_0^*(3, 1)$  (the complexity class corresponding to determining whether  $\text{val}^*(\mathfrak{G}) = 1$  for *three-player* nonlocal games) is equal to  $\Pi_2^0$ , which is the set of all languages  $L$  such that  $x \in L$  if and only if  $\forall y, \exists z R(x, y, z) = 1$  for a computable function  $R$  that depends on  $L$ .<sup>14</sup> Thus it is plausible that the complexity landscape of nonlocal games looks like the following:  $\text{MIP}^* = \text{RE}$ ,  $\text{MIP}_0^* = \Pi_2^0$ , and  $\text{MIP}^{\text{co}} = \text{MIP}_0^{\text{co}} = \text{coRE}$ . Such statements about the complexity of  $\text{MIP}^*$  versus  $\text{MIP}^{\text{co}}$ , in both the gapped and zero-gap cases, may reveal additional insights into the difference between the tensor product and commuting-operator models of correlations.

**Acknowledgments.** We thank Matthew Coudron, William Slofstra and Jalex Stark for enlightening discussions regarding possible consequences of our work. We thank William Slofstra and Jalex Stark for

<sup>12</sup> $\text{RE} \neq \text{coRE}$  follows from the fact that  $\text{RE} \cap \text{coRE}$  is the set of decidable languages and RE contains undecidable languages.

<sup>13</sup>We note that the “co” modifier on both sides of the equation  $\text{MIP}^{\text{co}} = \text{coRE}$  refer to different things!

<sup>14</sup>The class  $\Pi_2^0$  is also characterized as being part of the second level of the *arithmetical hierarchy* from computability theory, where  $\text{RE} = \Sigma_1^0$  and  $\text{coRE} = \Pi_1^0$  form the first level.

suggestions regarding explicit separations between  $C_{qa}$  and  $C_{qc}$ . We thank Peter Burton, William Slofstra and Jalex Stark for comments on a previous version. We thank Lewis Bowen and Mikael de la Salle for pointing out typos and minor errors in a previous version.

Zhengfeng Ji is supported by Australian Research Council (DP200100950). Anand Natarajan is supported by IQIM, an NSF Physics Frontiers Center (NSF Grant PHY-1733907). Thomas Vidick is supported by NSF CAREER Grant CCF-1553477, AFOSR YIP award number FA9550-16-1-0495, a CIFAR Azrieli Global Scholar award, MURI Grant FA9550-18-1-0161 and the IQIM, an NSF Physics Frontiers Center (NSF Grant PHY-1125565) with support of the Gordon and Betty Moore Foundation (GBMF-12500028). Henry Yuen is supported by NSERC Discovery Grant 2019-06636. Part of this work was done while John Wright was at the Massachusetts Institute of Technology. He is supported by IQIM, an NSF Physics Frontiers Center (NSF Grant PHY-1733907), and by ARO contract W911NF-17-1-0433.



## 2 Proof Overview

In this section we give an overview of the proof of the inclusion  $\text{RE} \subseteq \text{MIP}^*$ . Since all interactive proof systems considered in the paper involve a single-round interaction between a classical verifier and two quantum provers sharing entanglement we generally use the language of nonlocal games to describe such proof systems, and often refer to the provers as “players”. In a nonlocal game  $\mathfrak{G}$  (or simply “game” for short), the verifier can be described as the combination of two procedures: a *question sampling* procedure that samples a pair of questions  $(x, y)$  for the players according to a distribution  $\mu$  (known to the verifier and the players), and a *decision* procedure (also known to all parties) that takes as input the players’ questions and their respective answers  $a, b$  and evaluates a predicate  $D(x, y, a, b) \in \{0, 1\}$  to determine the verifier’s acceptance or rejection. Given a description of a nonlocal game  $\mathfrak{G}$ , recall that  $\text{val}^*(\mathfrak{G})$  denotes the *entangled value* of the game, which is defined as the supremum (6) of the players’ success probability in the game over all finite-dimensional tensor product strategies. (We refer to Section 5 for definitions regarding nonlocal games.)

Our results establish the existence of transformations on *families* of nonlocal games  $\{\mathfrak{G}_n\}_{n \in \mathbb{N}}$  having certain properties. In order to keep track of efficiency (and ultimately, computability) properties it is important to have a way to specify such families in a uniform manner. Towards this we introduce the following formalism. A *uniformly generated family of games* is specified through a pair of Turing machines  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  that satisfy certain conditions, in which case the pair is called a *normal form verifier*. The Turing machine  $\mathcal{S}$  (called a *sampler*) takes as input an index  $n \in \mathbb{N}$  and returns the description of a procedure that can be used to sample questions  $(x, y)$  in the game (this procedure itself obeys a certain format associated with “conditionally linear” distributions, defined below). The Turing machine  $\mathcal{D}$  (called a *decider*) takes as input an index  $n$ , questions  $(x, y)$ , and answers  $(a, b)$ , and returns a single-bit decision. For the sake of this proof overview we assume that the sampling and decision procedures run in time polynomial in the index  $n$ ; we refer to the running time of these procedures as the *complexity* of the verifier. Given a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  we associate to it an infinite family of nonlocal games  $\{\mathcal{V}_n\}$  indexed by positive integers in the natural way.

The main technical result of this paper is a *gap-preserving compression* transformation on normal form verifiers. The following theorem presents an informal summary of the properties of this transformation. Recall that for a game  $\mathfrak{G}$  and probability  $0 \leq p \leq 1$ ,  $\mathcal{E}(\mathfrak{G}, p)$  denotes the minimum local dimension of an entangled state shared by the players in order for them to succeed in  $\mathfrak{G}$  with probability at least  $p$ .

**Theorem 2.1** (Gap-preserving compression of normal form verifiers, informal). *There exists a polynomial-time Turing machine Compress that, when given as input the description of a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ , outputs the description of another normal form verifier  $\mathcal{V}' = (\mathcal{S}', \mathcal{D}')$  that satisfies the following properties: for all  $n \in \mathbb{N}$ , letting  $N = 2^n$ ,*

1. (**Completeness**) *If  $\text{val}^*(\mathcal{V}_N) = 1$  then  $\text{val}^*(\mathcal{V}'_n) = 1$ .*
2. (**Soundness**) *If  $\text{val}^*(\mathcal{V}_N) \leq \frac{1}{2}$  then  $\text{val}^*(\mathcal{V}'_n) \leq \frac{1}{2}$ .*
3. (**Entanglement lower bound**)  *$\mathcal{E}(\mathcal{V}'_n, \frac{1}{2}) \geq \max\{\mathcal{E}(\mathcal{V}_N, \frac{1}{2}), 2^{2^{\Omega(n)}}\}$ .*

The formal version of this theorem is stated in Section 12 as Theorem 12.1. The terminology *compression* is motivated by the fact, implicit in the informal statement of the theorem, that the time complexity of the verifier’s sampling and decision procedures in the game  $\mathcal{V}'_n$ , which is polynomial in  $n$ , is exponentially smaller than the time complexity of the verifier in the game  $\mathcal{V}_N$ , which is polynomial in  $N$  and thus exponential in  $n$ .

Before giving an overview of the proof of Theorem 2.1 we sketch how the existence of a Turing machine Compress with the properties stated in the theorem implies the inclusion  $\text{RE} \subseteq \text{MIP}^*$ . Recall that the complexity class RE consists of all languages  $L$  such that there is a Turing machine  $\mathcal{M}$  that accepts instances  $x$  in  $L$ , and does not accept instances  $x$  that are not in  $L$  (but is not required to terminate on such instances). To show  $\text{RE} \subseteq \text{MIP}^*$  we give an  $\text{MIP}^*$  protocol for the Halting Problem, which is a complete problem for RE. The Halting Problem is the language that consists of all Turing machine descriptions  $\mathcal{M}$  such that  $\mathcal{M}$  halts when run on an empty input tape. (For the purposes of this overview, we blur the distinction between a Turing machine and its description as a string of bits.) We give a procedure that given a Turing machine  $\mathcal{M}$  as input returns the description of a normal form verifier  $\mathcal{V}^{\mathcal{M}} = (\mathcal{S}^{\mathcal{M}}, \mathcal{D}^{\mathcal{M}})$  with the following properties. First, if  $\mathcal{M}$  does eventually halt on an empty input tape, then it holds that for all  $n \in \mathbb{N}$ ,  $\text{val}^*(\mathcal{V}_n^{\mathcal{M}}) = 1$ . Second, if  $\mathcal{M}$  does not halt then for all  $n \in \mathbb{N}$ ,  $\text{val}^*(\mathcal{V}_n^{\mathcal{M}}) \leq \frac{1}{2}$ .

We describe the procedure that achieves this. Informally, the procedure returns the specification of a verifier  $\mathcal{V}^{\mathcal{M}} = (\mathcal{S}^{\mathcal{M}}, \mathcal{D}^{\mathcal{M}})$  such that  $\mathcal{D}^{\mathcal{M}}$  proceeds as follows: on input  $(n, x, y, a, b)$  it first executes the Turing machine  $\mathcal{M}$  for  $n$  steps. If  $\mathcal{M}$  halts, then  $\mathcal{D}^{\mathcal{M}}$  accepts. Otherwise,  $\mathcal{D}^{\mathcal{M}}$  computes the description of the compressed verifier  $\mathcal{V}' = (\mathcal{S}', \mathcal{D}')$  that is the output of Compress on input  $\mathcal{V}^{\mathcal{M}}$ , then executes the decision procedure  $\mathcal{D}'(n, x, y, a, b)$  and accepts if and only if  $\mathcal{D}'$  accepts.<sup>15</sup>

To show that this procedure achieves the claimed transformation, consider two cases. First, observe that if  $\mathcal{M}$  eventually halts in some number of time steps  $T$ , then by definition  $\text{val}^*(\mathcal{V}_n^{\mathcal{M}}) = 1$  for all  $n \geq T$ . Using Theorem 2.1 along with an inductive argument it then follows that  $\text{val}^*(\mathcal{V}_n^{\mathcal{M}}) = 1$  for all  $n \geq 1$ . Second, if  $\mathcal{M}$  never halts, then observe that for any  $n \geq 1$  Theorem 2.1 implies two separate lower bounds on the amount of entanglement required to win the game  $\mathcal{V}_n^{\mathcal{M}}$  with probability at least  $\frac{1}{2}$ : the dimension is (a) at least  $2^{2^{\Omega(n)}}$ , and (b) at least the dimension needed to win the game  $\mathcal{V}_{2^n}^{\mathcal{M}}$  with probability at least  $\frac{1}{2}$ . Applying an inductive argument it follows that an *infinite* amount of entanglement is needed to win the game  $\mathcal{V}_n$  with any probability greater than  $\frac{1}{2}$ . Thus, a sequence of finite-dimension strategies for  $\mathcal{V}_n$  cannot lead to a limiting value larger than  $\frac{1}{2}$ , and  $\text{val}^*(\mathcal{V}_n^{\mathcal{M}}) \leq \frac{1}{2}$ .

We continue with an overview of the ideas behind the proof of Theorem 2.1.

**Compression by introspection.** To start, it is useful to review the protocol introduced in [NW19] to show the inclusion  $\text{NEEXP} \subseteq \text{MIP}^*$ . Fix an NEEXP-complete language  $L$ . The  $\text{MIP}^*$  protocol for NEXP from [NV18b], when scaled up to decide languages from NEEXP, yields a family of nonlocal games  $\{\mathfrak{G}_z\}$  that are indexed by instances  $z \in \{0, 1\}^*$ . The family of games decides  $L$  in the sense that for all  $z$ , the game  $\mathfrak{G}_z$  has entangled value 1 if  $z \in L$ , and has entangled value at most  $\frac{1}{2}$  if  $z \notin L$ . Furthermore, if  $n = |z|$  is the length of  $z$ , the verifier of the game  $\mathfrak{G}_z$  has complexity  $\text{poly}(N) = \exp(|z|)$  (recall that we use this terminology to refer to an upper bound on the running time of the verifier's sampling and decision procedure). Thus, this family of games does not by itself yield an  $\text{MIP}^*$  protocol for  $L$ . To overcome this the main contribution in [NW19] is the design of an efficient compression procedure  $\text{Compress}^{\text{NW}}$  that applies specifically to the family of games  $\{\mathfrak{G}_z\}$ . When given as input the description of  $\mathfrak{G}_z$ ,  $\text{Compress}^{\text{NW}}$  returns the description of a game  $\mathfrak{G}'_z$  such that if  $\text{val}^*(\mathfrak{G}_z) = 1$ , then  $\text{val}^*(\mathfrak{G}'_z) = 1$ , and if  $\text{val}^*(\mathfrak{G}_z) \leq \frac{1}{2}$ , then  $\text{val}^*(\mathfrak{G}'_z) \leq \frac{1}{2}$ . Furthermore, the complexity of the verifier for  $\mathfrak{G}'_z$  is  $\text{poly}(n)$ . Thus the family of games  $\{\mathfrak{G}'_z\}$  decides  $L$  and this shows that  $\text{NEEXP} \subseteq \text{MIP}^*$ , which is the best lower bound known on  $\text{MIP}^*$  prior to our work.

Presented in this way, it is natural to suggest iterating the procedure  $\text{Compress}^{\text{NW}}$  to achieve e.g. the inclusion  $\text{NEEXP} \subseteq \text{MIP}^*$ . To explain the difficulty in doing so, we give a little more detail on the

<sup>15</sup>The fact that the decider  $\mathcal{D}^{\mathcal{M}}$  can invoke the Compress procedure on itself follows from a well-known result in computability theory known as *Kleene's recursion theorem* (also called *Roger's fixed point theorem*) [Kle54, Rog87].

compression procedure. It consists of two main steps: starting from  $\mathfrak{G}_z$ , perform (1) *question reduction*, and (2) *answer reduction*. The goal of (1) is to reduce the length of the questions generated by the verifier in  $\mathfrak{G}_z$  from  $\text{poly}(N)$  to  $\text{poly}(n)$ . The goal of (2) is to achieve the same with respect to the length of answers expected from the players. Furthermore, the complexity of the verifier of the resulting game  $\mathfrak{G}'_z$  should be reduced from  $\text{poly}(N)$  to  $\text{poly}(n)$ .

Part (1) is achieved through a technique referred to as “introspection” where, rather than sampling questions  $(x, y)$  of length  $\text{poly}(N)$  as in the game  $\mathfrak{G}_z$ , the verifier instead executes a carefully crafted nonlocal game with the players that (a) requires questions of length  $\text{poly}(n)$ , (b) checks that the players share  $\text{poly}(N)$  EPR pairs, and (c) checks that the players measure the EPR pairs in such a way as to *sample for themselves* a question pair  $(x, y)$  such that one player gets  $x$  and the other player gets  $y$ . In other words, the players are essentially forced to introspectively ask themselves the questions of  $\mathfrak{G}_z$ .

After question reduction, the players still respond with  $\text{poly}(N)$ -length answers, which the verifier has to check satisfies the decision predicate of the original game  $\mathfrak{G}_z$ . The goal of Part (2) is to enable the decision procedure to implement the verification procedure while not requiring the entire full-length answers from the players. In the answer reduction scheme of [NW19] this is achieved by having the verifier run a *probabilistically checkable proof* (PCP) with the players so that they succinctly *prove* that first, they have introspected questions  $(x, y)$  from the correct distribution, and second, that they are able to generate  $\text{poly}(N)$ -length answers  $(a, b)$  that would satisfy the decision predicate of the original game  $\mathfrak{G}_z$  when executed on  $(x, y)$  and  $(a, b)$ . Since the questions and answers in the PCP are of length  $\text{poly}(n)$ , this achieves the desired answer length reduction.

Iterating this scheme presents a number of immediate difficulties that have to do with the fact that the sampling and decision procedures of the verifier in  $\mathfrak{G}'_z$  do not have such a nice form as those in  $\mathfrak{G}_z$ . First of all, the compression procedure of [NW19] can only “introspect” a specific question distribution of a nonlocal game from [NV18b]; we call this distribution a “low-degree test distribution”.<sup>16</sup> However, the resulting question distribution of the question-reduced verifier, which is used to check the introspection, has a much more complex structure. A similar issue arises with the modifications required to perform answer reduction. In the PCP employed to achieve this the question distribution appears to be much more complex than the low-degree test distribution (this is in large part due to the need for a specially tailored PCP procedure that encodes separately different chunks of the witness, corresponding to answers from different players). As a result it is entirely unclear at first whether the question distribution used by the verifier in  $\mathfrak{G}'_z$  can be “introspected” for a second time.

To overcome these difficulties we identify a natural class of question distributions, called *conditionally linear distributions*, that generalize the low-degree test distribution. We show that conditionally linear distributions can be “introspected” using conditionally linear distributions only, enabling recursive introspection. (In particular, they are a rich enough class to capture the types of question distributions produced by the compression scheme of [NW19].) We define normal form verifiers by restricting their sampling procedure to generate conditionally linear question distributions, and this allows us to obtain the compression procedure on normal form verifiers described in Theorem 2.1.

Conceptually, the identification of a natural class of distributions that is “closed under introspection” is a key step that enables the introspection technique to be applied recursively. (As we will see later, other

---

<sup>16</sup>The nonlocal game of [NV18b] is part of a more general family of games called “low-degree tests”, which have been studied extensively for their applications in complexity theory [BFL91], property testing [RS96], and PCPs [AS98]. Generally, the question distribution of a low-degree test is a random question pair  $(x, y)$  where  $y$  is a randomly chosen affine subspace in  $\mathbb{F}^m$  and  $x$  is a uniformly random point on  $y$ , where  $\mathbb{F}$  is a finite field and  $m \geq 2$  is an integer. The specific low-degree test game in [NV18b] (which is based on the low-degree test of [RS97]) uses two-dimensional subspaces; thus the question distribution of [RS97, NV18b] is referred to as the “plane-point distribution”.

closure properties of conditionally linear distributions, such as taking direct products, play an important role as well.) Since conditionally linear distributions are central to our construction we describe them next.

**Conditionally linear distributions.** Fix a vector space  $V$  that is identified with  $\mathbb{F}^m$ , for a finite field  $\mathbb{F}$  and integer  $m$ . Informally (see Definition 4.1 for a precise definition), a function  $L$  on  $V$  is *conditionally linear* (CL for short) if it can be evaluated by a procedure that takes the following form: (i) read a substring  $z^{(1)}$  of  $z$ ; (ii) evaluate a linear function  $L_1$  on  $z^{(1)}$ ; (iii) repeat steps (i) and (ii) with the remaining coordinates  $z \setminus z^{(1)}$ , such that the next steps are allowed to depend in an arbitrary way on  $L_1(z^{(1)})$  but not directly on  $z^{(1)}$  itself. What distinguishes a function of this form from an arbitrary function is that we restrict the number of iterations of (i)—(ii) to a constant number (at most 9, in our case). (One may also think of CL functions as “adaptively linear” functions, where the number of “levels” of adaptivity is the number of iterations of (i)—(ii).) A distribution  $\mu$  over pairs  $(x, y) \in V \times V$  is called conditionally linear if it is the image under a pair of conditionally linear functions  $L^A, L^B : V \rightarrow V$  of the uniform distribution on  $V$ , i.e.  $(x, y) \sim (L^A(z), L^B(z))$  for uniformly random  $z \in V$ .

An important class of CL distributions are low-degree test distributions, which are distributions over question pairs  $(x, y)$  where  $y$  is a randomly chosen affine subspace of  $\mathbb{F}^m$  and  $x$  is a uniformly random point on  $y$ . We explain this for the case where the random subspace  $y$  is one-dimensional (i.e. a line). Let  $V = V_X \oplus V_V$  where  $V_X = V_V = \mathbb{F}^m$ . Let  $L^A$  be the projection onto  $V_X$  (i.e. it maps  $(x, v) \rightarrow (x, 0)$  where  $x \in V_X$  and  $v \in V_V$ ). Define  $L^B : V \rightarrow V$  as the map  $(x, v) \mapsto (L_v^{L^N}(x), v)$  where  $L_v^{L^N} : V_X \rightarrow V_X$  is a linear map that, for every  $v \in V_V$ , projects onto a complementary subspace to the one-dimensional subspace of  $V_X$  spanned by  $v$  (one can think of this as an “orthogonal subspace” to the span of  $\{v\}$ ).  $L^B$  is conditionally linear because it can be seen as first reading the substring  $v \in V_V$  (which can be interpreted as specifying the *direction* of a line), and then applying a linear map  $L_v^{L^N}$  to  $x \in V_X$  (which can be interpreted as specifying a canonical point on the line  $\ell = \{x + tv : t \in \mathbb{F}\}$ ). It is not hard to see (and shown formally in Section 7.1.1) that the distribution of  $(L^A(z), L^B(z))$  for  $z$  uniform in  $V$ , is identical (up to relabeling) to the low-degree test distribution  $(x, \ell)$  where  $\ell$  is a uniformly random affine line in  $\mathbb{F}^m$ , and  $x$  is a uniformly random point on  $\ell$ .

Our main result about CL distributions, presented in Section 8, is that any CL distribution  $\mu$ , associated with a pair of CL functions  $(L^A, L^B)$  over a linear space  $V = \mathbb{F}^m$ , can be “introspected” using a CL distribution that is “exponentially smaller” than the initial distribution. Slightly more formally, to any CL distribution  $\mu$  we associate a two-player game  $\mathfrak{G}_\mu$  (called the “introspection game”) in which questions from the verifier are sampled from a CL distribution  $\mu'$  over  $\mathbb{F}^{m'}$  for some  $m' = \text{poly log}(m)$  and such that in any successful strategy for the game  $\mathfrak{G}_\mu$ , when the players are queried on a special question labeled INTRO, they must respond with a pair  $(x, y)$  that is approximately distributed according to  $\mu$ . (The game allows us to do more: it allows us to conclude *how* the players obtained  $(x, y)$  — by measuring shared EPR pairs in a specific basis — and this will be important when using the game as part of a larger protocol that involves other checks.) Crucially for us, the distribution  $\mu'$  only depends on a size parameter associated with  $(L^A, L^B)$  (essentially, the integer  $m$  together with the number of “levels” of adaptivity of  $L^A$  and  $L^B$ ), but not on any other structural property of  $(L^A, L^B)$ . Only the decision predicate for the introspection game  $\mathfrak{G}_\mu$  depends on the entire description of  $(L^A, L^B)$ .

We say a few words about the design of  $\mu'$  and the associated introspection game, which borrow heavily from [NW19]. Building on the “quantum low-degree test” introduced in [NV18a] it is already known how a verifier can force a pair of players to measure  $m$  EPR pairs in either the computational or Hadamard basis and report the (necessarily identical) outcome  $z$  obtained, all the while using questions of length polylogarithmic in  $m$  only. The added difficulty is to ensure that a player obtains, and returns, precisely the information about

$z$  that is contained in  $L^A(z)$  (resp.  $L^B(z)$ ), and not more. A simple example is the line-point distribution described earlier: there, the idea to ensure that e.g. the “point” player only obtains the first component,  $x$  of  $(x, v) \in V_X \oplus V_V$ , the verifier demands that the “point” player measures their qubits associated with the space  $V_V$  in the Hadamard, instead of computational, basis; due to the uncertainty principle this has the effect of “erasing” the outcome in the computational basis. The case of the “line” player is a little more complex: the goal is to ensure that, conditioned on the specification of the line  $\ell$  received by the “line” player, the point  $x$  received by the “point” player is uniformly random within  $\ell$ . This was shown to be possible in [NW19].

We can now describe how samplers of normal form verifiers are defined: these are Turing machines  $\mathcal{S}$  that specify an infinite family of CL distributions  $\{\mu_n\}$  by computing, for each index  $n$ , the CL functions  $(L^{A,n}, L^{B,n})$  associated with  $\mu_n$ . (See Definition 4.14 for a formal definition of samplers.) Thus, the question distributions of a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  are the CL distributions corresponding to  $\mathcal{S}$ .

**Question reduction.** Just like the compression procedure of [NW19], the compression procedure `Compress` of Theorem 2.1 begins with performing question reduction on the input game. Given a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ , the procedure `Compress` first computes a normal form verifier  $\mathcal{V}^{\text{INTRO}} = (\mathcal{S}^{\text{INTRO}}, \mathcal{D}^{\text{INTRO}})$  where for all  $n \in \mathbb{N}$ , the game  $\mathcal{V}_n^{\text{INTRO}}$  consists of playing the original game  $\mathcal{V}_N$  where  $N = 2^n$ , except that instead of sampling the questions according to the CL distribution  $\mu_N$  specified by the sampler  $\mathcal{S}_N$ , the verifier executes the introspection game  $\mathfrak{G}_{\mu_N}$  described in the previous subsection. Thus, in the game  $\mathcal{V}_n^{\text{INTRO}}$ , when both players receive the question labeled `INTRO` they are expected to sample  $(x, y)$  respectively according to  $\mu_N$ , and respond with the sampled question together with answers  $a, b$  respectively. The decider  $\mathcal{D}^{\text{INTRO}}$  on index  $n$  evaluates  $\mathcal{D}(N, x, y, a, b)$  and accepts if and only if  $\mathcal{D}$  accepts. As a result the time complexity of decider  $\mathcal{D}^{\text{INTRO}}$  on index  $n$  remains that of  $\mathcal{D}$ , i.e.  $\text{poly}(N)$ . However, the length of questions asked in  $\mathcal{V}_n^{\text{INTRO}}$  and the complexity of the sampler  $\mathcal{S}^{\text{INTRO}}$  are exponentially reduced, to  $\text{poly}(n)$ .

For convenience we refer to the questions asked by the verifier in the “question-reduced” game  $\mathcal{V}_n^{\text{INTRO}}$  as “small questions,” and the questions that are introspected by the players in  $\mathcal{V}_n^{\text{INTRO}}$  (equivalently, the questions asked in the original game  $\mathcal{V}_N$ ) as “big questions.”

**Answer reduction.** Having reduced the complexity of the question sampling, the next step in the compression procedure `Compress` is to reduce the complexity of decider  $\mathcal{D}^{\text{INTRO}}$  from  $\text{poly}(N)$  to  $\text{poly}(n)$  (which necessarily implies reducing the answer length to  $\text{poly}(n)$ ). To achieve this the compression procedure computes a normal form verifier  $\mathcal{V}^{\text{AR}} = (\mathcal{S}^{\text{AR}}, \mathcal{D}^{\text{AR}})$  from  $\mathcal{V}^{\text{INTRO}}$  such that both the sampler and decider complexity in  $\mathcal{V}^{\text{AR}}$  are  $\text{poly}(n)$  (here, `AR` stands for “answer reduction”).

Similarly to the answer reduction performed in [NW19], at a high level this is achieved by composing the game  $\mathcal{V}_n^{\text{INTRO}}$  with a probabilistically checkable proof (PCP). In our context a PCP is a proof encoding that allows a verifier to check whether, given Turing machine  $\mathcal{A}$  and time bound  $T$  provided as input, there exists some input  $x$  that  $\mathcal{A}$  accepts in time  $T$ . The PCP proof can be computed from  $\mathcal{A}$ ,  $T$ , and the accepting input (if it exists) and has length polynomial in  $T$  and the description length  $|\mathcal{A}|$  of  $\mathcal{A}$ . Crucially, the verifier can check a purported proof while only reading a constant number of symbols of it, each of length  $\text{polylog}(T, |\mathcal{A}|)$ , and executing a verification procedure that runs in time  $\text{polylog}(T, |\mathcal{A}|)$ .

We use PCPs for answer reduction as follows. The verifier in the game  $\mathcal{V}_n^{\text{AR}}$  samples questions as  $\mathcal{V}_n^{\text{INTRO}}$  would and sends them to the players. Instead of receiving the introspected questions and answers  $(x, y, a, b)$  for the original game  $\mathcal{V}_N$  and running the decision procedure  $\mathcal{D}(N, x, y, a, b)$ , the verifier instead asks the players to compute a PCP  $\Pi$  for the statement that the original decider  $\mathcal{D}$  accepts the input  $(N, x, y, a, b)$  in time  $T = \text{poly}(N)$ . The verifier then samples additional questions for the players that ask them to return

specific entries of the proof  $\Pi$ . Finally, upon receipt of the players' answers, the verifier executes the PCP verification procedure. Because of the efficiency of the PCP, both the sampling of the additional questions and the decision procedure can be executed in time  $\text{poly}(n)$ .<sup>17</sup>

This very rough sketch presents some immediate difficulties. A first difficulty is that in general no player by themselves has access to the entire input  $(N, x, y, a, b)$  to  $\mathcal{D}$ , so no player can compute the entire proof  $\Pi$ . We discuss this issue in the next paragraph. A second difficulty is that a black-box application of an existing PCP, as done in [NW19], results in a question distribution for  $\mathcal{V}_n^{\text{AR}}$  (i.e. the sampling of the proof locations to be queried) that is rather complex — and in particular, it may no longer fall within the framework of CL distributions for which we can do introspection. To avoid this, we design a bespoke PCP based on the classical MIP for NEXP (in particular, we borrow and adapt techniques from [BSS05, BSGH<sup>+</sup>06]). Two essential properties for us are that (i) the PCP proof is a collection of several low-degree polynomials, two of which are low-degree encodings of each player's big answer in the game  $\mathcal{V}_n^{\text{INTRO}}$ , and (ii) verifying the proof only requires (a) running low-degree tests, (b) querying all polynomials at a uniformly random point, and (c) performing simple consistency checks. Property (i) allows us to eliminate the extra layer of encoding in [NW19], who had to consider a PCP of proximity for a circuit applied to the low-degree encodings of the players' big answers. Property (ii) allows us to ensure that the question distribution employed by  $\mathcal{V}_n^{\text{AR}}$  remains conditionally linear.

**Oracularization.** The preceding paragraph raises a non-trivial difficulty. In order for the players to compute a proof for the claim that  $\mathcal{D}(N, x, y, a, b) = 1$  they need to know the entire input  $(x, y, a, b)$ . However, in general a player only has access to their own question and answer: one player only knows  $(x, a)$  and the other player knows  $(y, b)$ . The standard way of circumventing this difficulty is to consider an “oracularized” version of the game, where one player gets both questions  $(x, y)$  and is able to determine both answers  $(a, b)$ , while the other player only gets one of the questions at random, and is only asked for one of the answers, that is then checked for consistency with the first player's answer.

While this technique works well for games with classical players, when the players are allowed to use quantum strategies using entanglement oracularization does not, in general, preserve the completeness property of the game. To ensure that completeness is preserved we need an additional property of a completeness-achieving strategy for the original game: that there exists a *commuting and consistent strategy* on all pairs of questions  $(x, y)$  that are asked in the game with positive probability. Here commuting means that the measurement  $\{A_a^x\}_a$  performed by the player receiving  $x$  commutes with the measurement  $\{B_b^y\}_b$  performed by the player receiving  $y$ .<sup>18</sup> Consistent means that if both players perform measurements associated with the same question they obtain the same answer. If both properties hold then in the oracularized game when one player receives a pair  $(x, y)$  and the other player receives the question  $x$  (say), the first player can simultaneously measure both  $\{A_a^x\}_a$  and  $\{B_b^y\}_b$  on their own space to obtain a pair of answers  $(a, b)$ , and the second player can measure  $\{A_a^x\}_a$  to obtain a consistent answer  $a$ .

For answer reduction to be possible it is thus applied to the oracularized version of the introspection game  $\mathcal{V}_n^{\text{INTRO}}$ . This in turn requires us to ensure that the introspection game  $\mathcal{V}_n^{\text{INTRO}}$  has a commuting and consistent strategy achieving value 1 whenever it is the case that  $\text{val}^*(\mathcal{V}_n^{\text{INTRO}}) = 1$ . For this property to hold we verify that it holds for the initial game that is used to seed the compression procedure (this is true because we can start with an MIP\* protocol for NEXP for which there exists a perfect classical strategy)

<sup>17</sup>This idea is inspired by the technique of composition in the PCP literature, in which the complexity of a verification procedure can be reduced by composing a proof system (often a PCP itself) with another PCP.

<sup>18</sup>We stress that the commuting property only applies to question pairs that occur with positive probability, and does not mean that *all* pairs of measurement operators are required to commute; indeed this would imply that the strategy is effectively classical.

and we also ensure that each of the transformations of the compression protocol (question reduction, answer reduction, and parallel repetition described next) maintains it.

**Parallel repetition.** The combined steps of question reduction (via introspection) and answer reduction (via PCP composition) result in a game  $\mathcal{V}_n^{\text{AR}}$  such that the complexity of the verifier is  $\text{poly}(n)$ . Furthermore, if the original game  $\mathcal{V}_N$  has value 1, then  $\mathcal{V}_n^{\text{AR}}$  also has value 1. Unfortunately the sequence of transformations incurs a loss in the soundness parameters: if  $\text{val}^*(\mathcal{V}_N) \leq \frac{1}{2}$ , then we can only establish that  $\text{val}^*(\mathcal{V}_n^{\text{AR}}) \leq 1 - C$  for some positive constant  $C < \frac{1}{2}$  (we call  $C$  the *soundness gap*). Such a loss would prevent us from recursively applying the compression procedure Compress an arbitrary number of times, which is needed to obtain the desired complexity results for  $\text{MIP}^*$ .

To overcome this we need a final transformation to restore the soundness gap of the games after answer reduction to a constant larger than  $\frac{1}{2}$ . To achieve this we use the technique of parallel repetition. The parallel repetition of a game  $\mathcal{G}$  is another nonlocal game  $\mathcal{G}^k$ , for some number of repetitions  $k$ , which consists of playing  $k$  independent and simultaneous instances of  $\mathcal{G}$  and accepting if and only if all  $k$  instances accept. Intuitively, parallel repetition is meant to decrease the value of a game  $\mathcal{G}$  exponentially fast in  $k$ , provided  $\text{val}^*(\mathcal{G}) < 1$  to begin with. However, it is an open question of whether this is generally true for the entangled value  $\text{val}^*$ .

Nevertheless, some variants of parallel repetition are known to achieve exponential amplification. We use a variant called “anchored parallel repetition” and introduced in [BYY17]. This allows us to devise a transformation that efficiently amplifies the soundness gap to a constant. The resulting game  $\mathcal{V}_n^{\text{REP}}$  has the property that if  $\text{val}^*(\mathcal{V}_n^{\text{AR}}) = 1$ , then  $\text{val}^*(\mathcal{V}_n^{\text{REP}}) = 1$  (and moreover this is achieved using a commuting and consistent strategy), whereas if  $\text{val}^*(\mathcal{V}_n^{\text{AR}}) \leq 1 - C$  for some universal constant  $C > 0$  then  $\text{val}^*(\mathcal{V}_n^{\text{REP}}) \leq \frac{1}{2}$ . Furthermore, we have the additional property, essential for us, that good strategies in the game  $\mathcal{V}_n^{\text{REP}}$  require as much entanglement as good strategies in the game  $\mathcal{V}_n^{\text{AR}}$  (which in turn require as much entanglement as good strategies in the game  $\mathcal{V}_N$ ). The complexity of the verifier in  $\mathcal{V}_n^{\text{REP}}$  remains  $\text{poly}(n)$ .

The anchored parallel repetition procedure, when applied to a normal form verifier, also yields a normal form verifier: this is because the direct product of CL distributions is still conditionally linear.

**Putting it all together.** This completes the overview of the transformations performed by the compression procedure Compress of Theorem 2.1. To summarize, given an input normal form verifier  $\mathcal{V}$ , question reduction is applied to obtain  $\mathcal{V}^{\text{INTRO}}$ , answer reduction is applied to the oracularized version of  $\mathcal{V}^{\text{INTRO}}$  to obtain  $\mathcal{V}^{\text{AR}}$ , and anchored parallel repetition is applied to obtain  $\mathcal{V}^{\text{REP}}$ , which is returned by the compression procedure. Each of these transformations preserves completeness (including the commuting and consistent properties of a value-1 strategy) as well as the entanglement requirements of each game; moreover, the overall transformation preserves soundness.

### 3 Preliminaries

**Notation.** We use  $\Sigma$  to denote a finite alphabet.  $\mathbb{N}$  is the set of positive integers. For  $w \in \{0, 1\}$ ,  $\bar{w}$  denotes  $1 - w$ . For  $w \in \{A, B\}$ ,  $\bar{w} = B$  if  $w = A$  and  $\bar{w} = A$  otherwise. (For notational convenience we often implicitly make the identifications  $1 \leftrightarrow A$  and  $2 \leftrightarrow B$ .) We use  $\mathbb{F}$  to denote a finite field. We write  $M_n(\mathbb{F})$  to denote the set of  $n \times n$  matrices over  $\mathbb{F}$ . We write  $I$  to denote the identity operator on a vector space. We write  $\text{Tr}(\cdot)$  for the matrix trace. We write  $\mathcal{H}$  to denote a separable Hilbert space. For a linear operator  $T$ ,  $\|T\|$  denotes the operator norm.

**Asymptotics.** All logarithms are base 2. We use the notation  $O(\cdot)$ ,  $\text{poly}(\cdot)$ , and  $\text{polylog}(\cdot)$  in the following way. For  $f, g : \mathbb{N} \rightarrow \mathbb{R}_+$  we write  $f(n) = O(g(n))$  (omitting the integer  $n$  when it is clear from context) to mean that there exists a constant  $C > 0$  such that for all  $n \in \mathbb{N}$ ,  $f(n) \leq Cg(n)$ . When we write  $f(a_1, \dots, a_k) = \text{poly}(a_1, \dots, a_k)$ , this indicates that there exists a universal constant  $C > 0$  (which may vary each time the notation is used in the paper) such that  $f(a_1, \dots, a_k) \leq C(a_1 \cdots a_k)^C$  for all positive  $a_1, \dots, a_k$ . Similarly, when we write  $f(a_1, \dots, a_k) = \text{polylog}(a_1, \dots, a_k)$ , there exists a universal constant  $C$  such that  $f(a_1, \dots, a_k) \leq C \prod_{i=1}^k \log^C(1 + a_i)$  for all positive  $a_1, \dots, a_k$ . Finally, we write  $\log(a_1, \dots, a_k)$  as short hand for  $\prod_{i=1}^k \log(1 + a_i)$ .<sup>19</sup>

#### 3.1 Turing machines

Turing machines are a model of computation introduced in [Tur37], and play a central role in our modeling of verifiers for nonlocal games. Here we give an overview of certain aspects of Turing machines that are relevant for this paper. For an in-depth treatment of Turing machines, we refer the reader to the textbooks of Papadimitriou [Pap94] or Sipser [Sip12].

The tapes of a Turing machine are infinite one-dimensional arrays of cells that are indexed by natural numbers. A  $k$ -input Turing machine  $\mathcal{M}$  has  $k$  input tapes, one work tape, and one output tape. Each cell of a tape has symbols taken either from the set  $\{0, 1\}$  or the blank symbol  $\sqcup$ . At the start of the execution of a Turing machine, the work and output tapes are initialized to have all blank symbols. A Turing machine *halts* when it enters a designated halt state. The *output* of a Turing machine, when it halts, is the binary string that occupies the longest initial stretch of the output tape that does not have a blank symbol. If there are only blank symbols on the output tape, then by convention we say that the Turing machine's output is 0.

Every  $k$ -input Turing machine  $\mathcal{M}$  computes a (partial) function  $f : (\{0, 1\}^*)^k \rightarrow \{0, 1\}^*$  where the function is only defined on subset  $S \subseteq (\{0, 1\}^*)^k$  of inputs  $x$  on which  $\mathcal{M}$  halts. We use  $\mathcal{M}(x_1, x_2, \dots, x_k)$  to denote the output of a  $k$ -input Turing machine  $\mathcal{M}$  when  $x_i \in \{0, 1\}^*$  is written on the  $i$ -th input tape for  $i \in \{1, 2, \dots, k\}$ . If  $\mathcal{M}$  does not halt on an input  $x$ , then we define  $\mathcal{M}(x)$  to be  $\perp$ .

The *time complexity* of a Turing machine  $\mathcal{M}$  on input  $x = (x_1, x_2, \dots, x_k)$ , denoted by  $\text{TIME}_{\mathcal{M}, x}$ , is the number of time steps that  $\mathcal{M}$  takes on input  $x$  before it enters its designated halt state; if  $\mathcal{M}$  never halts on input  $x$ , then we define  $\text{TIME}_{\mathcal{M}, x} = \infty$ .

Every Turing machine  $\mathcal{M}$  has a canonical encoding as a bit string  $\overline{\mathcal{M}} \in \{0, 1\}^*$ , called the *canonical description* of  $\mathcal{M}$ . The canonical encoding describes the finite number of states and transition rules of  $\mathcal{M}$ . The details of this encoding are not important for this paper; we assume that some “reasonable” encoding scheme is used where the bit length of the encoding is at most some fixed polynomial in the number of states. The length of the description  $\overline{\mathcal{M}}$  is denoted by  $|\overline{\mathcal{M}}|$ . For every integer  $k \in \mathbb{N}$  and every string  $\alpha \in \{0, 1\}^*$ , the  $k$ -input Turing machine described by  $\alpha$  is denoted  $[\alpha]_k$ .

<sup>19</sup>The additional 1 in the argument of the  $\log(\cdot)$  is to ensure that this quantity is strictly positive.



Throughout this paper we frequently give high-level descriptions of Turing machines in English language, rather than explicitly describing them in terms of states and transition functions. In doing so we implicitly assume that all such descriptions can be formally converted into a description in terms of states and transition functions such that there is no hidden blow-up in the complexity of the representation.

We elaborate on several properties of Turing machines that are implicitly used throughout the paper.

**Turing machine simulation.** We frequently describe Turing machines as running or simulating other Turing machines as “subroutines”. We assume that such simulations can be performed with a polynomial overhead in terms of the time complexity. For example, suppose  $\mathcal{A}$  is a 1-input Turing machine that has the following high-level description: on input  $x$  (interpreted as an integer) it runs a 2-input Turing machine  $\mathcal{B}$  on the input  $(x, x^2)$  and obtains an output  $y$  (if  $\mathcal{B}$  halts), and then finally  $\mathcal{A}$  returns the output  $y$ . We assume that  $\mathcal{A}$  can efficiently simulate the Turing machine  $\mathcal{B}$ , even though  $\mathcal{B}$  has a different number of input tapes. This is because every  $k$ -input Turing machine  $\mathcal{M}$  can be efficiently simulated by a single *tape* Turing machine, as given by the following result [HS65]. We assume that there is an unambiguous, binary encoding  $\text{enc}_k(x)$  of  $k$ -tuples  $x \in (\{0, 1\}^*)^k$  that is computable in time  $O(k + |x_1| + \dots + |x_k|)$  where  $x_i$  is the  $i$ -th component of the tuple  $x$ .<sup>20</sup>

**Theorem 3.1** (Efficient universal Turing machine [HS65]). *For all  $k \in \mathbb{N}$  there exists a single tape Turing machine  $\mathcal{U}_k$  (called a universal Turing machine) with such that for every  $x \in (\{0, 1\}^*)^k$  and  $\alpha \in \{0, 1\}^*$ , if the tape of the Turing machine  $\mathcal{U}_k$  is initialized with  $\text{enc}_{k+1}(\alpha, x_1, \dots, x_k)$  and  $[\alpha]_k$  halts within  $T$  steps on input  $x$  then  $\mathcal{U}_k$  halts in at most  $C(k \cdot |\alpha| \cdot |x| \cdot T)^c$  steps and the contents of its tape is  $[\alpha]_k(x)$ . Here,  $C, c \geq 1$  are universal constants, and  $|x|$  denotes the sum of lengths  $|x_1| + \dots + |x_k|$ .*

Thus, the Turing machine  $\mathcal{A}$  can simulate  $\mathcal{B}$  on input  $(x, x^2)$  as follows: it first computes the encoding  $\text{enc}_3(\mathcal{B}, x, x^2)$  and writes this on a segment of the work tape. Then, it runs the Turing machine  $\mathcal{U}_k$  from Theorem 3.1 on this segment of the work tape, obtaining output  $y = \mathcal{B}(x, x^2)$  (if it eventually halts). Then, it writes  $y$  onto the output tape of  $\mathcal{A}$ . The total time complexity of  $\mathcal{A}$  on input  $x$  then includes the time complexity of computing the encoding, running Turing machine  $\mathcal{U}_k$ , and writing the final output.

We also assume that the length of the canonical description of  $\mathcal{A}$  depends only linearly on the canonical description of the Turing machine  $\mathcal{B}$ , so that  $|\overline{\mathcal{A}}| = O(|\overline{\mathcal{B}}|)$ ; the constant in the  $O(\cdot)$  notation hides the dependence on the description of the universal Turing machine  $\mathcal{U}_k$ , the computation of the encoding map  $\text{enc}_3(\cdot)$ , etc.

**Hardwiring constants.** Throughout this paper, we often define Turing machines  $\mathcal{M}$  with some number  $k$  of inputs, and then for some string  $a \in \{0, 1\}^*$  define a  $(k - 1)$ -input Turing machine  $\mathcal{M}_a$  whose behavior on input  $(y_1, \dots, y_{k-1})$  is to execute  $\mathcal{M}$  on input  $(a, y_1, \dots, y_{k-1})$ . Informally, the Turing machine  $\mathcal{M}_a$  “hardwires” the input  $a$  onto the first tape of  $\mathcal{M}$ . We informally write  $\mathcal{M}_a(y_1, \dots, y_{k-1}) = \mathcal{M}(a, y_1, \dots, y_{k-1})$ .

More precisely, the Turing machine  $\mathcal{M}_a$  on input  $(y_1, \dots, y_{k-1})$  first computes the encoding  $e = \text{enc}_{k+1}(\overline{\mathcal{M}}, a, y_1, \dots, y_{k-1})$ , and then runs the universal Turing machine  $\mathcal{U}_k$  on input  $e$ . The description of  $\mathcal{M}_a$  can be computed from  $\overline{\mathcal{M}}$  and  $a$  in time  $O(|a| + |\overline{\mathcal{M}}| + O(1))$ , where the  $O(1)$  comes from the description length of the universal Turing machine  $\mathcal{U}_k$ .

Furthermore, the time complexity of the Turing machine  $\mathcal{M}_a$  on input  $(y_1, \dots, y_{k-1})$  is at most  $\text{poly}(|a|, |y_1|, \dots, |y_{k-1}|, T)$  where  $T$  is the time complexity of  $\mathcal{M}$  on input  $(a, y_1, \dots, y_{k-1})$ . This bound

<sup>20</sup>A canonical choice of such an encoding is the following: a tuple  $(x_1, \dots, x_k)$  is encoded into a concatenation of a “dual rail” encoding of each  $x_i$ : every bit of  $x_i$  is expanded via the map  $0 \mapsto 01, 1 \mapsto 10$ , and the end of the string  $x_i$  is indicated by 00.

comes from the complexity of encoding the inputs from the  $k$  tapes into the  $\text{enc}_{k+1}(\cdot)$  format, and then simulating  $\mathcal{M}$  on input  $(\overline{\mathcal{M}}, a, y_1, \dots, y_{k-1})$ .

**Description complexity bounds.** When we bound the description lengths of the Turing machines presented in this paper, we do not worry about the exact details of how Turing machines are represented as binary strings – as mentioned we assume that *some* reasonable encoding is used – but instead we distinguish between whether the Turing machine description depends on any parameters used elsewhere in the paper. Note that whether the *description* depends on any parameters is different from whether the Turing machine takes parameters as *input*. For example, consider the following (high-level) descriptions of Turing machines  $\mathcal{A}$  and  $\mathcal{B}$ :

1. Turing machine  $\mathcal{A}$  takes two inputs  $(\alpha, x)$  and outputs the integer  $\alpha \times x$  where  $\alpha, x$  are interpreted as positive integers written in binary.
2. Turing machine  $\mathcal{B}$  takes one input  $x$  and outputs the integer  $\beta \times x$  where  $\beta$  is a fixed positive integer.

The description of  $\mathcal{A}$  does not depend on any “external” parameters, so its description length  $|\overline{\mathcal{A}}|$  is some universal constant, which we express as  $O(1)$ . On the other hand, the description of  $\mathcal{B}$  depends on some fixed integer  $\beta$ ; in other words, the parameter  $\beta$  is “hard-wired” into  $\mathcal{B}$ ’s description. The description of  $\mathcal{B}$  can be taken to be the following:  $\mathcal{B}$  (which is a 1-input Turing machine) simulates the execution of the 2-input Turing machine  $\mathcal{A}$  where the first input tape of  $\mathcal{A}$  has the binary representation of  $\beta$  written onto it and the second input tape has  $x$  (which is provided as input to  $\mathcal{B}$ ) written into it. Thus the description of  $\mathcal{B}$  includes the binary representation of  $\beta$ , the description of  $\mathcal{A}$  (whose length is a universal constant), and the description of the universal Turing machine (whose length is a universal constant). Thus the description length of  $\mathcal{B}$  is  $O(\log \beta) + O(1) = O(\log \beta)$ .

**Time complexity bounds.** When we bound the time complexity of the Turing machines presented in this paper, we similarly do not worry about the exact implementation details, but instead we assume that basic computations such as integer arithmetic, string comparisons, etc. are all performed using reasonably efficient algorithms that run in polynomial time in the length of the input. When running other Turing machines as subroutines, we assume that there is some polynomial overhead due to Theorem 3.1.

**Timeout counters.** We sometimes define Turing machines that are required to halt if either the Turing machine or some subroutine takes a number of time steps that exceeds some specified threshold. For example, we may write “Let  $\mathcal{M}$  be the Turing machine that on input the description  $\overline{\mathcal{R}}$  of a Turing machine  $\mathcal{R}$  and an integer  $T$ , simulates  $\mathcal{R}$  on the empty tape and halts if more than  $T$  steps are performed.”

This is useful for establishing an *a priori* bound on the time complexity of the Turing machine. In particular we will be able to claim that  $\mathcal{M}$ , as described above, always runs in time at most  $\text{poly}(T, |\overline{\mathcal{R}}|)$ , irrespective of the running time of  $\mathcal{R}$  itself.

We explain how such a Turing machine can be realized formally. We use a variant of the universal Turing machine of Theorem 3.1. Using the theorem, it is straightforward to show that given a (single tape) Turing machine  $\mathcal{R}$  there exists another two-tape Turing machine  $\mathcal{M}'$  such that, whenever the second tape is initialized to the binary representation of  $T$ ,  $\mathcal{M}'$  simulates  $\mathcal{R}$  and in-between every simulated step of  $\mathcal{R}$ , decrements the second tape and checks if it reaches zero. If it does, then  $\mathcal{M}'$  halts and rejects. Otherwise, it continues.

The time complexity of  $\mathcal{M}'$  is, by definition, at most  $O(T \log T)$  for all inputs. The description length of  $\mathcal{M}'$  is at most  $O(|\overline{\mathcal{R}}|)$ , because this is the description size of the universal Turing machine, whose initial tape has been hardwired with  $\overline{\mathcal{R}}$ . To convert back to a single tape Turing machine and obtain  $\mathcal{M}$  from  $\mathcal{M}'$  we can use Theorem 3.1 again, at the cost of a polynomial blow-up in the time complexity.

**Input representations.** Although the inputs and outputs of a Turing machine are strictly speaking binary strings, we oftentimes slightly abuse notation and specify Turing machines that treat their inputs and outputs as objects with more structure, such as finite field elements, integers, symbols from a larger alphabet, and so on. In this case we implicitly assume that the Turing machine specification uses a consistent convention to represent these structured objects as binary strings. Conventions for objects such as integers are straightforward. For representations of finite field elements, we refer the reader to Section 3.3.2.

### 3.2 Linear spaces

Linear spaces considered in the paper generally take the form  $V = \mathbb{F}^n$  for a finite field  $\mathbb{F}$  and integer  $n \geq 1$ . In particular, when we write “let  $V$  be a linear space”, unless explicitly stated otherwise we always mean a space of the form  $\mathbb{F}^n$ . Let  $\hat{E} = \{\hat{e}_1, \hat{e}_2, \dots, \hat{e}_n\}$  denote the standard basis of  $V$ , where for  $i \in \{1, 2, \dots, n\}$ ,

$$\hat{e}_i = (0, \dots, 0, 1, 0, \dots, 0)$$

has a 1 in the  $i$ -th coordinate and 0's elsewhere. We write  $\text{End}(V)$  to denote the set of linear transformations from  $V$  to itself.

**Definition 3.2** (Register subspace). A *register subspace*  $S$  of  $V$  is a subspace that is the span of a subset of the standard basis of  $V$ .<sup>21</sup> We often represent such a subspace as an indicator vector  $u \in \{0, 1\}^s$ , where  $s = \dim(V)$ , such that if  $\{\hat{e}_1, \dots, \hat{e}_s\}$  is the standard basis of  $V$  then  $S = \text{span}\{\hat{e}_i \mid u_i = 1\}$ .

**Definition 3.3.** Let  $\hat{E} = \{\hat{e}_i\}$  be the standard basis of  $V = \mathbb{F}^n$ . For two vectors  $u = \sum_{i=1}^n u_i \hat{e}_i$ ,  $v = \sum_{i=1}^n v_i \hat{e}_i$  in  $V$ , define the *dot product*

$$u \cdot v = \sum_{i=1}^n u_i v_i \in \mathbb{F}.$$

Let  $S$  be a subspace of  $V$ . The *subspace orthogonal to  $S$  in  $V$*  is

$$S^\perp = \{u \in V : u \cdot v = 0 \text{ for all } v \in S\}.$$

Although the notation  $S^\perp$  does not explicitly refer to  $V$ , the ambient space will always be clear from context.

We note that over finite fields, the notion of orthogonality does not possess all of the same intuitive properties of orthogonality over fields such as  $\mathbb{R}$  or  $\mathbb{C}$ ; for example, a non-zero subspace  $S$  may be orthogonal to itself (e.g.  $\text{span}\{(1, 1)\}$  over  $\mathbb{F}_2$ ). However, the following remains true over all fields.

**Lemma 3.4.** *Suppose  $S$  is a subspace of  $V$ . Then  $\dim(S) + \dim(S^\perp) = \dim(V)$ , and*

$$(S^\perp)^\perp = S.$$

---

<sup>21</sup>The use of the term “register” is meant to create an analogy for how the space of multiple qubits is often partitioned into “registers” containing a few qubits each.

*Proof.* The statement about the dimensions follows from the fact that vectors in  $S^\perp$  are the solution to a feasible linear system of equations with  $\dim(S)$  linearly independent rows; this implies that the solution space has dimension exactly  $\dim(V) - \dim(S)$ . Next, we argue that  $S \subseteq (S^\perp)^\perp$ . Let  $u \in S$ . Since all vectors  $v \in S^\perp$  are orthogonal to every vector in  $S$ , in particular  $u$ , this implies that  $u \in (S^\perp)^\perp$ . By dimension counting, it follows that  $(S^\perp)^\perp = S$ .  $\square$

**Definition 3.5.** Given a linear space  $V$ , two subspaces  $S$  and  $T$  of  $V$  are said to form a pair of *complementary subspaces* of  $V$  if

$$S \cap T = \{0\}, \quad S + T = V.$$

In this case, we write  $V = S \oplus T$ . Any  $x \in V$  can be written as  $x = x^S + x^T$  for  $x^S \in S$  and  $x^T \in T$  in a unique way. We refer to  $x^S$  (resp.  $x^T$ ) as the *projection of  $x$  onto  $S$  parallel to  $T$*  (resp. *onto  $T$  parallel to  $S$* ). We call the unique linear map  $L : V \rightarrow V$  that maps  $x \mapsto x^S$  the *projector onto  $S$  parallel to  $T$* .

A given subspace may have many different complementary subspaces: consider the example of  $S = \text{span}\{(1, 1)\}$  in  $\mathbb{F}_2^2$ . Different complementary subspaces include  $T = \text{span}\{(1, 0)\}$  and  $T' = \text{span}\{(0, 1)\}$ . It is convenient to define the notion of a *canonical complement* of a subspace  $S$ , given a basis for  $S$ .

**Definition 3.6.** Let  $E$  be the standard basis of linear space  $V = \mathbb{F}^n$ . Let  $F = \{v_1, v_2, \dots, v_m\} \subset V$  be a set of  $m$  linearly independent vectors in  $V$ . The *canonical complement*  $F^\perp$  of  $F$  is the set of  $n - m$  independent vectors defined as follows. Write  $v_i = \sum_{j=1}^n a_{ij} \hat{e}_j$ . Using a canonical algorithm for Gaussian elimination that works over arbitrary fields, transform the  $m \times n$  matrix  $(a_{ij})$  to reduced row echelon form  $(b_{ij})$ . Let  $J$  be the set of  $m$  column indices of the leading 1 entry in each row of  $(b_{ij})$ . The canonical complement is defined as  $F^\perp = \{\hat{e}_j : j \notin J\}$ .

**Remark 3.7.** We emphasize that the canonical complement  $F^\perp$  is a set (rather than a subspace), and it is defined with respect to a set of linearly independent vectors  $F$ . While there are equivalent ways of defining a canonical complement of a subspace in a basis-independent manner, we use this particular definition because it makes it clear that the canonical complement of a set  $F$  is efficiently computable.

**Remark 3.8.** Let  $\hat{E}$  be the standard basis of  $V$ . Suppose subspace  $S$  is a register subspace of  $V$  spanned by  $\hat{E}_0 \subseteq \hat{E}$ . Then it is not hard to verify that the canonical complement of  $\hat{E}_0$  is  $\hat{E} \setminus \hat{E}_0$  and the span of the canonical complement coincides with  $S^\perp$ .

**Lemma 3.9.** Let  $S$  be the span of linearly independent vectors  $F = \{v_1, \dots, v_m\} \subseteq V$  and let  $F^\perp$  be the canonical complement of  $F$  as defined in Definition 3.6. Let  $T = \text{span}(F^\perp)$ . Then

$$S + T = V, \quad S \cap T = \{0\}.$$

*Proof.* To show the first item, we must show that every vector in  $V$  can be written as a sum of an element of  $S$  and an element of  $T$ . To do this, let  $A = (a_{i,j})$  be the  $m \times n$  matrix over  $\mathbb{F}$  whose rows are the vectors  $v_i$ , as in Definition 3.6. Write  $A = UB$  where  $U$  is invertible and  $B$  is in reduced row echelon form. Let  $J$  be the set of column indices of the leading 1 entries in  $B$ , as in Definition 3.6. Now, the rows of  $B$  are linearly independent and span the subspace  $S$ , and the restriction of these rows to the columns in  $J$  are still linearly independent and span  $\mathbb{F}^m$ . Thus, for any vector  $u \in V$ , there is some linear combination  $v$  of rows of  $B$  such that  $u$  and  $v$  agree restricted to the columns in  $J$ . In other words, there exists  $v \in S$  such that  $u_j = v_j$  for all  $j \in J$ . Hence,  $u = v + w$  for  $w \in T$ , where  $T$  is the canonical complement. This shows  $S + T = V$ . Counting dimensions shows that necessarily  $S \cap T = \{0\}$ .  $\square$

**Definition 3.10.** Let  $F \subseteq V$  be a set of linearly independent vectors. Let  $F^\perp$  be the canonical complement of  $F$ . Define the *canonical linear map*  $L \in \text{End}(V)$  with kernel basis  $F$  as the projector onto  $T$  parallel to  $S$ , where  $S = \text{span}(F)$  and  $T = \text{span}(F^\perp)$ . When the basis  $F$  for  $S$  is clear from context, we refer to this map as the *canonical linear map with kernel*  $S$ .

**Definition 3.11.** Let  $L \in \text{End}(V)$  be a linear map, and let  $F$  be a basis for  $\ker(L)^\perp$ . Define  $L^\perp : V \rightarrow V$  as the canonical linear map with kernel basis  $F$ .

**Lemma 3.12.** Let  $L \in \text{End}(V)$  be a linear map and  $F$  a basis for  $\ker(L)^\perp$ . Let  $L^\perp \in \text{End}(V)$  be the linear map defined in Definition 3.11. Then  $\ker(L^\perp) = \ker(L)^\perp$ .

*Proof.* First, to set notation, let  $S = \text{span}(F) = \ker(L)^\perp$  and  $T = \text{span}(F^\perp)$ . By Lemma 3.9, any vector  $v \in V$  can be uniquely decomposed as  $v = v^S + v^T$ .

Now we will show the lemma. First we show that  $\ker(L)^\perp \subseteq \ker(L^\perp)$ . Let  $v \in \ker(L)^\perp$ , and write  $v = v^S + v^T$ . By the definition of  $L^\perp$ , it follows that  $v^T = 0$ , and hence  $v \in S = \ker(L)^\perp$ .

It remains to show the other direction, that is  $\ker(L^\perp) \subseteq \ker(L)^\perp$ . Let  $v \in \ker(L^\perp) = S$ . Then  $v = v^S + v^T$  where  $v^T = 0$ , and hence by the definition of  $L^\perp$ ,  $L^\perp v = 0$ . Hence  $v \in \ker(L)^\perp$ . □

### 3.3 Finite fields

Let  $p$  be a prime and  $q = p^k$  be a prime power. We denote the finite fields of  $p$  and  $q$  elements by  $\mathbb{F}_p$  and  $\mathbb{F}_q$  respectively. The prime  $p$  is the characteristic of field  $\mathbb{F}_q$ , and field  $\mathbb{F}_p$  is the *prime subfield* of  $\mathbb{F}_q$ . We sometimes omit the subscript and simply use  $\mathbb{F}$  to denote the finite field when the size of the field is implicit from context. For general background on finite fields, and explicit algorithms for elementary arithmetic operations, we refer to [MP13].

#### 3.3.1 Subfields and bases

Let  $q$  be a prime power and  $k$  an integer. The field  $\mathbb{F}_q$  is a subfield of  $\mathbb{F}_{q^k}$  and  $\mathbb{F}_{q^k}$  is a linear space of dimension  $k$  over  $\mathbb{F}_q$ . Let  $\{e_i\}_{i=1}^k$  be a basis of  $\mathbb{F}_{q^k}$  as a linear space over  $\mathbb{F}_q$ . Introduce a bijection  $\kappa_q : \mathbb{F}_{q^k} \rightarrow \mathbb{F}_q^k$  between  $\mathbb{F}_{q^k}$  and  $\mathbb{F}_q^k$  defined with respect to the basis  $\{e_i\}_{i=1}^k$  by

$$\kappa_q : a \mapsto (a_i)_{i=1}^k$$

where  $a = \sum_{i=1}^k a_i e_i$ . This map satisfies several nice properties. First, the map is an isomorphism of  $\mathbb{F}_q$ -vector spaces: it is  $\mathbb{F}_q$ -linear and addition in  $\mathbb{F}_{q^k}$  naturally corresponds to vector addition in  $\mathbb{F}_q^k$ . Namely, for all  $a, b \in \mathbb{F}_{q^k}$ ,

$$\kappa_q(a + b) = \kappa_q(a) + \kappa_q(b) .$$

Second, multiplication by a field element in  $\mathbb{F}_{q^k}$  corresponds to a linear map on  $\mathbb{F}_q^k$ . For all  $a \in \mathbb{F}_{q^k}$ , there exists a matrix  $K_a \in \text{M}_k(\mathbb{F}_q)$  such that for all  $b \in \mathbb{F}_{q^k}$ ,

$$\kappa_q(ab) = K_a \kappa_q(b) .$$

The matrix  $K_a$  is called the *multiplication table of  $a$  with respect to basis*  $\{e_i\}_{i=1}^k$ .

We extend the map  $\kappa_q$  to vectors, matrices and sets over  $\mathbb{F}_{q^k}$ . For  $v = (v_1, v_2, \dots, v_n) \in \mathbb{F}_{q^k}^n$ , define

$$\kappa_q(v) = (\kappa_q(v_i))_{i=1}^n \in \mathbb{F}_q^{kn} .$$

Similarly, for matrix  $M = (M_{i,j}) \in \mathbf{M}_{m,n}(\mathbb{F}_{q^k})$ , define

$$\chi_q(M) = (K_{M_{i,j}}) \in \mathbf{M}_{mk,nk}(\mathbb{F}_q),$$

the block matrix whose  $(i, j)$ -th block is the multiplication table  $K_{M_{i,j}}$  of  $M_{i,j}$  with respect to basis  $\{e_i\}_{i=1}^k$ . For a set  $S$  of vectors in  $\mathbb{F}_{q^k}^n$ , define

$$\kappa_q(S) = \{\kappa_q(v) : v \in S\}.$$

We omit the subscript and write  $\kappa$  and  $\chi$  for  $\kappa_q$  and  $\chi_q$  respectively when  $q$  equals to  $p$ , the characteristic of the field.

The *trace of  $\mathbb{F}_{q^k}$  over  $\mathbb{F}_q$*  is defined as

$$\mathrm{tr}_{q^k \rightarrow q} : a \mapsto \mathrm{Tr}(K_a) \tag{10}$$

for  $a \in \mathbb{F}_{q^k}$ , where  $\mathrm{Tr}(K_a)$  is the trace of the multiplication table of  $a$  with respect to the basis  $\{e_i\}$ . By definition, the trace is an  $\mathbb{F}_q$ -linear map from  $\mathbb{F}_{q^k}$  to  $\mathbb{F}_q$ . The trace is in fact independent of the choice of basis, which can be seen from the following equivalent definition [MP13, Definition 2.1.80]:

$$\mathrm{tr}_{q^k \rightarrow q}(a) = \sum_{j=0}^{k-1} a^{q^j}.$$

A *dual basis*  $\{e'_1, e'_2, \dots, e'_k\}$  of  $\{e_1, e_2, \dots, e_k\}$  is a basis such that  $\mathrm{tr}_{q^k \rightarrow q}(e_i e'_j) = \delta_{i,j}$  for all  $i, j \in \{1, 2, \dots, k\}$ . A *self-dual basis* is one that is equal to its dual. If for some  $\alpha \in \mathbb{F}_{q^k}$  the set  $\{\alpha^{q^j}\}_{j=0}^{k-1}$  forms a basis of  $\mathbb{F}_{q^k}$  over  $\mathbb{F}_q$ , the basis is called a *normal basis*. Self-dual and normal bases do not exist for all fields but are guaranteed to exist under certain conditions; in particular, if  $q = 2$  and  $k$  is odd, as will be shown in Lemma 3.16.

We record some convenient facts about the maps  $\kappa(\cdot)$  and  $\chi(\cdot)$  for self-dual bases.

**Lemma 3.13.** *Let  $q$  be a prime power,  $k$  an integer and  $\{e_i\}$  a self-dual basis for  $\mathbb{F}_{q^k}$  over  $\mathbb{F}_q$ . The map  $\kappa_q(\cdot)$  corresponding to  $\{e_i\}$  satisfies the following properties:*

1. For all  $x \in \mathbb{F}_{q^k}$ ,  $\kappa_q(x) = (\mathrm{tr}_{q^k \rightarrow q}(xe_1), \dots, \mathrm{tr}_{q^k \rightarrow q}(xe_k))$ .
2. For all  $x, y \in \mathbb{F}_{q^k}$ ,  $\mathrm{tr}_{q^k \rightarrow q}(xy) = \kappa_q(x) \cdot \kappa_q(y)$ .
3. For all  $M \in \mathbf{M}_{m,n}(\mathbb{F}_{q^k})$  and  $v \in \mathbb{F}_{q^k}^n$ , we have  $\chi_q(M)\kappa_q(v) = \kappa_q(Mv)$ .

*Proof.* We show each property by direct calculation.

1. Let  $\kappa_q(x) = (x_1, \dots, x_k)$ . Then by the definition of  $\kappa_q$ ,  $x = \sum_i x_i e_i$ . Multiplying both sides by  $e_j$  and taking the trace yields  $\mathrm{tr}_{q^k \rightarrow q}(xe_j) = \sum_i x_i \mathrm{tr}_{q^k \rightarrow q}(e_i e_j) = x_j$ , where the last equality is by self-duality of the basis.
2. Write  $\kappa_q(x) = (x_1, \dots, x_k)$  and  $\kappa_q(y) = (y_1, \dots, y_k)$ . Then

$$\begin{aligned} \mathrm{tr}_{q^k \rightarrow q}(xy) &= \mathrm{tr}_{q^k \rightarrow q}\left(\sum_{ij} x_i y_j e_i e_j\right) \\ &= \sum_{ij} x_i y_j \mathrm{tr}_{q^k \rightarrow q}(e_i e_j) \\ &= \kappa_q(x) \cdot \kappa_q(y), \end{aligned}$$

where the last equality uses the self-duality of the basis.

3. We compute the  $i$ th  $\mathbb{F}_q^k$ -block of  $\kappa_q(Mv)$ :

$$\begin{aligned}\kappa_q(Mv)_i &= \sum_{ij} \kappa_q(M_{ij}v_j) \\ &= \sum_{ij} K_{M_{ij}} \kappa_q(v_j) \\ &= \chi_q(M)_{ij} \kappa_q(v_j),\end{aligned}$$

where we have applied the definition of the multiplication table  $K_{M_{ij}}$  and the map  $\chi_q(\cdot)$ . □

For  $z \in \mathbb{F}^n$  and  $V, W$  a pair of complementary subspaces, recall from Definition 3.5 the notation  $z^V$  for the projection of  $z$  onto  $V$  and parallel to  $W$ .

**Lemma 3.14.** *Let  $\kappa_q(\cdot)$  denote the map corresponding to a basis  $\{e_i\}$  for  $\mathbb{F}_{q^k}$  over  $\mathbb{F}_q$ . Let  $V$  be a subspace of  $\mathbb{F}_{q^k}^n$  with linearly independent basis  $\{b_1, \dots, b_t\} \subseteq \mathbb{F}_{q^k}^n$ . Then the following hold:*

1.  $\kappa_q(V)$  is a subspace of  $\mathbb{F}_q^{nk}$ .
2.  $\{\kappa_q(e_i b_j)\}_{i,j}$  is a linearly independent basis of  $\kappa_q(V)$  over  $\mathbb{F}_q$ .
3. Let  $V, W$  be complementary subspaces of  $\mathbb{F}_{q^k}^n$ . Then  $V' = \kappa_q(V)$  and  $W' = \kappa_q(W)$  are complementary subspaces of  $\mathbb{F}_q^{kn}$ , and furthermore for all vectors  $z \in \mathbb{F}_{q^k}^n$ , we have  $\kappa_q(z^V) = \kappa_q(z)^{V'}$  and  $\kappa_q(z^W) = \kappa_q(z)^{W'}$ .

*Proof.* For the first item, we first verify that  $\kappa_q(V)$  is a subspace. Since  $V$  is a subspace, it contains  $0 \in \mathbb{F}_{q^k}^n$ , and therefore  $\kappa_q(0) = 0$  is also in  $\kappa_q(V)$ . Let  $u', v' \in \kappa_q(V)$ . Using that  $\kappa_q$  is a bijection there exist  $u, v \in \mathbb{F}_{q^k}^n$  such that  $u' = \kappa_q(u)$  and  $v' = \kappa_q(v)$ . Therefore

$$u' + v' = \kappa_q(u) + \kappa_q(v) = \kappa_q(u + v) \in \kappa_q(V),$$

where the inclusion follows because  $V$  is a subspace and thus contains  $u + v$ . Finally, for all  $x' \in \mathbb{F}_q$ , for all  $v \in V$ , we have that  $x' \kappa_q(v) = \kappa_q(x'v) \in \kappa_q(V)$  where we used that  $V$  is closed under scalar multiplication by  $\mathbb{F}_{q^k}$  and thus by  $\mathbb{F}_q$  (since  $\mathbb{F}_q$  is a subfield of  $\mathbb{F}_{q^k}$ ). Thus  $\kappa_q(V)$  is closed under scalar multiplication by  $\mathbb{F}_q$ .

For the second item, note that an element  $v \in V$  can be expressed uniquely as  $v = \sum_{i=1}^t v_i b_i$  for  $v_i \in \mathbb{F}_{q^k}$ . The element  $v_i$  can further be written as  $\sum_j v_{i,j} e_j$  where  $v_{i,j} \in \mathbb{F}_q$ . Thus  $v$  is a linear combination of the vectors  $\{e_j b_i\}$ , and therefore  $\kappa_q(v)$  is a linear combination of the vectors  $\{\kappa_q(e_j b_i)\}$ . To establish that the vectors  $\{\kappa_q(e_j b_i)\}$  are linearly independent, suppose towards contradiction that they are not. Then there would exist  $\alpha_{i,j} \in \mathbb{F}_q$  such that at least one  $\alpha_{i,j}$  is nonzero and

$$\begin{aligned}0 &= \sum_{i,j} \alpha_{i,j} \kappa_q(e_j b_i) \\ &= \kappa_q\left(\sum_i \left(\sum_j \alpha_{i,j} e_j\right) b_i\right) \\ &= \kappa_q\left(\sum_i \beta_i b_i\right),\end{aligned}$$

where we define  $\beta_i = \sum_j \alpha_{i,j} e_j$ . Since at least one  $\alpha_{i,j} \neq 0$  and the  $\{e_j\}$  are linearly independent over  $\mathbb{F}_q$ , there exists  $i$  such that  $\beta_i \neq 0$ , which means that there is a non-trivial linear combination of the basis elements  $b_i$  that equals 0 under  $\kappa_q(\cdot)$ . Since  $\kappa_q(\cdot)$  is injective, we get a contradiction with linear independence of the  $\{b_i\}$ .

For the third item, we observe that  $\kappa_q(V)$  and  $\kappa_q(W)$  must be complementary because  $\kappa_q(\cdot)$  is a linear map as well as a bijection. Let  $\{v_1, \dots, v_m\}$  and  $\{v_{m+1}, \dots, v_n\}$  denote bases for  $V$  and  $W$ , respectively. Thus the set  $\{v_1, \dots, v_n\}$  forms a basis for  $\mathbb{F}_q^n$ , and from the previous item, the set  $\{\kappa_q(e_j v_i)\}_{i,j}$  is a basis for  $\mathbb{F}_q^{kn}$ . Furthermore, the sets  $\{\kappa_q(e_j v_i)\}_{j,i=1,\dots,m}$  and  $\{\kappa_q(e_j v_i)\}_{j,i=m+1,\dots,n}$  are bases for  $\kappa_q(V)$  and  $\kappa_q(W)$ , respectively.

There is a unique choice of coefficients  $\alpha_{i,j} \in \mathbb{F}_q$  such that  $\kappa_q(z) = \sum_{i,j} \alpha_{i,j} \kappa_q(e_j v_i)$ . But then

$$\begin{aligned} \kappa_q(z) &= \kappa_q\left(\sum_i \left(\sum_j \alpha_{i,j} e_j\right) v_i\right) \\ &= \kappa_q\left(\sum_i \alpha_i v_i\right), \end{aligned}$$

where we define  $\alpha_i = \sum_j \alpha_{i,j} e_j$ . Since  $\kappa_q(\cdot)$  is a bijection, this implies that  $z = \sum_i \alpha_i v_i$ , and therefore  $z^V = \sum_{i=1}^m \alpha_i v_i$  (and similarly  $z^W = \sum_{i=m+1}^n \alpha_i v_i$ ). This implies that

$$\kappa_q(z)^{V'} = \sum_{i=1}^m \sum_j \alpha_{i,j} \kappa_q(e_j v_i) = \kappa_q(z^V),$$

and similarly  $\kappa_q(z)^{W'} = \kappa_q(z^W)$ . This completes the proof of the lemma.  $\square$

### 3.3.2 Bit string representations

As mentioned at the end of Section 3.1, we sometimes treat the inputs and outputs of Turing machines as representing elements of a finite field, or a vector space over a finite field. We discuss some important details about bit representations of finite field elements and arithmetic over finite fields.

In the paper we only consider fields  $\mathbb{F}_{2^k}$  where  $k$  is odd.

**Definition 3.15.** A field size  $q$  is called an *admissible* field size if  $q = 2^k$  for odd  $k$ .

Elements of  $\mathbb{F}_2$  are naturally represented using bits. To represent elements of  $\mathbb{F}_{2^k}$  as binary strings we require the specification of a basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$ . Given a basis  $\{e_i\}_{i=1}^k$  of  $\mathbb{F}_{2^k}$ , every element  $a \in \mathbb{F}_{2^k}$  has a unique expansion  $a = \sum_{i=1}^k a_i e_i$  and can be represented as the  $k$ -bit string corresponding to  $\kappa(a) \in \mathbb{F}_2^k$ . Note that we omitted the subscript 2 of  $\kappa$  as it maps to the linear space over the prime subfield  $\mathbb{F}_2$ . Thus the *binary representation* of  $a \in \mathbb{F}_{2^k}$  is defined as the natural binary representation of  $\kappa(a) \in \mathbb{F}_2^k$  (which in turn is the  $\mathbb{F}_2$ -representation of  $a$ ). Throughout the paper we freely associate between the binary representation of a field element  $a \in \mathbb{F}_{2^k}$  and its  $\mathbb{F}_2$ -representation, although—technically speaking—these are distinct objects.

Given the representations  $\kappa(a), \kappa(b)$  of  $a, b \in \mathbb{F}_{2^k}$ , to compute the binary representation of  $a + b$  it suffices to compute the addition bit-wise, modulo 2. Computing the multiplication of elements  $a, b$  requires the specification of the multiplication tables  $\{K_{e_i} \in \mathbf{M}_k(\mathbb{F}_2)\}_{i=1}^k$  for the basis  $\{e_i\}$ . Given representations  $\kappa(a) = (a_i)_{i=1}^k$ ,  $\kappa(b) = (b_i)_{i=1}^k$  for  $a, b \in \mathbb{F}_{2^k}$  respectively, the representation  $\kappa(ab)$  of the product  $ab$  is computed as

$$\kappa(ab) = \sum_{i=1}^k a_i \kappa(e_i b) = \sum_{i=1}^k a_i (K_{e_i} \kappa(b)). \quad (11)$$



Thus, using our representation for field elements, efficiently performing finite field arithmetic in  $\mathbb{F}_{2^k}$  reduces to having access to the multiplication table of some basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$ .

The following fact provides an efficient deterministic algorithm for computing a self-dual normal basis for  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$  and the corresponding multiplication tables for any odd  $k$ .

**Lemma 3.16.** *There exists a deterministic algorithm that given an odd integer  $k > 0$ , outputs a self-dual normal basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$  and the multiplication tables of the basis in  $\text{poly}(k)$  time.*

*Proof.* The algorithm of Shoup [Sho90, Theorem 3.2] shows that for prime  $p$ , an irreducible polynomial in  $\mathbb{F}_p[X]$  of degree  $k$  can be computed in time  $\text{poly}(p, k)$ . Then, the algorithm of Lenstra [LJ91, Theorem 1.1] shows that given such an irreducible polynomial, the multiplication table of a normal basis of  $\mathbb{F}_{p^k}$  over  $\mathbb{F}_p$  can be computed in  $\text{poly}(k, \log p)$  time. Finally, the algorithm of Wang [Wan89] shows that for odd  $k$  and a multiplication table  $K$  of a normal basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$ , a multiplication table  $K'$  for a self-dual normal basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$  can be computed in  $\text{poly}(k)$  time. Putting these three algorithms together yields the claimed statement.  $\square$

For  $q = 2^k$  for  $k$  odd (i.e.  $q$  is an admissible field size) we use the shorthand  $\text{tr}$  for  $\text{tr}_{q \rightarrow 2}$ .

**Lemma 3.17.** *Let  $k$  be an odd integer and  $\{e_i\}_{i=1}^k$  be a self-dual normal basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$ . Then  $\text{tr}(e_i) = 1$  for all  $i$ , and furthermore the representation  $\kappa(1)$  of the unit  $1 \in \mathbb{F}_{2^k}$  is the all ones vector in  $\mathbb{F}_2^k$ .*

*Proof.* Since  $\{e_i\}$  is a normal basis,  $e_i = \alpha^{2^i}$  for some element  $\alpha \in \mathbb{F}_{2^k}$ . Furthermore, for every element  $b \in \mathbb{F}_{2^k}$ , we have that  $\text{tr}(b^2) = \text{tr}(b)$ . This is because

$$\text{tr}(b^2) = \sum_{i=0}^{k-1} b^{2^{i+1}} = \sum_{i=0}^{k-1} b^{2^i} = \text{tr}(b),$$

where we use that  $b^{2^k} = b$  for all  $b \in \mathbb{F}_{2^k}$ . Since  $e_{i+1} = e_i^2$ , we get that  $\text{tr}(e_i) = \text{tr}(e_j)$  for all  $i, j$ . It cannot be the case that  $\text{tr}(e_i) = 0$  for all  $i$ . Suppose that this were the case. This would imply that  $\text{tr}(b) = 0$  for all  $b \in \mathbb{F}_{2^k}$ . But then for all  $j \in \{1, \dots, k\}$  and for some  $b \neq 0$ , we would also have by item 1 of Lemma 3.13 that  $b_j = \text{tr}(be_j) = 0$  where  $b = \sum_j b_j e_j$  with  $b_j \in \mathbb{F}_2$ . This implies that  $b$  is the all zero element of  $\mathbb{F}_{2^k}$ , which is a contradiction. Thus  $\text{tr}(e_i) = 1$  for all  $i = 1, 2, \dots, k$ .

The ‘‘furthermore’’ part follows item 1 of Lemma 3.13:

$$\kappa(1) = (\text{tr}(1 \cdot e_1), \dots, \text{tr}(1 \cdot e_k)) = (1, \dots, 1).$$

$\square$

**Lemma 3.18.** *For any odd integer  $k$ , let  $\{e_i\}_{i=1}^k$  denote the self-dual normal basis of  $\mathbb{F}_{2^k}$  over  $\mathbb{F}_2$  that is returned by the algorithm specified in Lemma 3.16 on input  $k$ . Then the following can be computed in time  $\text{poly}(k)$  on input  $k$ :*

1. The representation  $\kappa(a + b)$  of the sum  $a + b$  given the representations  $\kappa(a)$  and  $\kappa(b)$  of  $a, b \in \mathbb{F}_{2^k}$ .
2. The representation  $\kappa(ab)$  of the product  $ab$  given the representations  $\kappa(a)$  and  $\kappa(b)$  of  $a, b \in \mathbb{F}_{2^k}$ .
3. The multiplication table  $K_a \in M_k(\mathbb{F}_2)$  given the representation  $\kappa(a)$  of  $a \in \mathbb{F}_{2^k}$ .
4. The representation  $\kappa(a^{-1})$  of the multiplicative inverse of  $a \in \mathbb{F}_{2^k}$ , given the representation  $\kappa(a)$ .

5. The trace  $\text{tr}(a)$  given the multiplication table  $K_a$  of  $a \in \mathbb{F}_{2^k}$ .

Furthermore, for all integers  $n$ , the representations of projections  $\kappa(x^S)$  and  $\kappa(x^T)$  of  $x \in \mathbb{F}_{2^k}^n$  for complementary subspaces  $S, T$  of  $\mathbb{F}_{2^k}^n$  can be computed in  $\text{poly}(k, n)$  time, given the representations  $\kappa(x)$ ,  $\{\kappa(v_1), \kappa(v_2), \dots, \kappa(v_m)\}$  and  $\{\kappa(w_1), \kappa(w_2), \dots, \kappa(w_{n-m})\}$  where  $\{v_i\}$  and  $\{w_j\}$  are bases for  $S$  and  $T$  respectively.

*Proof.* Given an odd integer  $k$  as input, by Lemma 3.16 it is possible to compute the self-dual normal basis  $\{e_i\}_{i=1}^k$  together with the multiplication tables  $K_{e_i}$  for  $i = 1, 2, \dots, k$ . Addition is performed component-wise, and multiplication is done using Eq. (11). For the multiplication table  $K_a$  it suffices to compute the  $k$  products  $\kappa(ae_i)$  for  $i \in \{1, \dots, k\}$ . To compute inverses, observe that  $\kappa(1) = \kappa(aa^{-1}) = K_a \kappa(a^{-1})$ . The matrices  $K_a$  are invertible over  $\mathbb{F}_2$ , so therefore  $\kappa(a^{-1}) = K_a^{-1} \kappa(1)$ ; moreover,  $\kappa(1)$  is the all-ones vector by Lemma 3.17 and can thus be efficiently computed. Inverting the matrix can be done in  $\text{poly}(k)$  time via Gaussian elimination. The trace of an element  $a \in \mathbb{F}_{2^k}$  is by definition the trace of the multiplication table  $K_a$ .

For the ‘‘Furthermore’’ part, we observe that since  $\{v_1, v_2, \dots, v_m\} \cup \{w_1, w_2, \dots, w_{n-m}\}$  forms a basis for  $\mathbb{F}_{2^k}^n$ , there is a unique way to write  $x$  as a  $\mathbb{F}_{2^k}$  linear combination of  $\{v_i\}$  and  $\{w_j\}$ . Via Gaussian elimination over  $\mathbb{F}_{2^k}$ , the  $\mathbb{F}_2$ -representation of the coefficients of this linear combination can be computed in  $\text{poly}(n, k)$  time. Here we use that addition, multiplication and division over  $\mathbb{F}_{2^k}$  can be performed in time  $\text{poly}(k)$  using the previous items of the Lemma.  $\square$

**Remark 3.19.** Throughout this paper, whenever we refer to Turing machines that perform computations with elements of a field  $\mathbb{F}_q$  for an admissible field size  $q = 2^k$ , we mean that the Turing machines are representing elements of  $\mathbb{F}_q$  as vectors in  $\{0, 1\}^k$  using the basis specified by the algorithm of Lemma 3.16 and performing arithmetic as described in Lemma 3.18.

### 3.4 Polynomials and the low-degree code

In this section, we introduce some basic definitions about polynomials over finite fields. An  $m$ -variate polynomial  $f$  over  $\mathbb{F}_q$  is a function of the form

$$f(x_1, \dots, x_m) = \sum_{\alpha \in \{0, 1, \dots, q-1\}^m} c_\alpha x_1^{\alpha_1} \cdots x_m^{\alpha_m}$$

where  $\{c_\alpha\}$  is a collection of coefficients in  $\mathbb{F}_q$ . We say that  $f$  has *individual degree (at most)  $d$*  if  $c_\alpha \neq 0$  only if  $\alpha_i \leq d$  for all  $1 \leq i \leq m$ , and that it has *total degree (at most)  $d$*  if  $c_\alpha \neq 0$  only if  $\sum_i \alpha_i \leq d$ . (Affine) *multilinear* polynomials are polynomials with individual degree 1.

Low-degree polynomials play an important role in this paper. For one, the classical and quantum low-degree tests of Section 7 are nonlocal games that efficiently certify that the players’ answers are consistent with low-degree polynomials. Low-degree polynomials are also crucial in the probabilistically checkable proofs (PCP) construction that are used in ‘‘answer reduction’’ transformation (Section 10).

We recall the Schwartz-Zippel lemma:

**Lemma 3.20** (Schwartz-Zippel lemma [Sch80, Zip79]). *Let  $f, g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  be two unequal polynomials with total degree at most  $d$ . Then*

$$\Pr_{x \sim \mathbb{F}_q^m} [f(x) = g(x)] \leq d/q.$$

**The low-degree code.** The Schwartz-Zippel lemma implies that the set of low-degree polynomials form an error-correcting code with good distance, which we call the *low-degree code*.<sup>22</sup>

Fix an integer  $m \in \mathbb{N}$  and let  $M = 2^m$ . For every  $y \in \{0, 1\}^m$  define the following  $m$ -variate multilinear polynomial over  $\mathbb{F}_q$ :

$$\text{ind}_{m,y}(x) = \prod_{i:y_i=1} x_i \cdot \prod_{i:y_i=0} (1 - x_i).$$

Here, we identify  $\{0, 1\}$  as a subset of  $\mathbb{F}_q$ . Notice that, when restricted to the subcube  $\{0, 1\}^m$ ,  $\text{ind}_{m,y}$  is zero everywhere except for when  $x = y$ . For  $a \in \mathbb{F}_q^M$ , label the coordinates of  $a$  as  $a_y$  for  $y \in \{0, 1\}^m$  (identifying the latter set with  $\{1, \dots, M\}$  using, say, the lexicographic ordering on strings). For any such  $a$  define the multilinear polynomial  $g_a : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ , called the *low-degree encoding of  $a$* , as follows:

$$g_a(x) = \sum_{y \in \{0, 1\}^m} a_y \cdot \text{ind}_{m,y}(x). \quad (12)$$

Note that for any  $y \in \{0, 1\}^m$ ,  $g_a(y) = a_y$ . Furthermore, the map  $a \mapsto g_a$  is linear: for every  $x \in \mathbb{F}_q^m$ ,

$$g_a(x) = a \cdot \text{ind}_m(x), \quad (13)$$

where  $\text{ind}_m(x)$  is the vector  $(\text{ind}_{m,y}(x))_{y \in \{0, 1\}^m} \in \mathbb{F}_q^M$ .

**Lemma 3.21.** *Let  $q$  be an admissible field size, and let  $M = 2^m$  for some integer  $m$ . The complexity of computing the evaluation of the low-degree encoding  $g_a(x)$ , given  $a \in \mathbb{F}_q^M$  and  $x \in \mathbb{F}_q^m$  represented in binary, is  $\text{poly}(M, \log q)$ .*

*Proof.* Computing  $g_a(x)$  requires computing the sum of products  $a_y \cdot \text{ind}_{m,y}(x)$  over all  $y \in \{0, 1\}^m$ . Via Lemma 3.18, evaluating  $\text{ind}_{m,y}(x)$  takes time  $\text{poly}(m, \log q)$  because it requires performing  $m$  multiplications of  $\mathbb{F}_q$  elements, and therefore the product  $a_y \cdot \text{ind}_{m,y}(x)$  requires  $\text{poly}(m, \log q)$  time. Computing the sum over all  $y \in \{0, 1\}^m$  requires  $M \cdot \text{poly}(m, \log q) = \text{poly}(M, \log q)$  time, as claimed.  $\square$

Finally, for any  $H \subseteq \mathbb{F}_q$  we define the *decoding map*  $\text{Dec}_H(\cdot)$  which takes as input a polynomial  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  and returns the following vector  $a \in \mathbb{F}_q^M$ : for all  $y \in \{0, 1\}^m$ , if  $g(y) \in H$ , then set  $a_y = g(y)$ , and otherwise set  $a_y = 0$ . Note that for all  $a \in H^M$  we have  $\text{Dec}_H(g_a) = a$  where  $g_a$  is the low-degree encoding of  $a$ . For notational brevity we often write  $\text{Dec}(\cdot)$  to denote the boolean decoding map  $\text{Dec}_{\{0, 1\}}(\cdot)$ .

### 3.5 Linear spaces and registers

For a set  $V$ , we write  $\mathbb{C}^V$  for the complex vector space of dimension  $|V|$ . The space  $\mathbb{C}^V$  is endowed with a canonical orthonormal basis  $\{|x\rangle\}_{x \in V}$ . By “a quantum state on  $V$ ” we mean a unit vector

$$|\psi\rangle_V \in \mathbb{C}^V.$$

If  $V = \bigoplus_{i=1}^k V_k$  is the direct sum of subspaces  $V_k$  over  $\mathbb{F}$ , then  $\mathbb{C}^V$  can be identified with  $\bigotimes_{i=1}^k \mathbb{C}^{V_i}$ , by the identification of  $|x_1 \oplus \dots \oplus x_k\rangle$  with  $|x_1\rangle \otimes \dots \otimes |x_k\rangle$ . As a special case, if  $\{e_i\}$  is a basis of  $V$  the decomposition  $V = \bigoplus_{i=1}^k (\mathbb{F}e_i)$  yields the tensor product decomposition  $\mathbb{C}^V = \bigotimes_{i=1}^k \mathbb{C}^{|\mathbb{F}|}$ . We sometimes refer to the spaces  $\mathbb{C}^{|\mathbb{F}|}$  as the “qudits” of  $\mathbb{C}^V$  (or of a state on it).

<sup>22</sup>It is also known as the *generalized Reed-Muller code* in the coding theory literature.

**Definition 3.22.** For a linear space  $V$  over a finite field  $\mathbb{F}$ , define the EPR state on  $\mathbb{C}^V \otimes \mathbb{C}^V$  by

$$|\text{EPR}\rangle_V = \frac{1}{\sqrt{|V|}} \sum_{x \in V} |x\rangle \otimes |x\rangle .$$

We also write  $|\text{EPR}\rangle_{\mathbb{F}_q}$  as  $|\text{EPR}_q\rangle$  and  $|\text{EPR}_2\rangle$  as  $|\text{EPR}\rangle$ .

### 3.6 Measurements and observables

Quantum measurements are modeled as positive operator-valued measures (POVMs). A POVM consists of a set of positive semidefinite operators  $\{M_a\}_{a \in S}$  indexed by outcomes  $a \in S$  that satisfy the condition  $\sum_a M_a = I$ . If the latter condition is relaxed to  $\sum_a M_a \leq I$  then we refer to  $\{M_a\}$  as a *sub-measurement*. We sometimes use the same letter  $M$  to refer to the collection of operators defining the POVM. The probability that the measurement returns outcome  $a$  on state  $|\psi\rangle$  is given by

$$\Pr(a) = \langle \psi | M_a | \psi \rangle .$$

A POVM  $M = \{M_a\}$  is said to be *projective* if each operator  $M_a$  is a projector ( $M_a^2 = M_a$ ). This automatically implies that operators  $M_a, M_b$  for distinct outcomes  $a \neq b$  are orthogonal ( $M_a M_b = M_b M_a = 0$ ). An *observable* is a unitary matrix. A *binary observable* is an observable  $O$  such that  $O^2 = I$ , i.e.  $O$  has eigenvalues in  $\{-1, 1\}$ .

We follow the convention that subscripts of the measurement index the outcome and superscripts of the measurement are used to index different measurements. For example, we use  $\{M_{a,b}^x\}$  to represent a measurement indexed by  $x$  whose outcome consists of two parts  $a$  and  $b$ . In this case, by slightly abusing the notation, we use  $\{M_a^x\}$  and  $\{M_b^x\}$  to denote

$$M_a^x = \sum_b M_{a,b}^x, \quad M_b^x = \sum_a M_{a,b}^x .$$

For any  $x$ ,  $\{M_a^x\}$  and  $\{M_b^x\}$  are POVMs sometimes referred to as the ‘‘marginals’’ of  $\{M_{a,b}^x\}$ .

**Definition 3.23.** Let  $\{M_a^x\}_{a \in A}$  be a family of POVMs indexed by  $x \in \mathcal{X}$ . Let  $f : A \rightarrow B$  be an arbitrary function. We write  $\{M_{[f(\cdot)=b]}^x\}$  for the POVM derived from  $\{M_a^x\}$  by applying the function  $f$  before returning the outcome. More precisely,

$$M_{[f(\cdot)=b]}^x = \sum_{a: f(a)=b} M_a^x .$$

If  $b$  is not in the image of  $f$ , then we define  $M_{[f(\cdot)=b]}^x$  to be 0. In cases where the outcome  $a$  has a natural interpretation as a tuple  $(a_1, \dots, a_k)$  we sometimes slightly abuse notation and write  $M_{[a_i=b]}^x$  for some  $i \in \{1, \dots, k\}$  to denote

$$M_{[a_i=b]}^x = \sum_{(a_1, \dots, a_k): a_i=b} M_{(a_1, \dots, a_k)}^x .$$

We will make frequent use of Naimark dilation. In our setting, it can be formulated as follows. For a proof, see [JNV<sup>+</sup>20, Theorem 5.1]

**Theorem 3.24** (Naimark dilation). *Let  $|\psi\rangle$  be a state in  $\mathcal{H}_A \otimes \mathcal{H}_B$ . Let  $A = \{A_a^x\}$  be a sub-measurement acting on  $\mathcal{H}_A$  and  $B = \{B_b^y\}$  be a sub-measurement acting on  $\mathcal{H}_B$ . Then there exists*

1. Hilbert spaces  $\mathcal{H}_{A_{\text{aux}}}$  and  $\mathcal{H}_{B_{\text{aux}}}$ ,
2. a state  $|\text{aux}\rangle \in \mathcal{H}_{A_{\text{aux}}} \otimes \mathcal{H}_{B_{\text{aux}}}$ ,
3. and two measurements  $\hat{A} = \{\hat{A}_a^x\}$  and  $\hat{B} = \{\hat{B}_b^y\}$  acting on  $\mathcal{H}_A \otimes \mathcal{H}_{A_{\text{aux}}}$  and  $\mathcal{H}_B \otimes \mathcal{H}_{B_{\text{aux}}}$ , respectively,

such that the following is true. If we write  $|\hat{\psi}\rangle = |\psi\rangle \otimes |\text{aux}\rangle$ , then for all  $x, y, a, b$ ,

$$\langle \psi | A_a^x \otimes B_b^y | \psi \rangle = \langle \hat{\psi} | \hat{A}_a^x \otimes \hat{B}_b^y | \hat{\psi} \rangle.$$

In addition,  $|\text{aux}\rangle$  is a product state, meaning that we can write it as

$$|\text{aux}\rangle = |\text{aux}_A\rangle \otimes |\text{aux}_B\rangle,$$

for  $|\text{aux}_A\rangle$  in  $\mathcal{H}_{A_{\text{aux}}}$  and  $|\text{aux}_B\rangle$  in  $\mathcal{H}_{B_{\text{aux}}}$ .

The second of these is the ‘‘orthogonalization lemma’’ from [KV11]. In the setting of symmetric strategies, it states the following.

### 3.7 Generalized Pauli observables

For prime number  $p$ , the generalized Pauli operators over  $\mathbb{F}_p$  are a collection of observables indexed by a basis setting  $X$  or  $Z$  and an element  $a$  or  $b$  of  $\mathbb{F}_p$ , with eigenvalues that are  $p$ -th roots of unity. They are given by

$$\sigma^X(a) = \sum_{j \in \mathbb{F}_p} |j+a\rangle \langle j| \quad \text{and} \quad \sigma^Z(b) = \sum_{j \in \mathbb{F}_p} \omega^{bj} |j\rangle \langle j|, \quad (14)$$

where  $\omega = e^{\frac{2\pi i}{p}}$ , and addition and multiplication are over  $\mathbb{F}_p$ . These observables obey the ‘‘twisted commutation’’ relations

$$\forall a, b \in \mathbb{F}_p, \quad \sigma^X(a) \sigma^Z(b) = \omega^{-ab} \sigma^Z(b) \sigma^X(a). \quad (15)$$

Similarly, over a field  $\mathbb{F}_q$  with  $q$  a power of  $p$ , we can consider a set of generalized Pauli operators, indexed by a basis setting  $X$  or  $Z$  and an element of  $\mathbb{F}_q$  and with eigenvalues that are  $p$ -th roots of unity. For  $a, b \in \mathbb{F}_q$  they are given by

$$\tau^X(a) = \sum_{j \in \mathbb{F}_q} |j+a\rangle \langle j| \quad \text{and} \quad \tau^Z(b) = \sum_{j \in \mathbb{F}_q} \omega^{\text{tr}(bj)} |j\rangle \langle j|,$$

where addition and multiplication are over  $\mathbb{F}_q$ . For all  $W \in \{X, Z\}$ ,  $a, a' \in \mathbb{F}_q$ , and  $b \in \mathbb{F}_p$ , powers of these observables obey the relations

$$(\tau^W(a) \tau^W(a')) = \tau^W(a+a') \quad \text{and} \quad (\tau^W(a))^b = \tau^W(ab).$$

In particular, since  $pa = 0$  for any  $a \in \mathbb{F}_q$  we get that that  $(\tau^W(a))^p = I$  for any  $a \in \mathbb{F}_q$ . The observables obey analogous ‘‘twisted commutation’’ relations to (15),

$$\forall a, b \in \mathbb{F}_q, \quad \tau^X(a) \tau^Z(b) = \omega^{-\text{tr}(ab)} \tau^Z(b) \tau^X(a). \quad (16)$$

It is clear from the definition that all of the  $\tau^X$  operators commute with each other, and similarly all the  $\tau^Z$  operators commute with each other. Thus, it is meaningful to speak of a common eigenbasis for all  $\tau^X$

operators, and a common eigenbasis for all  $\tau^Z$  operators. The common eigenbasis for the  $\tau^Z$  operators is the computational basis  $\{|j\rangle\}_{j \in \mathbb{F}_q}$ . To map this basis to the common eigenbasis of the  $\tau^X$  operators, one can apply the Fourier transform

$$F = \frac{1}{\sqrt{q}} \sum_{a,b \in \mathbb{F}_q} \omega^{-\text{tr}(ab)} |a\rangle \langle b|. \quad (17)$$

Explicitly, the eigenbases consist of the vectors  $|e_W\rangle$  labeled by an element  $e \in \mathbb{F}_q$  and  $W \in \{X, Z\}$ , given by

$$|e_X\rangle = \frac{1}{\sqrt{q}} \sum_j \omega^{-\text{tr}(ej)} |j\rangle, \quad |e_Z\rangle = |e\rangle.$$

We denote the POVM whose elements are projectors onto basis vectors of the eigenbasis associated with the observables  $\tau^W$  by  $\{\tau_e^W\}_e$ . Then for all  $W \in \{X, Z\}$  and  $a \in \mathbb{F}_q$ , the observables  $\tau^W(a)$  can be written as

$$\tau^W(a) = \sum_{b \in \mathbb{F}_q} \omega^{\text{tr}(ab)} \tau_b^W. \quad (18)$$

To invert this relation, we first quote the following useful Fourier fact.

**Lemma 3.25** (Fact 3.2 of [NW19]). *Let  $V$  be a subspace of  $\mathbb{F}^k$ . For all  $v \notin V^\perp$ ,*

$$\mathbb{E}_{u \sim V} \omega^{\text{tr}(u \cdot v)} = 0,$$

where the expectation is over a uniformly random vector  $u$  from  $V$ .

We may now invert Equation (18):

$$\mathbb{E}_{a \in \mathbb{F}_q} \omega^{-\text{tr}(ab)} \tau^W(a) = \sum_{b' \in \mathbb{F}_q} \mathbb{E}_{a \in \mathbb{F}_q} \omega^{\text{tr}(a(b'-b))} \tau_{b'}^W = \tau_b^W, \quad (19)$$

where the second step follows from Lemma 3.25.

For systems with many qudits, we will consider tensor products of the operators  $\tau^W$ . Slightly abusing notation, for  $W \in \{X, Z\}$  and  $a \in \mathbb{F}_q^n$  we denote by  $\tau^W(a)$  the tensor product  $\tau^W(a_1) \otimes \dots \otimes \tau^W(a_n)$ . These obey the twisted commutation relations

$$\forall a, b \in \mathbb{F}_q^n, \quad \tau^X(a) \tau^Z(b) = \omega^{-\text{tr}(a \cdot b)} \tau^Z(b) \tau^X(a),$$

where  $a \cdot b = \sum_{i=1}^n a_i b_i \in \mathbb{F}_q$ . For  $W \in \{X, Z\}$  and  $e \in \mathbb{F}_q^n$  define the eigenstates

$$|e_W\rangle = |(e_1)_W\rangle \otimes \dots \otimes |(e_n)_W\rangle,$$

and associated rank-1 projectors  $\tau_e^W$ . Analogous versions of Equation (18) and Equation (19) relate the multi-qubit observables and projectors:

$$\tau^W(a) = \sum_{b \in \mathbb{F}_q^n} \omega^{\text{tr}(a \cdot b)} \tau_b^W, \quad (20)$$

$$\mathbb{E}_{a \in \mathbb{F}_q^n} \omega^{-\text{tr}(a \cdot b)} \tau^W(a) = \sum_{b' \in \mathbb{F}_q^n} \mathbb{E}_{a \in \mathbb{F}_q^n} \omega^{\text{tr}(a \cdot (b'-b))} \tau_{b'}^W = \tau_b^W. \quad (21)$$

Since we only consider finite fields  $\mathbb{F}_q$  such that  $q = 2^k$  the maximally entangled state  $|\text{EPR}_q\rangle$  and the corresponding qudit Pauli observables/projectors are isomorphic to a tensor product of maximally entangled states  $|\text{EPR}_2\rangle$  and *qubit* Pauli observables/projectors respectively; this is shown in the next lemma. This is used to argue that the Pauli basis test (described in Section 7.3) gives a self-test for Pauli observables and maximally entangled states over qubits.

**Lemma 3.26.** For all admissible field sizes  $q = 2^k$  and integers  $L$ , there exists an isomorphism  $\phi : (\mathbb{C}^q)^{\otimes L} \rightarrow (\mathbb{C}^2)^{\otimes Lk}$  such that<sup>23</sup>

$$\phi \otimes \phi |\text{EPR}_q\rangle^{\otimes L} = |\text{EPR}_2\rangle^{\otimes Lk}. \quad (22)$$

Moreover, for all  $W \in \{X, Z\}$  and for all  $u \in \mathbb{F}_q^L$

$$\tau_u^W = \phi^\dagger \left( \bigotimes_{i=1}^L \bigotimes_{j=1}^k \sigma_{u_{ij}}^W \right) \phi. \quad (23)$$

Here, the  $(u_{ij})_{i,j}$  denotes a vector of  $\mathbb{F}_2$  values such that  $u_i = \sum_j u_{ij} e_j$  for all  $i \in \{1, \dots, L\}$  with  $\{e_1, \dots, e_k\}$  being the self-dual normal basis of  $\mathbb{F}_q$  over  $\mathbb{F}_2$  specified by Lemma 3.16. For  $i \in \{1, \dots, L\}$  and  $j \in \{1, \dots, q\}$  the  $(i, j)$ -th factor  $\sigma_{u_{ij}}^W$  denotes the projector  $\frac{1}{2} (I + (-1)^{u_{ij}} \sigma^W(1))$  acting on the  $s$ -th qubit of  $|\text{EPR}_2\rangle^{\otimes Lk}$ , where  $s = (i-1)k + j$ .

*Proof.* Define the isometry  $\theta : \mathbb{C}^q \rightarrow (\mathbb{C}^2)^{\otimes k}$  as  $\theta |a\rangle = |a_1\rangle \otimes |a_2\rangle \cdots |a_k\rangle$  where  $\kappa(a) = (a_1, a_2, \dots, a_k) \in \mathbb{F}_2^k$  is the bijection introduced in Section 3.3 corresponding to the basis  $\{e_1, \dots, e_k\}$ . Observe that  $\theta \otimes \theta |\text{EPR}_{2^k}\rangle = |\text{EPR}_2\rangle^{\otimes k}$ .

Let  $a \in \mathbb{F}_q$ , and let  $\kappa(a) = (a_1, \dots, a_k) \in \mathbb{F}_2^k$ . Then from (21), we get

$$\begin{aligned} \tau_a^W &= \mathbb{E}_{b \in \mathbb{F}_q} (-1)^{\text{tr}(ab)} \tau^W(b) \\ &= \mathbb{E}_{b \in \mathbb{F}_q} (-1)^{\text{tr}(\sum_j a_j e_j b)} \tau^W(b) \\ &= \mathbb{E}_{b \in \mathbb{F}_q} (-1)^{\sum_j a_j b_j} \tau^W \left( \sum_j b_j e_j \right), \end{aligned} \quad (24)$$

where  $b = \sum_j b_j e_j$ ; since the basis  $\{e_j\}$  is self-dual, we have that  $b_j = \text{tr}(b e_j)$ . From (24) and the product relation in (3.7) we get that

$$\begin{aligned} \tau_a^W &= \mathbb{E}_{b_1, \dots, b_k \in \mathbb{F}_2} \prod_{j=1}^k (-1)^{a_j b_j} \tau^W(b_j e_j) \\ &= \prod_{j=1}^k \mathbb{E}_{b_j \in \mathbb{F}_2} (-1)^{a_j b_j} \tau^W(b_j e_j), \end{aligned} \quad (25)$$

where we have used the fact that the expectation is over iid  $b_1, \dots, b_k$  to take it inside the product in the second line. Next we claim that for all  $c \in \mathbb{F}_2$ , we have  $\tau^W(c e_j) = \theta^\dagger \sigma^{W,j}(c) \theta$  where  $\sigma^{W,j}(c) = I$  if  $c = 0$ , and otherwise is the Pauli  $W$  observable acting on the  $j$ -th qubit of  $(\mathbb{C}^2)^{\otimes k}$ . This can be verified by comparing the actions of both operators on the basis states of  $\mathbb{C}^q$ .

Thus we obtain that the right-hand side of (25) is equal to

$$\prod_{j=1}^k \mathbb{E}_{b_j \in \mathbb{F}_2} (-1)^{a_j b_j} \left[ \theta^\dagger \sigma^{W,j}(b_j) \theta \right] = \theta^\dagger \bigotimes_{j=1}^k \left( \mathbb{E}_{b_j \in \mathbb{F}_2} (-1)^{a_j b_j} \sigma^W(b_j) \right) \theta \quad (26)$$

$$= \theta^\dagger \left( \bigotimes_{j=1}^k \sigma_{b_j}^W \right) \theta. \quad (27)$$

<sup>23</sup>Here we have applied the natural identifications between  $(\mathbb{C}^q)^{\otimes L} \otimes (\mathbb{C}^q)^{\otimes L}$  and  $(\mathbb{C}^q \otimes \mathbb{C}^q)^{\otimes L}$ , and between  $(\mathbb{C}^2)^{\otimes Lk} \otimes (\mathbb{C}^2)^{\otimes Lk}$  and  $(\mathbb{C}^2 \otimes \mathbb{C}^2)^{\otimes Lk}$ .

Define  $\phi = \theta^{\otimes L}$ . It is evident that (22) holds for this choice of  $\phi$ , and it remains to show (23). The projector  $\tau_u^W$  can be decomposed as the tensor product  $\bigotimes_{i=1}^L \tau_{u_i}^W$  where  $\tau_{u_i}^W$  acts on the  $i$ -th factor of  $(\mathbb{C}^q)^{\otimes L}$ . Express each  $u_i$  as  $\sum_j u_{ij} e_j$  where  $u_{ij} \in \mathbb{F}_2$ . Then from Equation (27) we get that

$$\tau_u^W = \bigotimes_{i=1}^L \tau_{u_i}^W = \phi^\dagger \left( \bigotimes_{i=1}^L \bigotimes_{j=1}^k \sigma_{u_{ij}}^W \right) \phi, \quad (28)$$

which establishes Equation (23). □



## 4 Conditionally Linear Functions, Distributions, and Samplers

### 4.1 Conditionally linear functions and distributions

We first introduce *conditionally linear functions*, which are used to specify the question distribution for games considered in the paper in a way that the question distribution can be “introspected”, as described in Section 8. Intuitively, a conditionally linear function takes as input an element  $x \in V = \mathbb{F}^n$  for some  $n \geq 0$ , and applies linear maps  $L_j$  sequentially on  $x^{V_j}$  where  $V_1, V_2, \dots$  are a sequence of complementary register subspaces such that both the linear maps  $L_j$  and the subspace  $V_j$  for  $j \geq 2$  may depend on the values taken by previous linear maps  $L_1(x^{V_1}), L_2(x^{V_2}), \dots$ , etc.

In the remainder of the section we use  $V$  to denote the linear space  $\mathbb{F}^n$  for some integer  $n \geq 0$ . For ease of notation we extensively use the subscript range notation. For example, if  $V_1, V_2, \dots, V_\ell$  are fixed subspaces of  $V$  and  $k \in \{1, 2, \dots, \ell\}$  we write

$$V_{<k} = \bigoplus_{j:1 \leq j < k} V_j, \quad V_{>k} = \bigoplus_{j:\ell \geq j > k} V_j,$$

and it is understood that  $V_{\leq k}$  and  $V_{\geq k}$  are identical to  $V_{<k+1}$  and  $V_{>k-1}$ , respectively. Moreover, if  $V'$  is a register subspace of  $V$ ,  $F : V' \rightarrow V'$  is a function, and  $x \in V$ , we write  $x^F$  to denote  $F(x^{V'})$ . For example, in the following definition  $x^{L_1}$  is used as shorthand notation for  $L_1(x^{V_1})$ .

**Definition 4.1.** Let  $V$  be  $\mathbb{F}^n$  for some  $n \geq 0$ . For all integers  $\ell \geq 0$  the collection of  $\ell$ -level *conditionally linear functions* (implicitly, *on  $V$* ) is defined inductively as follows.

1. There is a single 0-level conditionally linear function, which is the 0 function on  $V$ .
2. Let  $\ell \geq 1$  and suppose the collection of  $(\ell - 1)$ -level conditionally linear functions has been defined. The collection of  $\ell$ -level conditionally linear functions on  $V$  consists of all functions  $L$  on  $V$  that can be expressed in the following form. There exist complementary register subspaces  $V_1$  and  $V_{>1}$  of  $V$ , a linear function  $L_1$  on  $V_1$ , and for all  $v \in L_1(V_1)$ , an  $(\ell - 1)$ -level conditionally linear function  $L_{>1, v}$  on  $V_{>1}$ , such that for all  $x \in V$ ,

$$L(x) = x^{L_1} + L_{>1, x^{L_1}}(x^{V_{>1}}).$$

The concept of conditionally linear functions is simple even though the notations seem complicated. It is best understood using an operational definition illustrated in Fig. 1. We give an example of a 2-level CL function in Example 4.3.

**Remark 4.2.** *It follows from the above definition that all 1-level CL functions are linear.*

**Example 4.3.** Consider  $V = \mathbb{F}_2^3$ . Then the following map

$$L : (x_0, x_1, x_2) \mapsto (0, x_0x_2 + x_1x_2 + x_0, x_2)$$

on  $V$  is a 2-level CL function by choosing  $V_1 = \text{span}\{(0, 0, 1)\}$ ,  $L_1$  the identity function on  $V_1$ ,  $V_2 = V_1^\perp$ , and

$$L_{2, x_2} : (x_0, x_1, 0) \mapsto (0, x_0x_2 + x_1x_2 + x_0, 0),$$

which is linear for all  $x_2 \in \mathbb{F}_2$ . Note that intuitively the function  $L$  selects either  $x_0$  or  $x_1$  in the second coordinate conditioning on the value of  $x_2$ .




Figure 1: An illustration of an  $\ell$ -level CL function  $L$ . First a register subspace  $V_1$  and a linear map  $L_1$  are chosen and applied to obtain  $x^{L_1} = L_1(x^{V_1}) \in V_1$ . Then depending the value of  $x^{L_1}$ , a register subspace  $V_2$  and a linear map  $L_{2, x^{L_1}}$  are chosen and applied to obtain  $x^{L_2} = L_{2, x^{L_1}}(x^{V_2}) \in V_2$  and so on. Finally,  $L(x)$  is defined to be  $\sum_{k=1}^{\ell} x^{L_k}$ .

**Remark 4.4.** Note that for any integer  $\ell \geq 1$  the collection of  $\ell$ -level CL functions trivially contains the collection of  $(\ell - 1)$ -level CL functions: for this it suffices to note that the 0 function, which is a 0-level CL function, is also a 1-level CL function by setting  $V_1 = V$ ,  $V_{>1} = \{0\}$ ,  $L_1(x) = 0$  for all  $x \in V$ , and  $L_{>1, x^{L_1}}$  is the 0 map for all  $x \in V$ .

**Definition 4.5.** Let  $L, R : V \rightarrow V$  be conditionally linear functions. The *conditionally linear distribution*  $\mu_{L,R}$  corresponding to  $(L, R)$  is defined as the distribution over pairs  $(L(x), R(x)) \in V \times V$  for  $x$  drawn uniformly at random from  $V$ .

Throughout the paper we abbreviate “conditionally linear functions” and “conditionally linear distributions” as *CL functions* and *CL distributions*, respectively.

The following lemma elucidates structural properties of  $\ell$ -level CL functions. Recall that using our shorthand notation,  $x^{L_{<k}}$  and  $x^{L_k}$  in the lemma denote  $L_{<k}(x)$  and  $L_{k,u}(x^{V_{k,u}})$  where  $u = L_{<k}(x)$ .

**Lemma 4.6.** Let  $\ell \geq 1$  and  $V = \mathbb{F}^n$  for some integer  $n \geq 0$ . A function  $L : V \rightarrow V$  is an  $\ell$ -level CL function if and only if the following collection of functions and subspaces exists:

- (i) For each  $k \in \{1, 2, \dots, \ell\}$ , a function  $L_{\leq k} : V \rightarrow V$  called the  $k$ -th marginal of  $L$ ;
- (ii) For each  $k \in \{1, 2, \dots, \ell\}$  and  $u \in L_{<k}(V)$ , a register subspace  $V_{k,u}$  of  $V$  called the  $k$ -th factor space with prefix  $u$ ;
- (iii) For each  $k \in \{1, 2, \dots, \ell\}$  and  $u \in L_{<k}(V)$ , a linear map  $L_{k,u} : V_{k,u} \rightarrow V_{k,u}$  called the  $k$ -th linear map of  $L$  with prefix  $u$ ;

such that the following conditions hold for all  $k \in \{1, 2, \dots, \ell\}$ .

1.  $L_{\leq k}$  is a  $k$ -level CL function on  $V$ ;

2.  $V = \bigoplus_{i=1}^{\ell} V_{i,x^{L_{<i}}}$  for all  $x \in V$ ;
3.  $L_{\leq k}(x) = \sum_{i=1}^k x^{L_i}$  for all  $x \in V$ , where  $L_i$  is shorthand notation for  $L_{i,x^{L_{<i}}}$ ;
4.  $L = L_{\leq \ell}$ .

As in Item 3, we sometimes use  $V_k$  and  $L_k$  to denote  $V_{k,u}$  and  $L_{k,u}$  respectively, leaving the prefix  $u$  implicit.

*Proof.* We first prove the “if” direction: if there exist spaces and functions satisfying the conditions in the lemma, the fact that  $L$  is an  $\ell$ -level CL function follows from Items 1 and 4 of the lemma statement.

We now prove the “only if” direction. Given a CL function  $L$  on  $V$ , we construct the  $k$ -th family of subspaces and functions for all  $k \in \{1, \dots, \ell\}$  by induction on the level  $\ell$ . First consider the base case  $\ell = 1$ . Since  $L_{<1} = 0$ , we omit the mentioning of the prefix  $u \in L_{<1}(V)$ . Define  $L_{\leq 1} = L$ , the factor space  $V_1 = V$ , and the linear map  $L_1 = L$ . It is straightforward to verify that the conditions in the lemma hold for these choices of linear maps and spaces.

Now, assume that the lemma holds for CL functions of level at most  $\ell - 1$ , and we prove the lemma for  $\ell$ -level CL functions. By definition, an  $\ell$ -level CL function  $L$  can be written as

$$L(x) = x^{L_1} + L_{>1,x^{L_1}}(x^{V_{>1}})$$

for some linear map  $L_1 : V_1 \rightarrow V_1$  and a family of  $(\ell - 1)$ -level CL functions

$$\{L_{>1,v} : V_{>1} \rightarrow V_{>1}\}_{v \in L_1(V_1)}$$

where  $V_1$  and  $V_{>1}$  are complementary register subspaces of  $V$ . Next, using the inductive hypothesis on the  $(\ell - 1)$ -level CL function  $L_{>1,v}$  we get that for all  $v \in L_1(V_1)$  and all  $k \in \{1, 2, \dots, \ell - 1\}$  there exist  $k$ -th marginal functions  $L'_{v,\leq k} : V_{>1} \rightarrow V_{>1}$ ,  $k$ -th factor spaces  $V'_{v,k,u}$ , and  $k$ -th linear maps  $L'_{v,k,u}$  of  $L_{>1,v}$  with prefix  $u \in L'_{v,<k}(V_{>1})$  such that the conditions of the lemma for  $L_{>1,v}$  hold.

Define the marginal functions  $L_{\leq k} : V \rightarrow V$ , factor spaces  $V_{k,u}$  and linear maps  $L_{k,u}$  for  $L$  as follows.

(i) Define  $L_{\leq 1} = L_1$  and the first factor space to be  $V_1$ ;

(ii) For all  $k \in \{2, 3, \dots, \ell\}$ , define

$$L_{\leq k} : x \mapsto x^{L_1} + L'_{x^{L_1},<k}(x^{V_{>1}}) \quad \text{for } x \in V; \quad (29)$$

(iii) For all  $k \in \{2, 3, \dots, \ell\}$  and  $u \in L_{<k}(V)$ , define  $V_{k,u} = V'_{v,k-1,w}$  and  $L_{k,u} = L'_{v,k-1,w}$  where  $v = u^{V_1} \in L_1(V_1)$  and  $w = u^{V_{>1}} \in L'_{v,<k-1}(V_{>1})$ .

We verify that the conditions of the lemma are satisfied. Since  $L'_{v,<k}$  is by assumption a  $(k - 1)$ -level CL function on  $V_{>1}$ , we get that  $L_{\leq k}$  is a  $k$ -level CL function on  $V$  from Eq. (29), establishing Item 1. By the induction hypothesis, we have for all  $v \in L_1(V_1)$  and  $y \in V_{>1}$ ,

$$V_{>1} = \bigoplus_{i=1}^{\ell-1} V'_{v,i,y^{L'_{v,<i}}}, \quad (30)$$

which implies that for all  $x \in V$  and  $v = x^{L_1}$ ,

$$V = V_1 \oplus \left( \bigoplus_{i=1}^{\ell-1} V'_{v,i,x^{L'_{v,<i}}} \right) = V_1 \oplus \left( \bigoplus_{i=2}^{\ell} V_{i,x^{L_{<i}}} \right) = \bigoplus_{i=1}^{\ell} V_{i,x^{L_{<i}}}.$$

The first equality follows from Eq. (30) while the second and third equalities follow from the definition of  $V_{k,u}$ . This establishes Item 2.

Next, we have that for all  $x \in V$ ,  $v = x^{L_1}$ , and  $k \in \{1, 2, \dots, \ell\}$ ,

$$L_{\leq k}(x) = v + L'_{v, < k}(x^{V_{>1}}) \quad (31)$$

$$= v + \sum_{i=1}^{k-1} x^{L'_{v,i}} = \sum_{i=1}^k x^{L_i}, \quad (32)$$

where  $L'_{v,i}$  is the  $i$ -th linear map of  $L_{>1,v}$  with prefix  $L'_{v, < i}(x)$  and  $L_i$  is the  $i$ -th linear map of  $L$  with prefix  $L_{< i}(x)$ . The second equality follows from the inductive hypothesis applied to  $L'_{v, < k}$  and the third equality follows from the definition of  $L_i$ . Line (32) implies Item 3 of the lemma.

Finally, Item 4 follows from (31) where we set  $k = \ell$  and observe that  $L'_{x^{L_1}, \leq \ell-1}$  is equal to  $L_{>1, x^{L_1}}$  by the inductive hypothesis. This shows that  $L_{\leq k}$ ,  $V_{k,u}$ , and  $L_{k,u}$  satisfy the conditions of the lemma and completes the induction.  $\square$

We note that the marginal functions, factor spaces, and linear maps of a given CL function  $L$  may not be unique; for example, consider the identity function on a linear space  $V = \mathbb{F}^n$ . This is clearly a 1-level CL function, but it can also be viewed as a  $k$ -level CL function for  $k \in \{2, \dots, n\}$  with an arbitrary partition of  $V$  into factor spaces.

**Lemma 4.7.** *Let  $\ell, k \geq 0$  be integers and  $U = \mathbb{F}^n$ ,  $V = \mathbb{F}^m$  be linear spaces. Suppose  $L$  is a  $k$ -level CL function on  $U$  and  $R_u$  is an  $\ell$ -level CL function on  $V$  for each  $u \in L(U)$ . Then the concatenation  $T$  of  $L$  and  $\{R_u\}_u$  defined as*

$$T(x) = L(x^U) + R_{L(x^U)}(x^V)$$

is a  $(k + \ell)$ -level conditionally linear function on  $U \oplus V$ .

*Proof.* We prove the claim by induction on  $k$ . The case  $k = 0$  follows from the Definition 4.1. Assume that the lemma holds for  $L$  being at most  $(k - 1)$ -level. By Definition 4.1, there are complementary register subspaces  $U_1$  and  $U_{>1}$  of  $U$ , a linear function  $L_1$  on  $U_1$ , and a family of  $(k - 1)$ -level CL functions  $L_{>1,v}$  for  $v \in L_1(U_1)$  such that

$$L(x^U) = x^{L_1} + L_{>1, x^{L_1}}(x^{U_{>1}}).$$

For all  $x^{L_1}$ , define the function  $T_{>1, x^{L_1}}$  on  $U_{>1} \oplus V$  as

$$T_{>1, x^{L_1}}(x^{U_{>1} \oplus V}) = L_{>1, x^{L_1}}(x^{U_{>1}}) + R_{x^{L_1} + x^{L_1}}(x^V),$$

the concatenation of  $L_{>1, x^{L_1}}$  and  $\{R_{L(x^U)}\}_{x^{L_1}}$  where  $L_{>1}$  is the shorthand notation of  $L_{>1, x^{L_1}}$ . By the induction hypothesis,  $T_{>1, x^{L_1}}$  is  $(k + \ell - 1)$ -level conditionally linear. The lemma follows from Definition 4.1.  $\square$

**Lemma 4.8** (Direct sums of CL functions). *Let  $V^{(1)}, V^{(2)}, \dots, V^{(m)}$  be register subspaces of  $V$  such that  $V = \bigoplus_{j=1}^m V^{(j)}$ . Suppose that, for each  $j \in \{1, 2, \dots, m\}$ ,  $L^{(j)}$  is an  $\ell_j$ -level conditionally linear function on  $V^{(j)}$ . Then the direct sum  $L = \bigoplus_{j=1}^m L^{(j)}$  is an  $\ell$ -level CL function over  $V$  for  $\ell = \max_j \{\ell_j\}$ , where  $L$  is defined by*

$$L(x) = \sum_{j=1}^m L^{(j)}(x^{(j)})$$

for all  $x = \sum_{j=1}^m x^{(j)} \in \bigoplus_{j=1}^m V^{(j)}$ .

*Proof.* It is easy to see that an  $\ell$ -level CL function is also  $k$ -level conditionally linear for all  $k \geq \ell$ . Hence, it suffices to prove the claim where  $\ell_j = \ell$  for  $j = 1, 2, \dots, m$ .

We prove the theorem by an induction on  $\ell$ . For  $\ell = 1$ , the functions  $L^{(j)}$  are linear and the claim follows by the fact that the direct sum of linear maps is linear.

Assume now the theorem holds for conditionally linear functions of level at most  $\ell - 1$  and  $L^{(j)}$  are  $\ell$ -level conditionally linear functions for  $j = 1, 2, \dots, m$ . By definition,  $L^{(j)}$  is the concatenation of conditionally linear functions  $L_1^{(j)}$  on  $V_1^{(j)}$  and  $\{L_{>1, v_j}^{(j)}\}_{v_j}$  on  $V_{>1}^{(j)}$  of levels 1, and  $\ell - 1$  respectively. Furthermore,

$$L(x) = \sum_{j=1}^m L^{(j)}(x^{(j)}) = \sum_{j=1}^m \left( v_j + L_{>1, v_j}^{(j)} \left( (x^{(j)})^{V_{>1}^{(j)}} \right) \right),$$

where  $v_j = L_1^{(j)} \left( (x^{(j)})^{V_1^{(j)}} \right)$ . By the induction hypothesis,

$$L_1(x^{V_1}) = \sum_{j=1}^m L_1^{(j)} \left( (x^{(j)})^{V_1^{(j)}} \right), \quad L_{>1, v}(x^{V_{>1}}) = \sum_{j=1}^m L_{>1, v_j}^{(j)} \left( (x^{(j)})^{V_{>1}^{(j)}} \right)$$

are 1-level and  $(\ell - 1)$ -level conditionally linear respectively for  $v = \sum_j v_j$ ,  $V_1 = \bigoplus_{j=1}^m V_1^{(j)}$ , and  $V_{>1} = \bigoplus_{j=1}^m V_{>1}^{(j)}$ . This proves that  $L$  is  $\ell$ -level conditionally linear.  $\square$

**Lemma 4.9.** *For each  $i \in \{1, \dots, m\}$  let  $L^{(i)}, R^{(i)} : V^{(i)} \rightarrow V^{(i)}$  be  $\ell_i$ -level conditionally linear functions and let  $L, R : V \rightarrow V$  be the direct sums  $L = \bigoplus_i L_i$  and  $R = \bigoplus_i R_i$ , respectively, as defined in Lemma 4.8. Then the conditionally linear distribution  $\mu_{L,R}$  is the product distribution  $\prod_{i=1}^m \mu_{L^{(i)}, R^{(i)}}$  over  $V \times V$ .*

*Proof.* The distribution  $\mu_{L,R}$  is the distribution over pairs  $(L(x), R(x))$  where  $x$  is sampled uniformly from  $V \times V$ . By Lemma 4.8, this is equivalent to the distribution over pairs  $((L_i(x^{V_i}))_{i=1}^m, (R_i(x^{V_i}))_{i=1}^m)$  where  $x$  is chosen uniformly at random from  $V$ . This distribution is exactly the product of the distributions  $\mu_{L^{(i)}, R^{(i)}}$  for  $i = 1, 2, \dots, m$ .  $\square$

CL functions used in the paper are frequently defined over a “large” field  $\mathbb{F}_{2^t}$  (e.g., the CL functions used in the low degree tests of Section 7). However, the introspection protocol in Section 8 handles CL functions defined over  $\mathbb{F}_2$ . The following definition and lemma show that CL functions over prime power fields can be viewed as CL functions over the prime field via a “downsizing” operation.

**Definition 4.10** (Downsizing CL functions). Let  $V = \mathbb{F}_q^n$  be a linear space for a prime power  $q = p^t$  for odd  $t$ . Let  $L : V \rightarrow V$  be a function. Let  $\kappa(\cdot)$  denote the downsize map from Section 3.3 corresponding to the basis  $\{e_1, \dots, e_t\}$  of  $\mathbb{F}_q$  over  $\mathbb{F}_p$  specified by Lemma 3.16. In particular,  $\kappa$  is linear over  $\mathbb{F}_p$ , and by Lemma 3.14, the set  $\kappa(V)$  is the linear space  $\mathbb{F}_p^{nt}$ . Define the *downsized function*  $L^\kappa : \kappa(V) \rightarrow \kappa(V)$  by  $L^\kappa = \kappa \circ L \circ \kappa^{-1}$ .

**Lemma 4.11.** *Let  $V = \mathbb{F}_q^n$  for a prime power  $q = p^t$ . Let  $L : V \rightarrow V$  be an  $\ell$ -level CL function over  $V$  for some integer  $\ell \geq 0$ . Let  $L_{\leq j}$ ,  $V_{j,u}$ , and  $L_{j,u}$  denote the  $j$ -th marginal functions, factor spaces, and linear maps corresponding to  $L$  as guaranteed by Lemma 4.6. Then  $L^\kappa : \kappa(V) \rightarrow \kappa(V)$  is an  $\ell$ -level CL function on  $V^\kappa = \kappa(V) = \mathbb{F}_p^{nt}$  with marginal functions  $L_{\leq j}^\kappa$ , factor spaces  $V_{j,v}^\kappa$ , and linear maps  $L_{j,v}^\kappa$  that satisfy the following for all  $j \in \{1, \dots, \ell\}$ .*

1. The  $j$ -th marginal function  $L_{\leq j}^\kappa$  of  $L^\kappa$  is equal to  $\kappa \circ L_{\leq j} \circ \kappa^{-1}$ .

2. For all  $u \in L_{<j}(V)$ , the  $j$ -th factor space  $V_{j,\kappa(u)}^\kappa$  and the  $j$ -th linear map  $L_{j,\kappa(u)}^\kappa$  of  $L^\kappa$  are equal to  $\kappa(V_{j,u})$  and  $\kappa \circ L_{j,\kappa(u)} \circ \kappa^{-1}$  respectively.

*Proof.* We prove the lemma by induction on  $\ell$ . Let  $L : V \rightarrow V$  be an  $\ell$ -level CL function. For the base case  $\ell = 1$ , observe that since  $\kappa$  is a linear bijection between  $\mathbb{F}_q$  and  $\mathbb{F}_p^t$  as linear spaces over  $\mathbb{F}_p$ , the function  $L^\kappa$  is linear, and thus a 1-level CL function over  $\kappa(V) = \mathbb{F}_p^{nt}$ . Furthermore, the first marginal function  $L_{\leq 1}^\kappa = L^\kappa = \kappa \circ L_{\leq 1} \circ \kappa^{-1}$ ; the factor space  $V_1^\kappa = \kappa(V_1) = \kappa(V)$ , and  $L_1^\kappa = L = \kappa \circ L_1 \circ \kappa^{-1}$ .

Assume that the statement of the lemma holds for some  $\ell - 1 \geq 1$ . Let  $L_{\leq j}$ ,  $V_{j,u}$ , and  $L_{j,u}$  denote the marginal functions, factor spaces, and linear maps corresponding to  $L$  as guaranteed by Lemma 4.6. Recursively define the following functions and spaces, for  $j \in \{1, \dots, \ell\}$ .

1.  $L_{\leq j}^\kappa = \kappa \circ L_{\leq j} \circ \kappa^{-1}$ .
2. For all  $u \in L_{<j}(V)$ , set  $V_{j,\kappa(u)}^\kappa = \kappa(V_{j,u})$  and set  $L_{j,\kappa(u)}^\kappa = \kappa \circ L_{j,\kappa(u)} \circ \kappa^{-1}$ .

We argue that  $\{L_{\leq j}^\kappa\}$ ,  $\{V_{j,v}^\kappa\}$ , and  $\{L_{j,v}^\kappa\}$  satisfy the conditions of Lemma 4.6 for the function  $L^\kappa$ , which implies that  $L^\kappa$  is an  $\ell$ -level CL function over  $\kappa(V)$ .

We first establish Item 4 of Lemma 4.6. Since  $L_{\leq \ell} = L$ , this implies

$$L^\kappa = \kappa \circ L \circ \kappa^{-1} = \kappa \circ L_{\leq \ell} \circ \kappa^{-1} = L_{\leq \ell}^\kappa,$$

as desired. Next, for all  $j \in \{1, 2, \dots, \ell\}$ , for all  $y \in \kappa(V)$  with  $y = \kappa(x)$  for some  $x \in V$ , letting  $u_j = L_{<j}(x)$ , we have

$$\kappa(V) = \kappa\left(\bigoplus_{j=1}^{\ell} V_{j,x^{L_{<j}}}\right) = \bigoplus_{j=1}^{\ell} \kappa(V_{j,x^{L_{<j}}}) = \bigoplus_{j=1}^{\ell} V_{j,\kappa(x^{L_{<j}})}^\kappa.$$

The first equality follows from Item 2 of Lemma 4.6, the second equality follows from Lemma 3.14, and the third equality follows by definition. Since  $\kappa(x^{L_{<j}}) = L_{<j}^\kappa \circ \kappa(x) = L_{<j}^\kappa(y)$ , this establishes Item 2 of Lemma 4.6.

Next, we have for all  $j \in \{1, 2, \dots, \ell\}$  and all  $y \in \kappa(V)$  with  $y = \kappa(x)$  for some  $x \in V$ ,

$$\begin{aligned} L_{\leq j}^\kappa(y) &= (\kappa \circ L_{\leq j} \circ \kappa^{-1})(y) = (\kappa \circ L_{\leq j})(x) = \sum_{i=1}^j \kappa(x^{L_{i,x^{L_{<i}}}}) \\ &= \sum_{i=1}^j L_{i,v_i}^\kappa(\kappa(x^{V_{i,x^{L_{<i}}}})) = \sum_{i=1}^j L_{i,v_i}^\kappa(y^{V_{i,v_i}^\kappa}) \end{aligned}$$

where  $v_i = L_{<i}^\kappa(y) = \kappa(x^{L_{<i}})$ . The first equality follows from definition of  $L_{\leq j}^\kappa$ , the second equality follows from  $y = \kappa(x)$ , the third equality follows from Item 3 of Lemma 4.6 applied to  $L_{\leq j}$ , the fourth equality follows from the definition of the linear map  $L_{i,v_i}^\kappa$ , and the fifth equality follows from Lemma 3.14. This establishes Item 3 of Lemma 4.6 for  $L_{\leq j}^\kappa$ .

Finally, since  $L_{\leq j}$  is a  $j$ -level CL function over  $V$ , using the inductive hypothesis we have that  $L_{\leq j}^\kappa$  is a  $j$ -level CL function over  $\kappa(V)$  when  $j \in \{1, 2, \dots, \ell - 1\}$ . It remains to establish that  $L_{\leq \ell}^\kappa$  is an  $\ell$ -level CL function. Since  $L$  is an  $\ell$ -level CL function, there exists register subspaces  $V_1, V_{>1}$  such that  $V = V_1 \oplus V_{>1}$ , a linear map  $L_1 : V_1 \rightarrow V_1$  and a collection of  $(\ell - 1)$ -level CL functions  $\{L_{>1,v} : V_{>1} \rightarrow V_{>1}\}_{v \in L_1(V_1)}$

such that  $L(x) = x^{L_1} + L_{>1, x^{L_1}}(x^{V_{>1}})$  for all  $x \in V$ . Observe that  $L_1^\kappa$  is a 1-level CL function on  $V_1^\kappa = \kappa(V_1)$ , and for  $v' = \kappa(v) \in L_1^\kappa(V_1^\kappa)$ , the inductive hypothesis implies the function  $L_{>1, v'}^\kappa$  is an  $(\ell - 1)$ -level CL function on  $V_{>1}^\kappa = \kappa(V_{>1})$ . Furthermore, since  $L_{\leq \ell}^\kappa = L^\kappa$ , we have that for all  $y \in \kappa(V)$  with  $y = \kappa(x)$  for some  $x \in V$ ,

$$L_{\leq \ell}^\kappa(y) = L^\kappa(y) = L_1^\kappa(y) + L_{>1, L_1^\kappa(y)}^\kappa(y^{V_{>1}^\kappa})$$

which implies that  $L_{\leq \ell}^\kappa$  is an  $\ell$ -level CL function over  $\kappa(V) = \kappa(V_1) \oplus \kappa(V_{>1})$ . This establishes Item 1 of Lemma 4.6, and completes the induction.  $\square$

**Lemma 4.12.** *Let  $V = \mathbb{F}_q^n$  for some integer  $n$  and prime power  $q = p^t$  for odd  $t$ . Let  $L, R : V \rightarrow V$  be CL functions. Let  $L^\kappa, R^\kappa : \kappa(V) \rightarrow \kappa(V)$  be the associated downsized CL functions, as defined in Definition 4.10. Then the distribution  $\mu_{L^\kappa, R^\kappa}$  over  $\kappa(V) \times \kappa(V)$  defined in Definition 4.5 is identical to the distribution of  $(x, y) \in \kappa(V) \times \kappa(V)$  obtained by first sampling  $(x', y')$  according to  $\mu_{L, R}$  and then returning  $(\kappa(x'), \kappa(y'))$ .*

*Proof.* The fact that  $L^\kappa, R^\kappa$  are well-defined CL functions follows from Lemma 4.11. The lemma is immediate from the definition of  $\mu_{L^\kappa, R^\kappa}$  and the fact that  $\kappa$  is a bijection.  $\square$

## 4.2 Conditionally linear samplers

Samplers are Turing machines that perform computations corresponding to CL functions defined in Section 4.1. The inputs and outputs of the sampler are binary strings that are interpreted as representing data of different types (integers, bits, vectors in  $\mathbb{F}_q^s$ , etc.). See Section 3.3.2 and in particular Remark 3.19 for an in-depth discussion of representing structured objects on a Turing machine.

**Definition 4.13.** A function  $q : \mathbb{N} \rightarrow \mathbb{N}$  is an *admissible field size function* if for all  $n \in \mathbb{N}$ ,  $q(n)$  is an admissible field size as defined in Definition 3.15.

**Definition 4.14** (Conditionally linear samplers). Let  $q : \mathbb{N} \rightarrow \mathbb{N}$  be an admissible field size function, and let  $s : \mathbb{N} \rightarrow \mathbb{N}$  be a function. A 6-input Turing machine  $\mathcal{S}$  is a  *$\ell$ -level conditionally linear sampler with field size  $q(n)$  and dimension  $s(n)$*  if for all  $n \in \mathbb{N}$ , letting  $q = q(n)$  and  $s = s(n)$ , there exist  $\ell$ -level CL functions  $L^{A, n}, L^{B, n} : \mathbb{F}_q^s \rightarrow \mathbb{F}_q^s$  with marginal functions  $\{L_{\leq j}^{w, n}\}$  and factor spaces  $\{V_{j, u}^{w, n}\}$  for  $w \in \{A, B\}$  satisfying the conditions of Lemma 4.6, such that for all  $w \in \{A, B\}, j \in \{1, \dots, \ell\}, z \in \mathbb{F}_q^s$ :

- On input  $(n, \text{DIMENSION})$ , the sampler  $\mathcal{S}$  outputs the dimension  $s(n)$ .
- On input  $(n, w, \text{MARGINAL}, j, z)$ , the sampler  $\mathcal{S}$  outputs the binary representation of  $L_{\leq j}^{w, n}(z)$ ,
- On input  $(n, w, \text{LINEAR}, j, u, y)$ , the sampler  $\mathcal{S}$  outputs the binary representation of  $L_{j, u}^{w, n}(y)$ , where  $u$  is interpreted as an element of  $V_{< j}^{w, n}$ ,
- On input  $(n, w, \text{FACTOR}, j, u)$ , the sampler  $\mathcal{S}$  outputs the  $j$ -th factor space  $V_{j, u}^{w, n}$  of  $L^{w, n}$  with prefix  $u \in L_{< j}^{w, n}(V)$ , represented as a vector in  $\{0, 1\}^s$  indicating which elementary basis vectors of  $\mathbb{F}_q^s$  span the factor space.

We call  $\mathbb{F}_q^{s(n)}$  the *ambient space* of  $\mathcal{S}$  on index  $n$ . We call the CL functions  $L^{w, n}$  for  $w \in \{A, B\}$  the *CL functions of  $\mathcal{S}$  on index  $n$* . The *time complexity* of  $\mathcal{S}$ , denoted as  $\text{TIME}_{\mathcal{S}}(n)$ , is the number of steps before  $\mathcal{S}$  halts for index  $n$ .

**Remark 4.15.** *Conditionally linear samplers are defined to have 6-input tapes, but depending on the input, not all input tapes are read. For example, if the second input tape has the input DIMENSION, then the remaining input tapes are ignored. Thus for notational convenience we write samplers with different numbers of arguments, depending on the type of argument it gets. The number of arguments is always at most 6, however.*

The following definition shows how samplers naturally correspond to conditionally linear distributions.

**Definition 4.16** (Distribution of a sampler). Let  $\mathcal{S}$  be a sampler with field size  $q(n)$ , dimension  $s(n)$ . For each  $n \in \mathbb{N}$ , let  $L^{A,n}, L^{B,n}$  denote the CL functions of  $\mathcal{S}$  on index  $n$ . Let  $\mu_{\mathcal{S},n}$  denote the CL distribution  $\mu_{L^{A,n}, L^{B,n}}$  corresponding to  $(L^{A,n}, L^{B,n})$ , as defined in Definition 4.5. We call  $\mu_{\mathcal{S},n}$  the *distribution of sampler  $\mathcal{S}$  on index  $n$* .

The following provides a definition of a “downsized” sampler that can be obtained from any sampler  $\mathcal{S}$  over an admissible field  $\mathbb{F}_q$ .

**Definition 4.17** (Downsized sampler). Let  $q : \mathbb{N} \rightarrow \mathbb{N}$  be an admissible field size function. Let  $\mathcal{S}$  be an  $\ell$ -level sampler with field size  $q(n)$  and dimension  $s(n)$ . Define  $\kappa(\mathcal{S})$  as the following Turing machine. For all  $n \in \mathbb{N}$ ,  $w \in \{A, B\}$ ,  $j \in \{1, \dots, \ell\}$ , and  $z \in \mathbb{F}_2^{s \log q}$  where  $q = q(n)$  and  $s = s(n)$ :

- On input  $(n, \text{DIMENSION})$ , the sampler returns the output of  $\mathcal{S}(n, \text{DIMENSION})$  multiplied by  $\log q$ .
- On input  $(n, w, \text{MARGINAL}, j, z)$ , the sampler  $\kappa(\mathcal{S})$  returns the output of  $\mathcal{S}(n, w, \text{MARGINAL}, j, z)$ .
- On input  $(n, w, \text{LINEAR}, j, u', y)$ , the sampler  $\kappa(\mathcal{S})$  computes  $u$  such that  $u' = \kappa(u)$  and returns the output of  $\mathcal{S}(n, w, \text{LINEAR}, j, u, y)$ .
- On input  $(n, w, \text{FACTOR}, j, u')$ , the sampler  $\kappa(\mathcal{S})$  computes  $u$  such that  $u' = \kappa(u)$  and the indicator vector

$$C = \mathcal{S}(n, w, \text{FACTOR}, j, u) \in \{0, 1\}^s,$$

and returns the expanded indicator vector  $(D_1, D_2, \dots, D_s) \in (\{0, 1\}^{\log q})^s$  where  $D_i$  is the all ones vector in  $\{0, 1\}^{\log q}$  if  $C_i = 1$  and  $D_i$  is the all zeroes vector otherwise.

The next lemma establishes that  $\kappa(\mathcal{S})$  is a well-defined CL sampler, in the sense that it can be derived from a family of CL functions as in Definition 4.14.

**Lemma 4.18.** *Let  $\ell \geq 1$  be such that  $\mathcal{S}$  is an  $\ell$ -level CL sampler, and let  $q(n)$  and  $s(n)$  be as in Definition 4.17. Then the Turing machine  $\kappa(\mathcal{S})$  is an  $\ell$ -level CL sampler with field size 2, dimension  $s'(n) = s(n) \log q(n)$ , and time complexity*

$$\text{TIME}_{\kappa(\mathcal{S})}(n) = O(\text{TIME}_{\mathcal{S}}(n) \log q(n)).$$

Furthermore, for every integer  $n \in \mathbb{N}$  the CL functions of  $\kappa(\mathcal{S})$  on index  $n$  are  $(L^{A,n})^\kappa$  and  $(L^{B,n})^\kappa$ , where  $L^{A,n}, L^{B,n}$  are the CL functions of  $\mathcal{S}$  on index  $n$ .

*Proof.* To show that  $\kappa(\mathcal{S})$  is an  $\ell$ -level CL sampler we first show the “Furthermore” part, i.e. verify that for any integer  $n \geq 1$  the CL functions  $(L^{A,n})^\kappa$  and  $(L^{B,n})^\kappa$  are its associated CL functions on index  $n$ , as defined in Definition 4.14.

Observe that for  $z \in V$ , the binary representation of  $z$  as an element of  $\{0, 1\}^{s \log q}$  passed as input to  $\mathcal{S}$  is, by definition (see Section 3.3.2), identical to the binary representation of  $\kappa(z)$ . Using the definition



$(L^{w,n})^\kappa = \kappa \circ L^{w,n} \circ \kappa^{-1}$  for  $w \in \{\mathbf{A}, \mathbf{B}\}$  this justifies that  $\kappa(\mathcal{S})$  returns the correct output when executed on inputs of the form  $(n, \text{DIMENSION})$ ,  $(n, w, \text{MARGINAL}, j, z)$  and  $(n, w, \text{LINEAR}, j, u', y)$ .

Next, if  $T$  is a register subspace of  $\mathbb{F}_q^s$  with indicator vector  $C \in \{0, 1\}^s$ , then  $\kappa(T)$  is a register subspace of  $\mathbb{F}_2^{s \log q}$  with indicator vector  $D$  defined from  $C$  as in Definition 4.17. Thus the output of  $\kappa(\mathcal{S})$  on input  $(n, w, \text{FACTOR}, j, u')$  is equal to the indicator vector of  $\kappa(V_{j,u}^{w,n})$ , which is the  $j$ -th factor space of  $L^{w,n}$  with prefix  $u' = \kappa(u)$ .

The time complexity of  $\kappa(\mathcal{S})$  is the same as with the sampler  $\mathcal{S}$ , except it takes  $O(\log q(n))$  times longer to output the factor space indicator vectors.

□

## 5 Nonlocal Games and MIP\*

We introduce definitions associated with nonlocal games and strategies that will be used throughout.

### 5.1 Games and strategies

**Definition 5.1** (Two-player one-round games). A *two-player one-round game*  $\mathfrak{G}$  is specified by a tuple  $(\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  where

1.  $\mathcal{X}$  and  $\mathcal{Y}$  are finite sets (called the *question alphabets*),
2.  $\mathcal{A}$  and  $\mathcal{B}$  are finite sets (called the *answer alphabets*),
3.  $\mu$  is a probability distribution over  $\mathcal{X} \times \mathcal{Y}$  (called the *question distribution*), and
4.  $D : \mathcal{X} \times \mathcal{Y} \times \mathcal{A} \times \mathcal{B} \rightarrow \{0, 1\}$  is a function (called the *decision predicate*).

**Definition 5.2** (Tensor product strategies). A *tensor product strategy*  $\mathcal{S}$  for a game  $\mathfrak{G} = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  is a tuple  $(|\psi\rangle, A, B)$  where

- $|\psi\rangle$  is a pure quantum state, i.e. a unit vector in  $\mathcal{H}_A \otimes \mathcal{H}_B$  for finite dimensional complex Hilbert spaces  $\mathcal{H}_A, \mathcal{H}_B$ ,
- $A$  is a set  $\{A^x\}$  such that for every  $x \in \mathcal{X}$ ,  $A^x = \{A_a^x\}_{a \in \mathcal{A}}$  is a POVM over  $\mathcal{H}_A$ , and
- $B$  is a set  $\{B^y\}$  such that for every  $y \in \mathcal{Y}$ ,  $B^y = \{B_b^y\}_{b \in \mathcal{B}}$  is a POVM over  $\mathcal{H}_B$ .

**Definition 5.3** (Tensor product value). The *tensor product value* of a tensor product strategy  $\mathcal{S} = (|\psi\rangle, A, B)$  with respect to a game  $\mathfrak{G} = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  is defined as

$$\text{val}^*(\mathfrak{G}, \mathcal{S}) = \sum_{x, y, a, b} \mu(x, y) D(x, y, a, b) \langle \psi | A_a^x \otimes B_b^y | \psi \rangle .$$

For  $v \in [0, 1]$  we say that the strategy  $\mathcal{S}$  *passes (or wins)*  $\mathfrak{G}$  with probability  $v$  if  $\text{val}^*(\mathfrak{G}, \mathcal{S}) \geq v$ . The *tensor product value* of  $\mathfrak{G}$  is defined as

$$\text{val}^*(\mathfrak{G}) = \sup_{\mathcal{S}} \text{val}^*(\mathfrak{G}, \mathcal{S}) ,$$

where the supremum is taken over all tensor product strategies  $\mathcal{S}$  for  $\mathfrak{G}$ .

**Remark 5.4.** Unless specified otherwise, all strategies considered in this paper are tensor product strategies, and we simply call them strategies. Similarly, we refer to  $\text{val}^*(\mathfrak{G})$  as the value of the game  $\mathfrak{G}$ .

**Definition 5.5** (Projective strategies). We say that a strategy  $\mathcal{S} = (|\psi\rangle, A, B)$  is *projective* if all the measurements  $\{A_a^x\}_a$  and  $\{B_b^y\}_b$  are projective.

**Definition 5.6.** A game  $\mathfrak{G} = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  is *symmetric* if the question and answer alphabets are the same for both players (i.e.  $\mathcal{X} = \mathcal{Y}$  and  $\mathcal{A} = \mathcal{B}$ ), the distribution  $\mu$  is symmetric (i.e.  $\mu(x, y) = \mu(y, x)$ ), and the decision predicate  $D$  treats both players symmetrically (i.e. for all  $x, y, a, b$ ,  $D(x, y, a, b) = D(y, x, b, a)$ ).

We call a strategy  $\mathcal{S} = (|\psi\rangle, A, B)$  *symmetric* if  $|\psi\rangle$  is a (pure) state in  $\mathcal{H} \otimes \mathcal{H}$ , for some Hilbert space  $\mathcal{H}$ , that is invariant under permutation of the two factors, and the measurement operators of both players are identical.

We often specify symmetric games  $\mathfrak{G}$  and symmetric strategies  $\mathcal{S}$  using a compact notation: we write  $\mathfrak{G} = (\mathcal{X}, \mathcal{A}, \mu, D)$  and  $\mathcal{S} = (|\psi\rangle, M)$  where  $M$  denotes the set of measurement operators for both players.

**Lemma 5.7.** *Let  $\mathfrak{G} = (\mathcal{X}, \mathcal{A}, \mu, D)$  be a symmetric game such that  $\text{val}^*(\mathfrak{G}) = 1 - \varepsilon$  for some  $\varepsilon \geq 0$ . Then there exists a symmetric and projective strategy  $\mathcal{S} = (|\psi\rangle, M)$  such that  $\text{val}^*(\mathfrak{G}, \mathcal{S}) \geq 1 - \varepsilon$ .*

*Proof.* By definition for any  $\varepsilon' > \varepsilon$  there exists a strategy  $\mathcal{S}' = (|\psi'\rangle, A, B)$  such that  $\text{val}^*(\mathfrak{G}, \mathcal{S}') \geq 1 - \varepsilon'$ . Using Naimark's theorem (Theorem 3.24) we can assume without loss of generality that  $|\psi'\rangle \in \mathbb{C}_{A'}^d \otimes \mathbb{C}_{B'}^d$ , for some integer  $d$  and that for every  $x$ ,  $A^x$  and  $B^x$  are projective measurements. Let

$$|\psi\rangle = \frac{1}{\sqrt{2}}(|0\rangle_A |1\rangle_B |\psi'\rangle_{A'B'} + |1\rangle_A |0\rangle_B |\psi'_\tau\rangle_{A'B'}) \in (\mathbb{C}_A^2 \otimes \mathbb{C}_{A'}^d) \otimes (\mathbb{C}_B^2 \otimes \mathbb{C}_{B'}^d),$$

where  $|\psi'_\tau\rangle$  is obtained from  $|\psi'\rangle$  by permuting the two players' registers. Observe that  $|\psi\rangle$  is invariant under permutation of  $AA'$  and  $BB'$ .

For any question  $x \in \mathcal{X}$  define the measurement  $M^x = \{M_a^x\}_{a \in \mathcal{A}}$  acting on the Hilbert space  $\mathbb{C}^2 \otimes \mathbb{C}^d$  as follows:

$$M_a^x = |0\rangle\langle 0| \otimes A_a^x + |1\rangle\langle 1| \otimes B_a^x,$$

i.e. for  $|\varphi\rangle \in \mathbb{C}^d$ ,  $M_a^x(|0\rangle|\varphi\rangle) = |0\rangle A_a^x |\varphi\rangle$  and  $M_a^x(|1\rangle|\varphi\rangle) = |1\rangle B_a^x |\varphi\rangle$ . When Alice receives question  $x$ , she measures  $M^x$  on registers  $AA'$ , and when Bob receives question  $y$ , he measures  $M^y$  on registers  $BB'$ . Using that by assumption the decision predicate  $D$  for  $\mathfrak{G}$  is symmetric, it is not hard to verify that  $\text{val}^*(\mathfrak{G}, \mathcal{S}) = \text{val}^*(\mathfrak{G}, \mathcal{S}')$ .  $\square$

**Definition 5.8.** Let  $\mathfrak{G} = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  be a game, and let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a strategy for  $\mathfrak{G}$  such that  $\mathcal{H}_A = \mathcal{H}_B$ . Let  $S \subseteq \mathcal{X} \times \mathcal{Y}$  denote the support of the question distribution  $\mu$ , i.e. the set of  $(x, y)$  such that  $\mu(x, y) > 0$ . We say that  $\mathcal{S}$  is a *commuting strategy* for  $\mathfrak{G}$  if for all question pairs  $(x, y) \in S$ , we have  $[A_a^x, B_b^y] = 0$  for all  $a \in \mathcal{A}, b \in \mathcal{B}$ , where  $[A, B] = AB - BA$  denotes the commutator.

**Definition 5.9** (Consistent measurements). Let  $\mathcal{A}$  be a finite set, let  $|\psi\rangle \in \mathcal{H} \otimes \mathcal{H}$  be a state, and let  $\{M_a\}_{a \in \mathcal{A}}$  be a projective measurement on  $\mathcal{H}$ . We say that  $\{M_a\}_{a \in \mathcal{A}}$  is *consistent on  $|\psi\rangle$*  if and only if

$$\forall a \in \mathcal{A}, \quad M_a \otimes I_B |\psi\rangle = I_A \otimes M_a |\psi\rangle.$$

When the state  $|\psi\rangle$  is clear from context we simply say that  $\{M_a\}_{a \in \mathcal{A}}$  is *consistent*.

The terminology ‘‘consistent’’ arises from the fact that when the measurement  $\{M_a\}$  is performed on both registers of  $|\psi\rangle$ , the probability of obtaining twice the same outcome is

$$\sum_a \langle \psi | M_a \otimes M_a | \psi \rangle = \sum_a \langle \psi | I_A \otimes (M_a)^2 | \psi \rangle = \sum_a \langle \psi | I_A \otimes M_a | \psi \rangle = 1,$$

where the first equality is by definition of consistency, the second uses that  $\{M_a\}$  is projective, and the last uses  $\sum_a M_a = I$ .

**Definition 5.10** (Consistent strategies). Let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a projective strategy with state  $|\psi\rangle \in \mathcal{H} \otimes \mathcal{H}$ , for some Hilbert space  $\mathcal{H}$ , which is defined on question alphabets  $\mathcal{X}$  and  $\mathcal{Y}$  and answer alphabets  $\mathcal{A}$  and  $\mathcal{B}$ , respectively. We say that the strategy  $\mathcal{S}$  is *consistent* if for all  $x \in \mathcal{X}$ , the measurement  $\{A_a^x\}_{a \in \mathcal{A}}$  is consistent on  $|\psi\rangle$  and if for all  $y \in \mathcal{Y}$ , the measurement  $\{B_b^y\}_{b \in \mathcal{B}}$  is consistent on  $|\psi\rangle$ .

We almost always restrict our attention to symmetric strategies. In this case, consistency implies that whenever both players are sent the same question, they provide the same answer.

**Definition 5.11.** We say that a strategy  $\mathcal{S}$  for a game  $\mathfrak{G}$  is *PCC* if it is projective, consistent, and commuting for  $\mathfrak{G}$ . Additionally, we say that a PCC strategy  $\mathcal{S}$  is *SPCC* if it is furthermore symmetric.

Recall that for a (pure) state  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ , the *Schmidt rank* of  $|\psi\rangle$  is the smallest integer  $k$  such that

$$|\psi\rangle = \sum_{i=1}^k \alpha_i |u_i\rangle \otimes |v_i\rangle,$$

for some orthonormal families  $\{|u_i\rangle\} \subset \mathcal{H}_A$  and  $\{|v_i\rangle\} \subset \mathcal{H}_B$  and  $\alpha_i > 0$ . The coefficients  $\alpha_i$  are uniquely defined by  $|\psi\rangle$  and called *Schmidt coefficients*. The state  $|\psi\rangle$  is called *maximally entangled* if  $\alpha_i = \frac{1}{\sqrt{k}}$  for all  $i = 1, \dots, k$ .

**Definition 5.12** (Entanglement requirements of a game). For all games  $\mathfrak{G}$  and  $\nu \in [0, 1]$ , let  $\mathcal{E}(\mathfrak{G}, \nu)$  denote the minimum integer  $d$  such that there exists a finite dimensional tensor product strategy  $\mathcal{S}$  that achieves success probability at least  $\nu$  in the game  $\mathfrak{G}$  with a state  $|\psi\rangle$  whose Schmidt rank is at most  $d$ . If there is no finite dimensional strategy that achieves success probability  $\nu$ , then define  $\mathcal{E}(\mathfrak{G}, \nu)$  to be  $\infty$ .

**Remark 5.13.** A strategy  $\mathcal{S} = (|\psi\rangle, A, B)$  for a symmetric game  $\mathfrak{G} = (\mathcal{X}, \mathcal{X}, \mathcal{A}, \mathcal{A}, \mu, D)$  is called *synchronous* if it holds that for every  $x \in \mathcal{X}$  and  $a \neq b \in \mathcal{A}$ ,  $\langle \psi | A_a^x \otimes B_b^x | \psi \rangle = 0$ ; in other words, the players never return different answers when simultaneously asked the same question. As shown in [PSS<sup>+</sup>16] the condition for a finite-dimensional strategy of being synchronous is equivalent to the condition that it is projective, consistent, and moreover  $|\psi\rangle$  is a maximally entangled state. (The equivalence is extended to infinite-dimensional strategies, as well as correlations induced by limits of finite-dimensional strategies, in [KPS18].)

## 5.2 Distance measures

We introduce several distance measures that are used throughout.

**Definition 5.14** (Distance between states). Let  $\{|\psi_n\rangle\}_{n \in \mathbb{N}}$  and  $\{|\psi'_n\rangle\}_{n \in \mathbb{N}}$  be two families of states in the same space  $\mathcal{H}$ . For some function  $\delta : \mathbb{N} \rightarrow [0, 1]$  we say that  $\{|\psi_n\rangle\}$  and  $\{|\psi'_n\rangle\}$  are  $\delta$ -close, denoted as  $|\psi\rangle \approx_\delta |\psi'\rangle$ , if  $\| |\psi_n\rangle - |\psi'_n\rangle \|^2 = O(\delta(n))$ . (For convenience we generally leave the dependence of the states and  $\delta$  on the indexing parameter  $n$  implicit, writing e.g.  $|\psi\rangle$  for  $|\psi_n\rangle$ .)

**Definition 5.15** (Consistency between POVMs). Let  $\mathcal{X}$  be a finite set and  $\mu$  a distribution on  $\mathcal{X}$ . Let  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  be a quantum state, and for all  $x \in \mathcal{X}$ ,  $\{A_a^x\}$  and  $\{B_a^x\}$  POVMs. We write

$$A_a^x \otimes I_B \simeq_\delta I_A \otimes B_a^x$$

on state  $|\psi\rangle$  and distribution  $\mu$  if

$$\mathbb{E}_{x \sim \mu} \sum_{a \neq b} \langle \psi | A_a^x \otimes B_b^x | \psi \rangle = O(\delta).^{24}$$

In this case, we say that  $\{A_a^x\}$  and  $\{B_a^x\}$  are  $\delta$ -consistent on  $|\psi\rangle$ .

<sup>24</sup>Here the use of the  $O(\cdot)$  notation refers to the fact that the notation will often be used to measure consistency for *families* of POVM and states indexed by an implicit parameter  $n \in \mathbb{N}$ , which will always be clear from context but omitted for legibility, see Definition 5.14. The  $O(\cdot)$  is taken as  $n \rightarrow \infty$ .

Note that a consistent measurement on  $|\psi\rangle$  according to Definition 5.9 is 0-consistent with itself on the same  $|\psi\rangle$  and under the singleton distribution, according to Definition 5.15 (and vice-versa). This is because

$$\sum_a \langle \psi | M_a \otimes M_a | \psi \rangle = 1 \implies \forall a, \langle \psi | I \otimes M_a | \psi \rangle = M_a \otimes \langle \psi | \psi \rangle,$$

because using that  $\{M_a\}$  is a POVM there must be equality in the Cauchy-Schwarz inequality

$$1 = \sum_a \langle \psi | M_a \otimes M_a | \psi \rangle \leq \left( \sum_a \|I \otimes M_a | \psi \rangle\|^2 \right) \left( \sum_a \|M_a \otimes I | \psi \rangle\|^2 \right)^{1/2} \leq 1.$$

**Definition 5.16** (Distance between POVMs). Let  $\mathcal{X}$  be a finite set and  $\mu$  a distribution on  $\mathcal{X}$ . Let  $|\psi\rangle \in \mathcal{H}$  be a quantum state, and for all  $x \in \mathcal{X}$ ,  $\{M_a^x\}$  and  $\{N_a^x\}$  two POVMs on  $\mathcal{H}$ . We say that  $\{M_a^x\}$  and  $\{N_a^x\}$  are  $\delta$ -close on state  $|\psi\rangle$  and under distribution  $\mu$  if

$$\mathbb{E}_{x \sim \mu} \sum_a \|(M_a^x - N_a^x) | \psi \rangle\|^2 \leq O(\delta),$$

and we write  $M_a^x \approx_\delta N_a^x$  to denote this when the state  $|\psi\rangle$  and distribution  $\mu$  are clear from context. This distance is referred to as the *state-dependent distance*.

**Definition 5.17** (Distance between strategies). Let  $\mathfrak{G} = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  be a nonlocal game and let  $\mathcal{S} = (\psi, A, B)$ ,  $\mathcal{S}' = (\psi', A', B')$  be strategies for  $\mathfrak{G}$ . For  $\delta \in [0, 1]$  we say that  $\mathcal{S}$  is  $\delta$ -close to  $\mathcal{S}'$  if the following conditions hold.

1. The states  $|\psi\rangle, |\psi'\rangle$  are states in the same Hilbert space  $\mathcal{H}_A \otimes \mathcal{H}_B$  and are  $\delta$ -close.
2. For all  $x \in \mathcal{X}, y \in \mathcal{Y}$ , we have  $A_a^x \approx_\delta (A'_a)^x$  and  $B_b^y \approx_\delta (B'_b)^y$ , with the approximations holding under the distribution  $\mu$ , and on either  $|\psi\rangle$  or  $|\psi'\rangle$ .

We record several useful facts about the consistency measure and the state-dependent distance without proof. Readers are referred to Sections 4.4 and 4.5 in [NW19] for additional discussion and proofs.

**Fact 5.18** (Fact 4.13 and Fact 4.14 in [NW19]). For POVMs  $\{A_a^x\}$  and  $\{B_a^x\}$ , the following hold.

1. If  $A_a^x \otimes I_B \simeq_\delta I_A \otimes B_a^x$  then  $A_a^x \otimes I_B \approx_\delta I_A \otimes B_a^x$ .
2. If  $A_a^x \otimes I_B \approx_\delta I_A \otimes B_a^x$  and  $\{A_a^x\}$  and  $\{B_a^x\}$  are projective measurements, then  $A_a^x \otimes I_B \simeq_\delta I_A \otimes B_a^x$ .
3. If  $A_a^x \otimes I_B \approx_\delta I_A \otimes B_a^x$  and either  $\{A_a^x\}$  or  $\{B_a^x\}$  is a projective measurement, then  $A_a^x \otimes I_B \simeq_{\delta^{1/2}} I_A \otimes B_a^x$ .

**Fact 5.19** (Fact 4.20 in [NW19]). Let  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  be finite sets, and let  $\mu$  be a distribution over question pairs  $(x, y)$ . Let  $\{A_{a,b}^x\}$  and  $\{B_{a,b}^x\}$  be operators whose outcomes range over the product set  $\mathcal{A} \times \mathcal{B}$ . Suppose a set of operators  $\{C_{a,c}^y\}$ , whose outcomes range over the product set  $\mathcal{A} \times \mathcal{C}$ , satisfies the condition  $\sum_c (C_{a,c}^y)^\dagger C_{a,c}^y \leq I$  for all  $a$  and all  $y$ . If  $A_{a,b}^x \approx_\delta B_{a,b}^x$  on average over  $x$  sampled from the corresponding marginal of distribution  $\mu$ , then  $C_{a,c}^y A_{a,b}^x \approx_\delta C_{a,c}^y B_{a,b}^x$  on average over  $(x, y)$  sampled from  $\mu$ .

*Proof.* Fix questions  $x, y$  and answers  $a \in \mathcal{A}, b \in \mathcal{B}$ . We have then that

$$\sum_c \left\| (C_{a,c}^y A_{a,b}^x - C_{a,c}^y B_{a,b}^x) |\psi\rangle \right\|^2 = \sum_c \langle \psi | (A_{a,b}^x - B_{a,b}^x)^\dagger (C_{a,c}^y)^\dagger (C_{a,c}^y) (A_{a,b}^x - B_{a,b}^x) | \psi \rangle \quad (33)$$

$$\leq \langle \psi | (A_{a,b}^x - B_{a,b}^x)^\dagger (A_{a,b}^x - B_{a,b}^x) | \psi \rangle \quad (34)$$

$$= \left\| (A_{a,b}^x - B_{a,b}^x) |\psi\rangle \right\|^2 \quad (35)$$

where the inequality follows from the assumption that  $\sum_c (C_{a,c}^y)^\dagger C_{a,c}^y \leq I$ . Thus we obtain the desired conclusion

$$\mathbb{E}_{(x,y) \sim \mu} \sum_{a,b,c} \left\| (C_{a,c}^y A_{a,b}^x - C_{a,c}^y B_{a,b}^x) |\psi\rangle \right\|^2 \leq \mathbb{E}_{(x,y) \sim \mu} \sum_{a,b} \left\| (A_{a,b}^x - B_{a,b}^x) |\psi\rangle \right\|^2 \leq \delta.$$

□

**Fact 5.20.** Let  $\mathcal{X}, \mathcal{A}$  denote finite sets, and let  $\mathcal{G}$  denote a set of functions  $g : \mathcal{X} \rightarrow \mathcal{A}$ . Let  $\{A_a^x\}, \{B_a^x\}$  be POVMs indexed by  $\mathcal{X}$  and outcomes in  $\mathcal{A}$ . Let  $\{S_g^x\}_{x \in \mathcal{X}, g \in \mathcal{G}}$  denote a set of operators such that for all  $x \in \mathcal{X}$ ,  $\sum_g (S_g^x)^\dagger S_g^x \leq I$ . If  $A_a^x \approx_\delta B_a^x$  on average over  $x$ , then  $S_g^x A_{g(x)}^x \approx_\delta S_g^x B_{g(x)}^x$ .

*Proof.* We expand:

$$\begin{aligned} \mathbb{E}_x \sum_g \left\| S_g^x (A_{g(x)}^x - B_{g(x)}^x) |\psi\rangle \right\|^2 &= \mathbb{E}_x \sum_g \langle \psi | (A_{g(x)}^x - B_{g(x)}^x)^\dagger (S_g^x)^\dagger S_g^x (A_{g(x)}^x - B_{g(x)}^x) | \psi \rangle \\ &= \mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x)^\dagger \left( \sum_{g: g(x)=a} (S_g^x)^\dagger S_g^x \right) (A_a^x - B_a^x) | \psi \rangle \\ &\leq \mathbb{E}_x \sum_a \langle \psi | (A_a^x - B_a^x)^\dagger (A_a^x - B_a^x) | \psi \rangle \\ &= \mathbb{E}_x \sum_a \left\| (A_a^x - B_a^x) |\psi\rangle \right\|^2. \end{aligned}$$

The inequality follows from the fact that  $\sum_{g: g(x)=a} (S_g^x)^\dagger S_g^x \leq \sum_g (S_g^x)^\dagger (S_g^x) \leq I$ . The last line is at most  $\delta$  by assumption, and we obtain the desired conclusion. □

**Lemma 5.21.** Let  $\mathcal{X}$  and  $\mathcal{A}$  be finite sets and  $\mu$  a distribution on  $\mathcal{X}$ . Let  $\{A_a^x\}_{a \in \mathcal{A}}$  be a projective measurement and let  $\{B_a^x\}_{a \in \mathcal{A}}$  be a set of matrices. If  $A_a^x \approx_\delta B_a^x$ , then for all subsets  $S \subseteq \mathcal{A}$ , we have

$$\sum_{a \in S} A_a^x \approx_\delta \sum_{a \in S} A_a^x \cdot B_a^x.$$

*Proof.* We expand:

$$\begin{aligned} \mathbb{E}_x \left\| \sum_{a \in S} (A_a^x - A_a^x \cdot B_a^x) |\psi\rangle \right\|^2 &= \mathbb{E}_x \left\| \sum_{a \in S} A_a^x \cdot (A_a^x - B_a^x) |\psi\rangle \right\|^2 \\ &= \mathbb{E}_x \sum_{a \in S} \langle \psi | (A_a^x - B_a^x)^\dagger A_a^x (A_a^x - B_a^x) | \psi \rangle \\ &\leq \mathbb{E}_x \sum_a \left\| (A_a^x - B_a^x) |\psi\rangle \right\|^2, \end{aligned}$$

where all expectations are taken according to  $\mu$ . In the second line we used the projectivity of  $\{A_a^x\}$ , and in the third line we used that  $A_a^x \leq I$ . □

**Fact 5.22** (Triangle inequality, Fact 4.28 in [NW19]). *If  $A_a^x \approx_\delta B_a^x$  and  $B_a^x \approx_\epsilon C_a^x$ , then  $A_a^x \approx_{\delta+\epsilon} C_a^x$ .*

**Fact 5.23** (Triangle inequality for “ $\simeq$ ”, Proposition 4.29 in [JNV<sup>+</sup>20]). *If  $A_a^x \otimes I_B \simeq_\epsilon I_A \otimes B_a^x$ ,  $C_a^x \otimes I_B \simeq_\delta I_A \otimes B_a^x$ , and  $C_a^x \otimes I_B \simeq_\gamma I_A \otimes D_a^x$ , then  $A_a^x \otimes I_B \simeq_{\epsilon+2\sqrt{\delta+\gamma}} I_A \otimes D_a^x$ .*

**Fact 5.24** (Data processing, Fact 4.26 in [NW19]). *Suppose  $A_a^x \otimes I_B \simeq_\delta I_A \otimes B_a^x$ . Then  $A_{[f(\cdot)=b]}^x \otimes I_B \simeq_\delta I_A \otimes B_{[f(\cdot)=b]}^x$ .*

The state-dependent distance is the right tool for reasoning about the closeness of measurement operators in a strategy. The following lemma ensures that, when two families of measurements are close on a state, changing from one family of measurement to the other only introduces a small error to the value of the strategy.

**Lemma 5.25.** *Let  $\{A_{a,b}^x\}$ ,  $\{B_{a,b,c}^x\}$ ,  $\{C_{a,c}^x\}$  be POVMs. Suppose  $\{B_{a,b,c}^x\}$  is projective, and*

$$\begin{aligned} A_{a,b}^x \otimes I_B &\approx_\delta I_A \otimes B_{a,b}^x, \\ C_{a,c}^x \otimes I_B &\approx_\delta I_A \otimes B_{a,c}^x, \end{aligned}$$

where recall that  $B_{a,b}^x = \sum_c B_{a,b,c}^x$  and similarly  $B_{a,c}^x = \sum_b B_{a,b,c}^x$ . Then the following approximate commutation relation holds:

$$[A_{a,b}^x, C_{a,c}^x] \otimes I_B \approx_\delta 0.$$

*Proof.* Applying Fact 5.19 to  $C_{a,c}^x \otimes I_B \approx_\delta I_A \otimes B_{a,c}^x$  and  $\{A_{a,b}^x \otimes I_B\}$ , we have

$$A_{a,b}^x C_{a,c}^x \otimes I_B \approx_\delta A_{a,b}^x \otimes B_{a,c}^x. \quad (36)$$

Similarly, applying Fact 5.19 to  $A_{a,b}^x \otimes I_B \approx_\delta I_A \otimes B_{a,b}^x$  and  $\{I_A \otimes B_{a,c}^x\}$ , and using the fact that  $\{B_{a,b,c}^x\}$  is projective, we have

$$\begin{aligned} A_{a,b}^x \otimes B_{a,c}^x &\approx_\delta I_A \otimes B_{a,c}^x B_{a,b}^x \\ &= I_A \otimes B_{a,b,c}^x. \end{aligned} \quad (37)$$

Combining Equations (36) and (37), we have

$$A_{a,b}^x C_{a,c}^x \otimes I_B \approx_\delta I_A \otimes B_{a,b,c}^x. \quad (38)$$

A similar argument gives

$$C_{a,c}^x A_{a,b}^x \otimes I_B \approx_\delta I_A \otimes B_{a,b,c}^x. \quad (39)$$

The claim follows from Equations (38) and (39).  $\square$

The following lemma is a slightly modified version of [NW19, Fact 4.34].

**Lemma 5.26.** *Let  $k \geq 0$  be an integer and let  $\epsilon > 0$ . Let  $\mathcal{X}$  be a finite set and  $\mu$  a distribution over  $\mathcal{X}$ . For each  $1 \leq i \leq k$  let  $\mathcal{G}_i$  be a finite set of functions  $g_i : \mathcal{Y} \rightarrow \mathcal{R}_i$  and for each  $x \in \mathcal{X}$  let  $\{G_g^{i,x}\}_{g \in \mathcal{G}_i}$  be a projective measurement. Suppose that for all  $i \in \{1, \dots, k\}$ ,  $\mathcal{G}_i$  satisfies the following property: for any two  $g_i \neq g'_i \in \mathcal{G}_i$ , the probability that  $g_i(y) = g'_i(y)$  over a uniformly random  $y \in \mathcal{Y}$  is at most  $\epsilon$ .*

*Let  $\{A_{g_1, g_2, \dots, g_k}^x\}$  be a projective measurement with outcomes  $(g_1, \dots, g_k) \in \mathcal{G}_1 \times \dots \times \mathcal{G}_k$ . For each  $1 \leq i \leq k$ , suppose that on average over  $x \sim \mu$  and  $y \in \mathcal{Y}$  sampled uniformly at random,*

$$A_{[\text{eval}_y(\cdot)=a_i]}^x \otimes I_B \simeq_\delta I_A \otimes G_{[\text{eval}_y(\cdot)=a_i]}^{i,x}. \quad (40)$$

Define the POVM family  $\{C_{g_1, g_2, \dots, g_k}^x\}$ , for  $x \in \mathcal{X}$ , by

$$C_{g_1, g_2, \dots, g_k}^x = G_{g_1}^{k, x} \dots G_{g_{k-1}}^{2, x} G_{g_k}^{1, x} G_{g_{k-1}}^{2, x} \dots G_{g_1}^{k, x}.$$

Then on average over  $x \sim \mu$  and  $y \in \mathcal{Y}$  sampled uniformly at random,

$$A_{[\text{eval}_y(\cdot)=(a_1, a_2, \dots, a_k)]}^x \otimes I_B \simeq_{k(\delta+\varepsilon)^{1/2}} I_A \otimes C_{[\text{eval}_y(\cdot)=(a_1, a_2, \dots, a_k)]}^x. \quad (41)$$

*Proof.* The proof is identical to the one given in [NW19, Fact 4.34], with the only modification needed to insert the dependence on  $x$  for all measurements considered.  $\square$

**Lemma 5.27** (Fact 4.35 in [NW19]). *Let  $\mathcal{D}$  be a distribution on  $(x, y_1, y_2) \in \mathcal{X} \times \mathcal{Y}_1 \times \mathcal{Y}_2$ . For  $i \in \{1, 2\}$  let  $\mathcal{G}_i$  be a collection of functions  $g_i : \mathcal{Y}_i \rightarrow \mathcal{R}_i$  and let  $\{(G_i)_g^x\}_{g \in \mathcal{G}_i}$  be families of measurements such that  $\{(G_2)_g^x\}_g$  is projective for every  $x$ . Suppose further that for every  $(x, y_1)$  it holds that for  $g_2 \neq g'_2 \in \mathcal{G}_2$  the probability, on average over  $y_2$  chosen from  $\mathcal{D}$  conditioned on  $(x, y_1)$ , that  $g_2(y_2) = g'_2(y_2)$  is at most  $\eta$ . Let  $\{A_{a_1, a_2}^{x, y_1, y_2}\}$  be a family of projective measurements with outcomes  $(a_1, a_2) \in \mathcal{R}_1 \times \mathcal{R}_2$  such that for  $i \in \{1, 2\}$ ,*

$$A_{a_i}^{x, y_1, y_2} \otimes I \simeq_\delta I \otimes (G_i)_{[\text{eval}_{y_i}(\cdot)=a_i]}^x \quad (42)$$

and

$$A_{a_1, a_2}^{x, y_1, y_2} \otimes I \simeq_\delta I \otimes A_{a_1, a_2}^{x, y_1, y_2}. \quad (43)$$

Define a family of measurements  $\{J_{g_1, g_2}^x\}$  as

$$J_{g_1, g_2}^x = (G_2)_{g_2}^x (G_1)_{g_1}^x (G_2)_{g_2}^x. \quad (44)$$

Then there is a

$$\delta_{\text{pasting}} = \delta_{\text{pasting}}(\eta, \delta) = \text{poly}(\eta, \delta) \quad (45)$$

such that

$$A_{a_1, a_2}^{x, y_1, y_2} \otimes I \simeq_{\delta_{\text{pasting}}} I \otimes J_{[\text{eval}_{y_1}(\cdot)=a_1, \text{eval}_{y_2}(\cdot)=a_2]}^x. \quad (46)$$

**Lemma 5.28** (Fact 4.32 of [NW19]). *Let  $\mathfrak{G}$  be a nonlocal game, and let  $\mathcal{S}, \mathcal{S}'$  be two strategies that are  $\delta$ -close (in the sense of Definition 5.17) for  $\delta \in [0, 1]$ , and use the same state  $|\psi\rangle$ . If either  $\mathcal{S}$  or  $\mathcal{S}'$  is projective, then  $|\text{val}^*(\mathfrak{G}, \mathcal{S}) - \text{val}^*(\mathfrak{G}, \mathcal{S}')| \leq O(\delta^{1/2})$ .*

### 5.3 The class MIP\*

The complexity class MIP\* of *multi-prover interactive proof systems with entangled provers* is the class of languages that can be decided by a *proof system* in which a polynomial-time classical *verifier* interacts with noncommunicating, computationally unbounded *provers* who may share a *finite-dimensional* entangled state. In general there may be polynomially many provers, and the verifier may interact with them over polynomially many rounds. In each round, the verifier generates a question for each prover and receives a response from them. At the end of the interaction the verifier decides to either accept or reject. A language  $L$  is in MIP\* if for every input  $z \in L$ , there exists a strategy that the provers can use to convince the verifier to accept with high probability ( $\geq 2/3$ ), and for every input  $z \notin L$ , there is *no* strategy that the provers can use to convince the verifier to accept with more than low probability ( $\geq 1/3$ ). The former property is called *completeness* of the proof system, and the latter *soundness*. The completeness and soundness probabilities  $2/3$  and  $1/3$  may be amplified by sequentially repeating the protocol.



For a formal definition of the class  $\text{MIP}^*$ , see e.g. [VW16, Section 6.1]. The main result of this paper is a lower bound, and for this purpose, it suffices to restrict our attention to proof systems which involve only two provers, one round of interaction, and completeness probability 1. This class is often denoted  $\text{MIP}_{1,1/2}^*(2,1)$ . A formal definition for this restricted setting follows.

**Definition 5.29.** A language  $L$  is in  $\text{MIP}_{1,1/2}^*$  if and only if there exist two randomized Turing machines  $\mathcal{S}$  and  $\mathcal{D}$  with the following properties.

1. **Efficiency:** For every  $z \in \{0,1\}^*$  there is a game  $\mathfrak{G}_z = (\mathcal{X}, \mathcal{Y}, \mathcal{A}, \mathcal{B}, \mu, D)$  such that:
  - (a) The Turing machine  $\mathcal{S}$  given input  $z$  runs in time  $\text{poly}(|z|)$  and returns a pair  $(x, y) \in \mathcal{X} \times \mathcal{Y}$  such that the distribution of  $(x, y)$ , over the random choices of  $\mathcal{S}$ , is  $\mu$ .
  - (b) The Turing machine  $\mathcal{D}$  given as input  $z$  and a tuple  $(x, y, a, b) \in \mathcal{X} \times \mathcal{Y} \times \mathcal{A} \times \mathcal{B}$  runs in time  $\text{poly}(|z|)$  and returns  $D(x, y, a, b)$ .<sup>25</sup>
2. **Completeness:** If  $z \in L$ , then  $\text{val}^*(\mathfrak{G}_z) = 1$
3. **Soundness:** If  $z \notin L$ , then  $\text{val}^*(\mathfrak{G}_z) \leq 1/2$ .

We say that the pair  $(\mathcal{S}, \mathcal{D})$  form an  $\text{MIP}^*$  *verifier* for the language  $L$ , and the associated family of games  $\mathfrak{G}_z$  are an  $\text{MIP}^*$  protocol for  $L$ .

It is clear that  $\text{MIP}_{1,1/2}^*(2,1) \subseteq \text{MIP}^*$  (the probability 1/2 of acceptance of inputs not in the language may be reduced to 1/3 by sequential repetition as noted above). The main result of this paper is that  $\text{RE} \subseteq \text{MIP}_{1,1/2}^*(2,1)$ . Since it is known that  $\text{MIP}^* \subseteq \text{RE}$  it follows that  $\text{MIP}_{1,1/2}^*(2,1) = \text{MIP}^* = \text{RE}$ .

To show our lower bound it will suffice to consider an even more restricted class of  $\text{MIP}^*$  verifiers, for which the Turing machines  $\mathcal{S}$  and  $\mathcal{D}$  have a special structure which we refer to as *normal form*. This is defined in the following subsection.

## 5.4 Normal form verifiers

We introduce a normal form for verifiers in nonlocal games. The normal form uses Turing machines to specify the two actions performed by the verifier in a game: the generation of questions and the verification of answers. For the generation of questions, we use the formalism of samplers introduced in Section 4.2. The normal form for verifiers gives a uniform method to specify an infinite family of nonlocal games.

**Definition 5.30 (Decider).** A *decider* is a 5-input Turing machine  $\mathcal{D}$  that on all inputs of the form  $(n, x, y, a, b)$  where  $n$  is an integer and  $x, y, a, b \in \{0,1\}^*$ ,  $\mathcal{D}$  halts and returns a single bit. Let  $\text{TIME}_{\mathcal{D}}(n)$  denote the maximum time complexity of  $\mathcal{D}$  over all inputs of the form  $(n, x, y, a, b)$ . When the decider  $\mathcal{D}$  outputs 0 we say that it *rejects*, otherwise we say that it *accepts*. Furthermore, we call the input  $n$  to a decider the *index*.

**Definition 5.31.** A *normal form verifier* is a pair  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  where  $\mathcal{S}$  is a sampler with field size  $q(n) = 2$  and  $\mathcal{D}$  is a decider. The description length of  $\mathcal{V}$  is defined to be  $|\mathcal{V}| = \max\{|\mathcal{S}|, |\mathcal{D}|\}$ , the maximum of the description lengths of  $\mathcal{S}$  and  $\mathcal{D}$ . The number of levels of verifier  $\mathcal{V}$  is defined to be the number of levels of its sampler  $\mathcal{S}$ .

---

<sup>25</sup>Note that the running time of  $\mathcal{D}$  should be  $\text{poly}(|z|)$ , even for long inputs  $a, b$ . This can be ensured by having  $\mathcal{D}$  return 0 whenever  $x, y, a, b$  are too long with respect to  $|z|$ .

Next we introduce a notion of *bounded normal form verifiers*, in which a single parameter (in this case, an integer  $\lambda$ ) specifies a bound on the time complexity of  $\mathcal{V}$ , as well as on the description length of the verifier.

**Definition 5.32** ( $\lambda$ -bounded verifiers). Let  $\lambda \in \mathbb{N}$  be an integer. A normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is  $\lambda$ -bounded if the following two conditions hold

1. The time complexity bounds of the verifier  $\text{TIME}_{\mathcal{S}}(n)$ ,  $\text{TIME}_{\mathcal{D}}(n)$  are at most  $n^\lambda$  for  $n \geq 2$ .<sup>26</sup>
2. The description length of the verifier  $|\mathcal{V}|$  is bounded by  $\lambda$ .

Normal form verifiers specify an infinite family of nonlocal games indexed by natural numbers in the following way.

**Definition 5.33.** Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier. For  $n \in \mathbb{N}$ , we define the following nonlocal game  $\mathcal{V}_n$  to be the  $n$ -th game corresponding to the verifier  $\mathcal{V}$ . The question sets  $\mathcal{X}$  and  $\mathcal{Y}$  are  $\{0, 1\}^{\text{TIME}_{\mathcal{S}}(n)}$ . The answer sets  $\mathcal{A}$  and  $\mathcal{B}$  are  $\{0, 1\}^{\text{TIME}_{\mathcal{D}}(n)}$ . The question distribution is the distribution  $\mu_{\mathcal{S}, n}$  specified in Definition 4.16. The decision predicate is the function computed by  $\mathcal{D}(n, \cdot, \cdot, \cdot, \cdot)$ , when the last four inputs are restricted to  $\mathcal{X} \times \mathcal{Y} \times \mathcal{A} \times \mathcal{B}$ . The value of the game is denoted by  $\text{val}^*(\mathcal{V}_n)$ .

We note that the game  $\mathcal{V}_n$  is well-defined since for a normal form verifier the distribution  $\mu_{\mathcal{S}, n}$  is supported on  $\{0, 1\}^{\text{TIME}_{\mathcal{S}}(n)} \times \{0, 1\}^{\text{TIME}_{\mathcal{S}}(n)}$  and a normal form decider always halts with a single-bit output.

**Definition 5.34** (Verifier with commuting strategy). Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier. For  $v : \mathbb{N} \rightarrow [0, 1]$  say that  $\mathcal{V}$  has a *value- $v$  commuting strategy* if for all  $n \in \mathbb{N}$ , the game  $\mathcal{V}_n$  has a value- $v(n)$  commuting strategy.

---

<sup>26</sup>We do not require the bound to hold for  $n = 1$  as  $n^\lambda$  is always 1 and hence it is usually not satisfied.

## 6 Types

We augment the definition of conditionally linear functions with a construct we call *types*. A type  $t$  is an element of a *type set*  $\mathcal{T}$ , and a  $\mathcal{T}$ -typed family of conditionally linear functions is a collection  $\{L_t\}_{t \in \mathcal{T}}$  containing a CL function  $L_t$  for each type  $t \in \mathcal{T}$ . The utility of this definition is that it allows us to define another object, namely conditionally linear distributions parameterized by an undirected graph  $G = (\mathcal{T}, E)$  on the set of types known as a *type graph*. Given two  $\mathcal{T}$ -typed families of conditionally linear functions  $\{L_u\}_{u \in \mathcal{T}}, \{R_v\}_{v \in \mathcal{T}}$ , the  $(\mathcal{T}, G)$ -typed conditionally linear distribution corresponding to them is the distribution which samples a pair of types  $(u, v)$  uniformly at random from the edges of  $G$  (with each endpoint having equal probability as being chosen for  $u$  or  $v$ , respectively) and then samples  $(x, y)$  from  $\mu_{L_u, R_v}$ . The output is the pair  $((u, x), (v, y))$ .

The normal form verifiers we present in the paper frequently use typed CL distributions to sample their questions, rather than untyped CL distributions. Types allow us to model the parts of their question distributions which are unstructured and unsuitable for being sampled from CL distributions. A common use of types is to allow the verifier to use previously defined games as subroutines. Here, the type helps indicate which subroutine the verifier selects, and an edge in the type graph between two different types allows us to introduce a test that cross-checks the results of one subroutine with the results of another.

Finally, we show how to convert any typed CL distribution into an equivalent (in the precise sense defined below) untyped CL distribution with two additional levels, a technique we call *detying*. This entails showing how to “simulate” the *graph distribution* of  $G = (\mathcal{T}, E)$ , i.e. the uniform distribution on its edges, using an untyped CL distribution. The simulation we give is based on rejection sampling and is only approximate: its quality degrades exponentially with the number of types in  $\mathcal{T}$ . As a result, we will ensure throughout the paper that all type sets we consider are of a small, in fact generally constant, size.

This section is organized as follows. In Section 6.1 we define typed variants of CL distributions, samplers, deciders, and verifiers. In Section 6.2 we define a CL distribution which samples from the graph distribution of a given graph  $G = (\mathcal{T}, E)$ . In Section 6.3 we define a canonical way to detype typed samplers, deciders, and verifiers using the graph sampler from Section 6.2. We then prove the main result of the section, Lemma 6.18, which relates the value of the detyped normal form verifier to the value of the original typed verifier.

### 6.1 Typed samplers, deciders, and verifiers

**Definition 6.1** (Typed conditionally linear functions). Let  $\mathcal{T}$  be a finite set and  $V$  be  $\mathbb{F}^n$  for some integer  $n \geq 0$ . A  $\mathcal{T}$ -typed family of  $\ell$ -level conditionally linear functions (implicitly, on  $V$ ) is a collection  $\{L_t\}_{t \in \mathcal{T}}$  such that, for each  $t \in \mathcal{T}$ ,  $L_t$  is an  $\ell$ -level conditionally linear function on  $V$ .

**Definition 6.2** (Graph distribution). Let  $G = (U, E)$  be an undirected graph with vertex set  $U$  and edge set  $E$ . Edges in  $E$  are written as multisets  $\{u, v\}$  of two vertices; the case  $u = v$  represents a self-loop. Suppose there are  $m$  edges,  $k$  of which are self-loops. Then the *graph distribution*  $\mu_G$  of  $G$  is the distribution over  $U \times U$  such that for every  $(u, v) \in U \times U$ ,

$$\mu_G(u, v) = \begin{cases} 1/(2m - k) & \text{if } \{u, v\} \in E, \\ 0 & \text{otherwise.} \end{cases}$$

This is identical to the uniform distribution over pairs  $(u, v) \in U \times U$  such that  $\{u, v\} \in E$ .

**Definition 6.3** (Typed conditionally linear distributions). Let  $\mathcal{T}$  be a type set and  $L = \{L_u\}_{u \in \mathcal{T}}, R = \{R_v\}_{v \in \mathcal{T}}$  be  $\mathcal{T}$ -typed families of conditionally linear functions on  $V$ . Let  $G = (\mathcal{T}, E)$  be a graph with vertex set  $\mathcal{T}$ . The  $(\mathcal{T}, G)$ -typed conditionally linear distribution  $\mu_{L,R}^G$  corresponding to  $(L, R)$  is the distribution over pairs  $((u, x), (v, y))$ , where  $(u, v)$  is drawn from  $\mu_G$  and  $(x, y)$  is drawn from  $\mu_{L_u, R_v}$ .

**Definition 6.4** (Typed conditionally linear samplers). Let  $q : \mathbb{N} \rightarrow \mathbb{N}$  be an admissible field size function and  $s : \mathbb{N} \rightarrow \mathbb{N}$  be a function. Let  $\mathcal{T}$  be a finite type set and let  $G = (\mathcal{T}, E)$  be a graph with vertex set  $\mathcal{T}$ . A 7-input Turing machine  $\mathcal{S}$  is a  $(\mathcal{T}, G)$ -typed,  $\ell$ -level conditionally linear sampler with field size  $q(n)$  and dimension  $s(n)$  if for all  $n \in \mathbb{N}$ , letting  $q = q(n)$  and  $s = s(n)$ , there exist  $\mathcal{T}$ -typed families of  $\ell$ -level conditionally linear functions  $\{L_t^{A,n}\}_{t \in \mathcal{T}}$  and  $\{L_t^{B,n}\}_{t \in \mathcal{T}}$  on  $V = \mathbb{F}_q^s$  where  $t \in \mathcal{T}, w \in \{A, B\}$ , the conditionally linear function  $L_t^{w,n}$  has marginal functions  $\{L_{t, \leq j}^{w,n}\}$  and factor spaces  $\{V_{t,j,u}^{w,n}\}$  satisfying the conditions of Lemma 4.6, and for all  $t \in \mathcal{T}, w \in \{A, B\}, j \in \{1, \dots, \ell\}$ , and  $z \in V$ :

- On input  $(n, \text{DIMENSION})$ , the sampler  $\mathcal{S}$  returns the dimension  $s(n)$ .
- On input  $(n, w, \text{MARGINAL}, j, z, t)$ , the sampler  $\mathcal{S}$  returns the binary representation of  $L_{t, \leq j}^{w,n}(z)$ .
- On input  $(n, w, \text{LINEAR}, j, u, y, t)$ , the sampler  $\mathcal{S}$  outputs the binary representation of  $L_{t,j,u}^{w,n}(y)$ ,
- On input  $(n, w, \text{FACTOR}, j, u, t)$ , the sampler  $\mathcal{S}$  returns the factor space  $V_{t,j,u}^{w,n}$  of  $L_t^{w,n}$ , represented as an indicator vector in  $\{0, 1\}^s$ .

We call  $\mathbb{F}_{q(n)}^{s(n)}$  the ambient space of  $\mathcal{S}$ . We call  $\{L_t^{A,n}\}, \{L_t^{B,n}\}$  the CL functions of  $\mathcal{S}$  on index  $n$ . The time complexity of  $\mathcal{S}$ , denoted  $\text{TIME}_{\mathcal{S}}(n)$ , is the number of steps before  $\mathcal{S}$  halts for index  $n$ .

We assume that types  $t \in \mathcal{T}$  are represented using binary strings of length at most  $\lceil \log |\mathcal{T}| \rceil$ ; if a type  $t$  is given as input to the sampler  $\mathcal{S}$  and is not an element of  $\mathcal{T}$ , then the sampler returns 0. Furthermore, as described in Remark 4.15 for un-typed samplers, we write typed samplers with different numbers of arguments depending on the input. We note that Definition 6.4 includes the graph  $G$  as a parameter, even though it is not explicitly used anywhere; this is so we can define the following concept, i.e. the *distribution* of a typed sampler, which *does* depend on  $G$ .

**Definition 6.5** (Distribution of a typed sampler). Let  $\mathcal{S}$  be a  $(\mathcal{T}, G)$ -typed sampler. Let  $L^w = \{L_t^w\}_t$  for  $w \in \{A, B\}$  be the CL functions of  $\mathcal{S}$  on index  $n$ . The *distribution of sampler  $\mathcal{S}$  on index  $n$* , denoted  $\mu_{\mathcal{S},n}^G$ , is the  $(\mathcal{T}, G)$ -typed conditionally linear distribution corresponding to  $(L^A, L^B)$ .

**Definition 6.6** (Downsizing typed CL samplers). Let  $\mathcal{S}$  be a typed sampler. The typed downsized sampler  $\kappa(\mathcal{S})$  is defined as in Definition 4.17 with the only difference that the type  $t$  is included as part of the input to the sampler, as in Definition 6.4.

**Lemma 6.7.** *Let  $\mathcal{S}$  be a  $(\mathcal{T}, G)$ -typed  $\ell$ -level CL sampler, for some finite set  $\mathcal{T}$ , graph  $G$ , and integer  $\ell \geq 0$ . Let  $q(n)$  and  $s(n)$  be as in Definition 6.4. Then  $\kappa(\mathcal{S})$  defined in Definition 6.6 is a  $(\mathcal{T}, G)$ -typed  $\ell$ -level CL sampler with field size 2, dimension  $s(n) \log q(n)$ , and time complexity*

$$\text{TIME}_{\kappa(\mathcal{S})}(n) = O(\text{TIME}_{\mathcal{S}}(n) \log q(n)).$$

*Furthermore, for every integer  $n \geq 1$ , the CL functions of  $\kappa(\mathcal{S})$  on index  $n$  are  $\{(L_t^{w,n})^\kappa\}_{w \in \{A, B\}, t \in \mathcal{T}}$ , as defined in Definition 4.10.*

*Proof.* The proof is analogous to the proof of Lemma 4.18, and we omit it.  $\square$

**Definition 6.8** (Typed decider). A *typed decider* is a 7-input Turing machine  $\mathcal{D}$  that on all inputs of the form  $(n, u, x, v, y, a, b)$  where  $n$  is an integer and  $u, x, v, y, a, b \in \{0, 1\}^*$ ,  $\mathcal{D}$  halts and returns a single bit. When  $\mathcal{D}$  returns 0 we say that it *rejects*, otherwise we say that it *accepts*. We use  $\text{TIME}_{\mathcal{D}}(n)$  to denote the time complexity of  $\mathcal{D}$  on inputs of the form  $(n, \dots)$ .

**Definition 6.9.** Let  $\mathcal{T}$  be a finite set and let  $G = (\mathcal{T}, E)$  be a graph. A  $(\mathcal{T}, G)$ -typed normal form verifier is a pair  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  where  $\mathcal{S}$  is a  $(\mathcal{T}, G)$ -typed sampler with field size  $q(n) = 2$  and  $\mathcal{D}$  is a typed decider.

**Definition 6.10.** Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a  $(\mathcal{T}, G)$ -typed normal form verifier. For  $n \in \mathbb{N}$ , we define the following nonlocal game  $\mathcal{V}_n$  to be the  $n$ -th game corresponding to the verifier  $\mathcal{V}$ . The question sets  $\mathcal{X}$  and  $\mathcal{Y}$  are  $\mathcal{T} \times \{0, 1\}^{\text{TIME}_{\mathcal{S}}(n)}$ . The answer sets  $\mathcal{A}$  and  $\mathcal{B}$  are  $\{0, 1\}^{\text{TIME}_{\mathcal{D}}(n)}$ . The question distribution is the distribution  $\mu_{\mathcal{S}, n}^G$  specified in Definition 6.5. The decision predicate is the function computed by  $\mathcal{D}(n, u, x, v, y, a, b)$ , when the inputs  $((u, x), (v, y), a, b)$  are restricted to  $\mathcal{X} \times \mathcal{Y} \times \mathcal{A} \times \mathcal{B}$ . The value of the game is denoted by  $\text{val}^*(\mathcal{V}_n)$ .

For  $w \in \{A, B\}$  and a question  $(u, x)$  to player  $w$  we refer to  $u$  as the *question type* and  $x$  as the *question content*.

## 6.2 Graph distributions

We describe a construction of conditionally linear distributions which sample from the graph distribution (see Definition 6.2) of a graph  $G = (U, E)$ . We begin with a technical definition, followed by the definition of the conditionally linear distribution.

**Definition 6.11** (Neighbor indicator). Given a graph  $G = (U, E)$ , the *neighbor indicator* of a vertex  $u \in U$  is the vector  $\text{neigh}_G(u) \in \mathbb{F}_2^U$  in which, for all  $v \in U$ ,

$$\text{neigh}_G(u)_v = \begin{cases} 1 & \text{if } \{u, v\} \in E, \\ 0 & \text{otherwise.} \end{cases}$$

In addition, the  $\mathbb{F}_2$ -encoding of a vertex  $u \in U$  is the vector  $\text{enc}_G(u) \in \mathbb{F}_2^U \times \mathbb{F}_2^U$  given by  $\text{enc}_G(u) = (e_u, \text{neigh}_G(u))$ , where  $e_u$  is the standard basis vector with a 1 in the  $u$ -th position and 0's everywhere else.

**Definition 6.12** (Graph sampler). Let  $G = (U, E)$  be a graph with  $n$  vertices. Then the *conditionally linear functions corresponding to  $G$*  are the pair of functions  $L_G^A, L_G^B$  on linear space  $V_G$  specified in Fig. 2 where  $V_G = V_{VA} \oplus V_{NA} \oplus V_{VB} \oplus V_{NB}$ .

These conditionally linear functions do not simulate the graph distribution in the sense of sampling directly from it. The following proposition, however, does show a sense in which these functions simulate the graph distribution, namely via rejection sampling.

**Proposition 6.13** (Simulating the graph distribution). *Let  $G = (U, E)$  be a graph with  $n$  vertices and  $m$  edges,  $k$  of which are self-loops. Let  $L_G^A, L_G^B$  be the conditionally linear functions corresponding to  $G$  (see Figure 2 for the definition and associated notation). Let  $(x, y) \sim \mu_{L_G^A, L_G^B}$ . Consider the event  $\mathcal{E}_G$  that there exists  $u, v \in U$  such that the following two statements are true.*

(i)  $x^{V_{VA} \oplus V_{NA}} = \text{enc}_G(u)$  and  $y^{V_{VB} \oplus V_{NB}} = \text{enc}_G(v)$ ,

<b>Subspaces</b>			
$V_{VA}$	$V_{NA}$	$V_{VB}$	$V_{NB}$
$\mathbb{F}_2^U$	$\mathbb{F}_2^U$	$\mathbb{F}_2^U$	$\mathbb{F}_2^U$
<b>Conditionally linear function <math>L_G^A</math></b>			
1st factor subspace	$V_{VA} \oplus V_{NA}$		
1st linear function	Identity function		
2nd factor subspace	$V_{VB} \oplus V_{NB}$		
2nd linear functions	For all $x \in V_{VA} \oplus V_{NA}$ , suppose there exists a $u \in U$ such that $x = \text{enc}_G(u)$ . Then for all $y \in V_{VB} \oplus V_{NB}$ , $L_{G,2,x}^A$ zeroes out all entries of $y$ except for $(y^{V_{NB}})_u$ . Otherwise, $L_{G,2,x}^A = 0$ .		
<b>Conditionally linear function <math>L_G^B</math></b>			
1st factor subspace	$V_{VB} \oplus V_{NB}$		
1st linear function	Identity function		
2nd factor subspace	$V_{VA} \oplus V_{NA}$		
2nd linear functions	Similarly defined as those for $L_G^A$ by swapping $V_{VA}$ and $V_{NA}$ with $V_{VB}$ and $V_{NB}$ respectively.		

Figure 2: Specification of the conditionally linear functions corresponding to  $G$ .

(ii)  $(x^{V_{NB}})_u = (y^{V_{NA}})_v = 1$ .

Then

1.  $\Pr_{x,y}(\mathcal{E}_G) = (2m - k)/16^n$ .
2. Conditioned on  $\mathcal{E}_G$ ,  $(u, v)$  are distributed as the graph distribution of  $G$  (see Definition 6.2).

Note that  $\mathcal{E}_G$  occurs if and only if both  $x^{V_{NB}}$  and  $y^{V_{NA}}$  are nonzero. In particular, if  $x$  and  $y$  are sampled from  $\mu_{L_G^A, L_G^B}$  and given to the respective players, then at least one of them knows when the event  $\mathcal{E}_G$  does not occur.

*Proof.* Let  $z$  be drawn uniformly at random from  $V_{VA} \oplus V_{NA} \oplus V_{VB} \oplus V_{NB}$ , and let  $x = L_G^A(z)$  and  $y = L_G^B(z)$ . Then with probability  $n^2/16^n$ , there exist  $u, v \in U$  such that

$$x^{V_{VA} \oplus V_{NA}} = \text{enc}_G(u) \text{ and } y^{V_{VB} \oplus V_{NB}} = \text{enc}_G(v) .$$

Conditioned on this occurring,  $u$  and  $v$  are distributed as independent, uniformly random vertices in  $U$ . If we further condition on  $\{u, v\} \in E$ , which occurs with probability  $(2m - k)/n^2$ , then by definition,  $(u, v)$  is distributed as the graph distribution of  $G$ . But this event is exactly the event that  $\mathcal{E}_G$  holds on  $(x, y)$ , establishing the proposition.  $\square$

### 6.3 Detying typed verifiers

We give a canonical method for taking a typed normal form verifier and producing an untyped normal form verifier which simulates it. Throughout this section,  $\mathcal{T}$  denotes a finite set,  $G = (\mathcal{T}, E)$  denotes a graph, and  $L_G^A, L_G^B$  denote the conditionally linear functions corresponding to  $G$  acting on the vector space  $V_G = V_{VA} \oplus V_{NA} \oplus V_{VB} \oplus V_{NB}$  of dimension  $4 \cdot |\mathcal{T}|$  over  $\mathbb{F}_2$ , as in Definition 6.12.

**Definition 6.14** (Detyed CL functions). Let  $L^A = \{L_t^A\}, L^B = \{L_t^B\}$  be  $\mathcal{T}$ -typed families of  $\ell$ -level conditionally linear functions on  $V$ . We define the *detyed CL functions corresponding to  $(L^A, L^B)$  on  $G$*  to be the pair of  $(\ell + 2)$ -level CL functions  $(R^A, R^B) = \text{Detype}_G(L^A, L^B)$  on linear space  $V_{\text{DETYPE}} = V_G \oplus V$  as follows. For  $w \in \{A, B\}$ , and  $z \in L_G^w(V_G)$ , define the family of  $\ell$ -level CL functions  $\{L_z^w\}$  on  $V$  as

$$L_z^w = \begin{cases} 0 & \text{if } z^{V_{N\bar{w}}} = 0, \\ L_t^w & \text{otherwise, for } z^{V_{Vw}} = e_t. \end{cases}$$

We note that when  $z^{V_{N\bar{w}}}$  is nonzero, it is always the case that  $z^{V_{Vw}} = e_t$  for a unique type  $t$ , by Definition 6.12. For  $w \in \{A, B\}$ ,  $R^w$  is the concatenation of  $L_G^w$  and  $\{L_z^w\}_z$  (cf. Lemma 4.7).

**Definition 6.15** (Detyed samplers). Let  $\mathcal{S}$  be a  $(\mathcal{T}, G)$ -typed sampler. For each  $n \in \mathbb{N}$ , let  $\{L_t^{A,n}\}, \{L_t^{B,n}\}$  be the CL functions of  $\mathcal{S}$  on index  $n$ , and set  $(R^{A,n}, R^{B,n}) = \text{Detype}_G(L^{A,n}, L^{B,n})$ . Then the *detyed sampler*  $\text{Detype}(\mathcal{S})$  is the (standard) sampler whose CL functions on index  $n$  are  $R^{A,n}, R^{B,n}$ . Its dimension function is  $s_{\text{DETYPE}}(n) = 4|\mathcal{T}| + s(n)$ .

We note that  $\text{Detype}(\mathcal{S})$  is indeed a sampler in line with Definition 4.14, i.e. a sampler without types.

**Definition 6.16** (Detyed deciders). Let  $\mathcal{D}$  be a typed decider. We define the *detyed decider*  $\text{Detype}_G(\mathcal{D})$  to be the (standard) decider that behaves as follows: on input  $(n, x, y, a, b)$ , it attempts to parse  $x = (x', x'')$ ,  $y = (y', y'') \in V_G \times \{0, 1\}^*$  (using a canonical scheme for representing pairs of strings). If it cannot, it accepts. Otherwise, suppose that there exists  $\{u, v\} \in E_G$  such that, using notation from Definition 6.11,

$$x' = (e_u, \text{neigh}_G(u), 0, e_u), \quad y' = (0, e_v, e_v, \text{neigh}_G(v)) \in V_{VA} \oplus V_{NA} \oplus V_{VB} \oplus V_{NB}.$$

Then it returns the output of  $\mathcal{D}$  on input  $(n, u, x'', v, y'', a, b)$ . Otherwise, it accepts.

We note that we include the subscript “ $G$ ” in  $\text{Detype}_G(\mathcal{D})$  because we do not specify a graph when defining a typed decider (cf. Definition 6.8). This is different than the situation for samplers  $\mathcal{S}$ , where we do not include a subscript in  $\text{Detype}(\mathcal{S})$  because the graph  $G$  is already specified when defining  $\mathcal{S}$ .

**Definition 6.17** (Detyed verifiers). Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a  $(\mathcal{T}, G)$ -typed normal form verifier. We define the *detyed verifier*, denoted by  $\text{Detype}(\mathcal{V})$ , to be the (standard) normal form verifier  $(\text{Detype}(\mathcal{S}), \text{Detype}_G(\mathcal{D}))$ .

**Lemma 6.18** (Typed verifiers to detyed verifiers). *Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a  $(\mathcal{T}, G)$ -typed normal form verifier. The detyed verifier  $\text{Detype}(\mathcal{V}) = (\text{Detype}(\mathcal{S}), \text{Detype}_G(\mathcal{D}))$  satisfies the following properties: for all  $n \in \mathbb{N}$ ,*

1. (**Completeness**) *If  $\mathcal{V}_n$  has a value-1 PCC strategy, then  $\text{Detype}(\mathcal{V})_n$  has a value-1 PCC strategy.*
2. (**Soundness**) *If  $\text{val}^*(\text{Detype}(\mathcal{V})_n) \geq 1 - \varepsilon$ , then  $\text{val}^*(\mathcal{V}_n) \geq 1 - 16^{|\mathcal{T}|} \cdot \varepsilon$ . Furthermore,*

$$\mathcal{E}(\text{Detype}(\mathcal{V})_n, 1 - \varepsilon) \geq \mathcal{E}(\mathcal{V}_n, 1 - 16^{|\mathcal{T}|} \cdot \varepsilon).$$

3. (**Sampler parameters**) If  $\mathcal{S}$  is an  $\ell$ -level sampler, then  $\text{Detype}(\mathcal{S})$  is an  $(\ell + 2)$ -level sampler. The time complexity of  $\text{Detype}(\mathcal{S})$  satisfies the following:

$$\text{TIME}_{\text{Detype}(\mathcal{S})}(n) = \text{poly}(|\mathcal{T}|, \text{TIME}_{\mathcal{S}}(n)).$$

4. (**Decider complexity**) The decider  $\text{Detype}_G(\mathcal{D})$  has time complexity  $\text{poly}(|\mathcal{T}|, \text{TIME}_{\mathcal{D}}(n))$ .
5. (**Efficient computability**) The descriptions of  $\text{Detype}(\mathcal{S})$  and  $\text{Detype}_G(\mathcal{D})$  are polynomial time computable from the description of  $G$  and the descriptions of  $\mathcal{S}$  and  $\mathcal{D}$ , respectively.

*Proof.* Throughout this proof, we fix an index  $n$ . Let  $s = s(n)$  be the dimension of  $\mathcal{S}$ . The ambient space of  $\mathcal{S}$  is  $V = \mathbb{F}_2^s$  and the ambient space of  $\text{Detype}(\mathcal{S})$  is  $V_{\text{DETYPE}} = V_G \oplus V$ . Let  $u, v \in \mathcal{T}$ . For this proof, we introduce the notation

$$\text{view}^A(u) = (e_u, \text{neigh}_G(u), 0, e_u), \text{view}^B(v) = (0, e_v, e_v, \text{neigh}_G(v)) \in V_{VA} \oplus V_{NA} \oplus V_{VB} \oplus V_{NB}.$$

Supposing that players A and B receive  $x$  and  $y$  in  $V_{\text{DETYPE}}$ , and supposing that  $(x, y)$  satisfies event  $\mathcal{E}_G$  from Proposition 6.13, then  $x^{V_G} = \text{view}^A(u)$  and  $y^{V_G} = \text{view}^B(v)$  for a unique  $\{u, v\} \in E$ .

**Completeness.** Let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a value-1 PCC strategy for  $\mathcal{V}_n$ . We construct a PCC strategy  $\mathcal{S}^{\text{DETYPE}}$  for  $\text{Detype}(\mathcal{V})_n$  with value 1. This strategy also uses the state  $|\psi\rangle$ . When a player receives a question, they perform measurements described as follows.

Player A: given  $x \in V_{\text{DETYPE}}$  the player checks if for some  $u \in \mathcal{T}$ ,  $x^{V_G} = \text{view}^A(u)$ . If so, they perform the measurement

$$\left\{ A_a^{(u, x^V)} \right\}$$

to obtain an outcome  $a$ , which they use as their answer. If not, they reply with the empty string. (This entails performing the measurement whose POVM element corresponding to the empty string is the identity matrix.)

Player B: given  $y \in V_{\text{DETYPE}}$ , the player checks if for some  $v \in \mathcal{T}$ ,  $y^{V_G} = \text{view}^B(v)$ . If so, they perform the measurement

$$\left\{ B_b^{(v, y^V)} \right\}$$

to obtain an outcome  $b$ , which they use as their answer. If not, they reply with the empty string.

This strategy is projective and consistent because the only measurements it uses are those in  $\mathcal{S}$  and “trivial” measurements containing the identity matrix. Suppose the players receive questions  $x$  and  $y$  such that both  $x^{V_G} = \text{view}^A(u)$  and  $y^{V_G} = \text{view}^B(v)$ . In this case, the questions  $(u, x^V)$  and  $(v, y^V)$  are in the support of the question distribution of  $\mathcal{V}$ . As a result, the players succeed with probability 1 on these questions, and their measurements always commute. For the remaining pairs of questions, the decider  $\text{Detype}_G(\mathcal{D})$  always accepts, and the measurements always commute by virtue of the fact that at least one is trivial, i.e. containing the identity matrix as a POVM element.



**Soundness.** Let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a strategy for  $\text{Detype}(\mathcal{V})_n$  with value  $1 - \varepsilon$ . Suppose  $G$  has  $m$  edges,  $k$  of which are self-loops. For any  $(x, y)$  drawn from  $\mu_{\text{Detype}(\mathcal{S}), n}$ , the decider  $\text{Detype}_G(\mathcal{D})$  automatically accepts unless  $(x^{V_G}, y^{V_G})$  satisfies event  $\mathcal{E}_G$  from Proposition 6.13, which occurs with probability  $(2m - k)/16^{|\mathcal{T}|}$ . When this happens,  $x^{V_G}$  and  $y^{V_G}$  are distributed as  $\text{view}^A(u)$  and  $\text{view}^B(v)$ , where  $(u, v)$  are distributed as the graph distribution on  $G$ . As a result, conditioned on  $\mathcal{E}_G$ , the probability that  $\mathcal{S}$  succeeds on  $\text{Detype}(\mathcal{V})_n$  is equal to the probability that the strategy  $\mathcal{S}' = (|\psi\rangle, A', B')$  succeeds on  $\mathcal{V}_n$ , where

$$(A')_a^{u, x^V} = A_a^{(\text{view}^A(u), x^V)}, \quad (B')_b^{v, y^V} = B_b^{(\text{view}^B(v), y^V)}.$$

This means that

$$\begin{aligned} \text{val}^*(\text{Detype}(\mathcal{V})_n, \mathcal{S}) &= \left(1 - \frac{2m - k}{16^{|\mathcal{T}|}}\right) + \frac{2m - k}{16^{|\mathcal{T}|}} \cdot \text{val}^*(\mathcal{V}_n, \mathcal{S}') \\ &\leq \left(1 - \frac{1}{16^{|\mathcal{T}|}}\right) + \frac{1}{16^{|\mathcal{T}|}} \cdot \text{val}^*(\mathcal{V}_n, \mathcal{S}'). \end{aligned}$$

Thus,  $\mathcal{S}'$  has value at least  $1 - 16^{|\mathcal{T}|} \cdot \varepsilon$ . This proves the first statement in the soundness. As for the second,  $\mathcal{S}$  and  $\mathcal{S}'$  use the same state  $|\psi\rangle$ , and therefore both strategies have the same Schmidt rank, which by definition is at least  $\mathcal{E}(\mathcal{V}_n, 1 - 16^{|\mathcal{T}|} \cdot \varepsilon)$ .

**Complexity.** Definition 6.14 implies that  $\text{Detype}(\mathcal{S})$  is an  $(\ell + 2)$ -level sampler by Lemma 4.7 and that  $V_{\text{DETYPE}}$  has dimension  $4|\mathcal{T}| + s$ . The claimed time bounds of  $\text{Detype}(\mathcal{S})$  and  $\text{Detype}_G(\mathcal{D})$  follow from the fact that these perform simple,  $\text{poly}(|\mathcal{T}|)$ -time computations followed by running  $\mathcal{S}$  and  $\mathcal{D}$  as subroutines.  $\square$

## 7 Classical and Quantum Low-degree Tests

In this section we introduce the classical and quantum low individual degree tests, that form the basis of our classical, quantum-sound and quantum PCP constructions. For quantum soundness of the classical test we refer to [JNV<sup>+</sup>20]. For the quantum test, we refer to [NV18a] and its adaptation to the low individual degree test given in Appendix A. The protocol in this work combines both of these tests, using the quantum low individual degree test for question reduction and the classical low individual degree test for answer reduction. Section 7.1 below introduces the classical test, and Section 7.3 does the same for the quantum test. Prior to doing this, we introduce the *Magic Square game* in Section 7.2, a key subroutine in the quantum test.

### 7.1 The classical low-degree test

We begin with a generalization of the classical low-degree test known as the “simultaneous individual low-degree test”. We sometimes refer to this as the “classical low-degree test” for short. The low-degree test is used as a subroutine in the Pauli Basis test (see Section 7.3) as well as the answer-reduction normal form verifier (see Section 10). We describe the test as a nonlocal game in Section 7.1.1 and show that the question distribution can be implemented as a (typed) conditionally linear distribution (see Section 6 for the definition of typed CL distributions).

#### 7.1.1 The game

The game  $\mathfrak{G}^{\text{LD}}$  is parametrized by a tuple  $\text{ldparams} = (q, m, d, k)$  where  $m, d, k \in \mathbb{N}$  are integers,  $q \in \mathbb{N}$  is an admissible field size, and  $m$  divides  $q$ . We sometimes write  $\mathfrak{G}_{\text{ldparams}}^{\text{LD}}$  to emphasize the dependence of the classical low-degree test on the parameter tuple  $\text{ldparams}$ .

We first provide a high-level description of the game for  $k = 1$ . It is based on the low-individual degree variant of the multilinearity test of [BFL91], where one player (called the “points player”) receives a random point  $u \in \mathbb{F}_q^m$ , and the other player (called the “lines player”) receives a random axis-parallel line  $\ell$  that contains  $u$ , i.e. the line parallel to the  $i$ -th axis which goes through point  $u$ , for uniformly random  $i \in \{1, \dots, m\}$ . The points player is supposed to respond with a value  $a \in \mathbb{F}_q$ , and the lines player is supposed to respond with a univariate polynomial  $f_\ell$  of degree at most  $d$  such that  $f_\ell(u_i) = a$ , where  $u_i$  is the  $i$ -th coordinate of  $u$ .

Suppose the players agree beforehand on a global polynomial  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  such that every variable has degree at most  $d$ . Then a winning strategy is the following: the points player responds with  $g(u)$ , and the lines player responds with the univariate polynomial  $f_\ell$  that is the restriction of  $g$  to line  $\ell$ ; formally, for  $t \in \mathbb{F}_q$ ,

$$f_\ell(t) = g(u_1, u_2, \dots, u_{i-1}, t, u_{i+1}, \dots, u_m).$$

It is easy to see that  $f_\ell(t)$  has degree at most  $d$  in the variable  $t$ . Thus in this case the players will pass the low-degree test with probability 1. The low-degree testing theorem states that the converse approximately holds: if the players pass the low-degree test with probability close to 1, then their responses must be approximately consistent (in some sense) with a global polynomial  $g$  with individual degree  $d$ .

When  $k > 1$ , the players are supposed to respond with a *tuple* of answers; and the test is intended to check that the players’ responses are consistent with a  $k$ -tuple of functions  $g_i : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  that are polynomials where every variable has degree at most  $d$ .

The game that we use is slightly more elaborate than the “axis-parallel lines-versus-points” test just described. In addition, we have two other subtests. One is a “diagonal line-vs-points” test: here, the points

player receives a point  $u \in \mathbb{F}_q^m$  as before, but the lines player receives a line  $\ell$  that goes through  $u$  but is not necessarily axis-aligned. Instead, it's chosen by first picking an index  $i \in \{1, 2, \dots, m\}$  uniformly at random, picking  $v \in \mathbb{F}_q^m$  uniformly at random, and letting  $v' = \pi_{i-1}(v) = (0, 0, \dots, 0, v_i, v_{i+1}, \dots, v_m)$  (i.e.,  $\pi_{i-1}$  zeroes out the first  $i - 1$  coordinates of its input). Then, the line is defined as the set

$$\ell = \{u + tv' : t \in \mathbb{F}_q\},$$

which is a uniformly random line in the affine subspace of  $\mathbb{F}_q^m$  corresponding to all points  $v$  whose first  $(i - 1)$  coordinates match those of  $u$ . The points player responds with a value  $a \in \mathbb{F}_q$ , the lines player responds with a univariate polynomial  $f_\ell$  of degree  $md$ ,<sup>27</sup> and the verifier checks that  $f_\ell(t_u) = a$  (where  $t_u \in \mathbb{F}_q$  is the projection of  $u$  onto the line  $\ell$  with respect to a canonical parameterization of  $\ell$ ).

The other additional test is a ‘‘consistency test’’, where both players receive either the same points question, the same axis-parallel lines question, or the same diagonal lines question. In all cases they are expected to respond with the same answer.

**Axis-parallel lines and diagonal lines.** Before presenting a formal description of the game  $\mathfrak{G}^{\text{LD}}$ , we first define what we mean by lines, and their properties.

**Definition 7.1 (Lines).** A line  $\ell$  in  $\mathbb{F}_q^m$  specified by a pair  $(u, v) \in (\mathbb{F}_q^m)^2$  is the subset

$$\ell(u, v) = \{u + tv : t \in \mathbb{F}_q\} \subseteq \mathbb{F}_q^m. \quad (47)$$

We call the vector  $v$  a *direction* of line  $\ell(u, v)$ .<sup>28</sup>

A line  $\ell(u, v)$  is *axis-parallel* if there is a coordinate  $i \in \{1, 2, \dots, m\}$  for which  $v$  is equal to  $e_i$ , the  $i$ -th elementary basis vector in  $\mathbb{F}_q^m$ . We say that such a line  $\ell$  is *parallel to the  $i$ -th direction*. A line  $\ell(u, v)$  is *diagonal* if there exists an  $i \in \{0, 1, \dots, m - 1\}$  for which  $v_1 = v_2 = \dots = v_i = 0$ .

The following Proposition establishes that, for a fixed direction, a line  $\ell$  is an equivalence class of points on the line.

**Proposition 7.2.** *Let  $u, v \in \mathbb{F}_q^m$ . Then for all  $u' \in \ell(u, v)$ , we have that  $\ell(u, v) = \ell(u', v)$ .*

*Proof.* Fix  $u' \in \ell(u, v)$ . We have that  $u' = u + tv$  for some  $t \in \mathbb{F}_q$ . Let  $a \in \ell(u, v)$ . This means that  $a = u + sv$  for some  $s \in \mathbb{F}_q$ . We also have that  $a = u + (s - t + t)v = u + tv + (s - t)v = u' + (s - t)v$ , so  $\ell(u, v) \subseteq \ell(u', v)$ . A similar argument shows that  $\ell(u', v) \subseteq \ell(u, v)$ .  $\square$

We now specify a canonical representation of lines  $\ell = \ell(u, v)$  as pairs  $(u_0, v)$  where  $u_0$  is a *canonical representative* of  $\ell$  specified as follows. Let  $L_v^{\text{LN}} : \mathbb{F}_q^m \rightarrow \mathbb{F}_q^m$  denote the canonical linear map with the one-dimensional kernel basis  $\{v\}$  (see Definition 3.10 for a definition of canonical linear map). Note that for all  $u \in \ell$ , we have that  $L_v^{\text{LN}}(u)$  is a point in  $\ell$ . This is because since the image and kernel of  $L_v^{\text{LN}}$  are complementary subspaces by definition (see Definition 3.10), any  $u \in \mathbb{F}_q^m$  has a unique decomposition  $u = u_0 + u_1$  where  $u_0$  is in the image of  $L_v^{\text{LN}}$  and  $u_1 \in \ker(L_v^{\text{LN}})$ , which by definition is the subspace spanned by  $v$ . Thus  $u_1 = tv$  for some  $t \in \mathbb{F}_q$  and we have that  $u_0 = u - tv \in \ell$ . In particular, if  $u \in \ell$  then  $u_0 = L_v^{\text{LN}}(u) \in \ell$  as well. Furthermore, since  $L_v^{\text{LN}}$  is a projection onto its image by definition, we have  $L_v^{\text{LN}}(u_0) = u_0$ .

Next, note that for any  $u, u' \in \ell$ , since  $u - u' = sv$  for some  $s \in \mathbb{F}_q$  it follows that  $L_v^{\text{LN}}(u) = L_v^{\text{LN}}(u')$ . Thus  $u_0 = u'_0$  and this shows that  $L_v^{\text{LN}}$  gives a means to compute a canonical representative of lines.

<sup>27</sup>Since  $\ell$  is no longer axis-parallel, the restriction of an individual degree- $d$  polynomial  $g$  to a line  $\ell$  may yield a degree  $md$  polynomial.

<sup>28</sup>For convenience we explicitly allow  $v = 0$ , in which case the line is reduced to a singleton.

**Definition 7.3** (Canonical representative of a line). Let  $\ell$  be a line in direction  $v$ . A canonical representative of  $\ell$  is the point  $u_0 \in \mathbb{F}_q^m$  such that

$$u_0 = L_v^{\text{LN}}(u)$$

for all  $u \in \ell$ .

**Question distribution.** We now present the question distribution of the game  $\mathfrak{G}^{\text{LD}}$  as a typed CL distribution (see Section 6 for the definition of typed CL functions and typed CL distributions). The type set  $\mathcal{T}^{\text{LD}}$  of the game  $\mathfrak{G}^{\text{LD}}$  we use is

$$\mathcal{T}^{\text{LD}} = \{\text{POINT}, \text{ALINE}, \text{DLINE}\}.$$

The distribution over question types is uniform over  $\mathcal{T}^{\text{LD}} \times \mathcal{T}^{\text{LD}}$ .

We introduce the corresponding typed CL functions  $L_{\text{POINT}}, L_{\text{ALINE}}, L_{\text{DLINE}}$  where the corresponding CL distributions  $\mu_{L_{\text{ALINE}}, L_{\text{POINT}}}$  and  $\mu_{L_{\text{DLINE}}, L_{\text{POINT}}}$  implement the axis line-versus-point and diagonal line-versus-point distributions, respectively. The functions are parametrized by a field size  $q$ , a dimension  $m$ , and three complementary register subspaces  $V_X, V_I, V_V$  of some ambient space. The register  $V_X$  (which is isomorphic to  $\mathbb{F}_q^m$ ) is called the *point register*,  $V_I$  (which is isomorphic to  $\mathbb{F}_q$ ) is called the *coordinate register*, and  $V_V$  (which is isomorphic to  $\mathbb{F}_q^m$ ) is called the *direction register*, respectively. We let  $V$  denote the direct sum  $V_X \oplus V_I \oplus V_V$ . We identify elements of  $V$  as triples  $(u, s, v) \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$ .

Define  $L_{\text{POINT}}$  to be the 1-level CL function that projects onto  $V_X$ : for every  $(u, s, v) \in V$ ,

$$L_{\text{POINT}}(u, s, v) = (u, 0, 0). \quad (48)$$

Define  $L_{\text{ALINE}}$  to be the following 2-level CL function:

$$L_{\text{ALINE}}(u, s, v) = (L_{e_i}^{\text{LN}}(u), s, 0) \text{ for all } (u, s, v) \in V, \quad (49)$$

where  $L^{\text{LN}}$  is the linear function used to compute the canonical representative of lines (see Definition 7.3), and  $i = \chi(s)$  where we define  $\chi(s)$  to be the unique integer such that

$$s = (\chi(s) - 1) \frac{q}{m} + r \quad (50)$$

for some  $0 \leq r < q/m$ , and we interpret the field element  $s \in \mathbb{F}_q$  as an integer between 0 and  $q - 1$ ). The function  $L_{\text{ALINE}}$  is the concatenation of the 1-level CL function that projects the subspaces  $V_I \oplus V_V$  down to  $V_I$  (i.e. zeroes out the entire  $V_V$  register), and the family of 1-level CL functions  $\{L_{e_i}^{\text{LN}}\}$  acting on  $V_X$ , indexed by  $i \in \{1, 2, \dots, m\}$ . (See Lemma 4.7 for definition of CL function concatenation.) Thus  $L_{\text{ALINE}}$  is a 2-level CL function.

Define  $L_{\text{DLINE}}$  to be the following 3-level CL function: for all  $(u, s, v) \in V$ ,

$$L_{\text{DLINE}}(u, s, v) = (L_{\pi_{i-1}(v)}^{\text{LN}}(u), s, \pi_{i-1}(v)) \quad (51)$$

where  $i = \chi(s)$ . The function  $L_{\text{DLINE}}$  is the concatenation of the following CL functions (again, see Lemma 4.7 for definition of CL function concatenation): (i) the identity function on  $V_I$  (which is a 1-level CL function); (ii) for every  $i \in \{1, 2, \dots, m\}$  the 1-level CL function on  $V_V$  that maps  $v \mapsto \pi_{i-1}(v)$  (i.e. it zeroes out the first  $i - 1$  coordinates of  $v$ ), and (iii) for every pair  $(s, v') \in V_I \oplus V_V$ , the 1-level CL function  $L_{v'}^{\text{LN}}$  acting on  $V_X$ . Thus  $L_{\text{DLINE}}$  is a 3-level CL function.

The following two lemmas establish that the CL functions  $L_{\text{POINT}}, L_{\text{ALINE}}, L_{\text{DLINE}}$  give rise to the axis-parallel line-versus-point and diagonal line-versus-point distributions.

**Lemma 7.4.** Consider the pair  $(\ell, u)$  sampled in the following way: first, sample the tuple  $((u_0, s, 0), (u, 0, 0))$  from the CL distribution  $\mu_{L_{\text{ALINE}}, L_{\text{POINT}}}$ , and let  $\ell = \ell(u_0, e_i)$  where  $i = \chi(s)$ . Then  $u$  is a uniformly random point in  $\mathbb{F}_q^m$  and  $\ell$  is a line that goes through  $u$  and is parallel to a uniformly random axis  $i \in \{1, 2, \dots, m\}$ .

*Proof.* Since  $s$  is uniformly random in  $\mathbb{F}_q$  and  $m$  divides  $q$  by assumption, we have that  $i$  is uniformly random between  $\{1, 2, \dots, m\}$  and therefore  $(u_0, s)$  specifies a uniformly random axis-parallel line  $\ell(u_0, e_i)$ . Since  $u_0 = L_{e_i}^{\text{LN}}(u)$ , we have that  $u \in \ell(u_0, e_i)$ .  $\square$

**Lemma 7.5.** Consider the pair  $(\ell, u)$  sampled in the following way: first, sample the tuple  $((u_0, s, v), (u, 0, 0))$  from the CL distribution  $\mu_{L_{\text{DLINE}}, L_{\text{POINT}}}$ , and let  $\ell = \ell(u_0, v)$  where  $i = \chi(s)$ . Then  $u$  is a uniformly random point in  $\mathbb{F}_q^m$  and  $\ell$  is a random diagonal line that goes through  $u$  and shares the first  $i - 1$  coordinates with  $u$ , for uniformly random  $i$ .

*Proof.* The proof of this follows identically to Lemma 7.4.  $\square$

For later convenience we make the following definition.

**Definition 7.6** (Line-point distributions). The axis line-point distribution  $D_{\text{ALINE}}$  is the distribution  $\mu_{L_{\text{ALINE}}, L_{\text{POINT}}}$  over pairs  $(\ell = (u_0, e_i), u)$  (we omit the 0 elements for simplicity). The diagonal line-point distribution  $D_{\text{DLINE}}$  is the distribution  $\mu_{L_{\text{DLINE}}, L_{\text{POINT}}}$  over  $(\ell = (u_0, s, v), u)$ . The line-point distribution  $D_{\text{LINE}}$  is the equal mixture of the axis-parallel line-point distribution and the diagonal line-point distribution. In other words, a sample from  $D_{\text{LINE}}$  is distributed as follows: with probability  $\frac{1}{2}$ , output a sample from  $D_{\text{ALINE}}$ , and with probability  $\frac{1}{2}$ , output a sample from  $D_{\text{DLINE}}$ .

**Decision procedure** The decision procedure  $\mathcal{D}^{\text{LD}}$  for the game  $\mathfrak{G}^{\text{LD}}$  is presented in Figure 3. The table at the top specifies a parsing scheme for the questions and answers, depending on the type of question. For example, when a player receives a question with type POINT, the question content  $x$ , a bit string of length  $m \log q$ , should be interpreted by the decision procedure and the players as an element of the vector space  $\mathbb{F}_q^m$ , as indicated in Section 3.3.2. Similarly the answer to a question with type POINT is expected to be a bit string of length  $k \log q$ , and is interpreted as an element of  $\mathbb{F}_q^k$ . For questions with type ALINE, the question content is a  $(m \log q + \log q)$ -bit string, which is parsed as a pair  $(u_0, s) \in \mathbb{F}_q^m \times \mathbb{F}_q$ , which in turn can be interpreted as a specification for an axis-parallel line  $\ell(u_0, e_i)$  in  $\mathbb{F}_q^m$  where  $i = \chi(s)$ , as described in Lemma 7.4. The answer is interpreted as the description of  $k$  degree- $d$  univariate polynomials defined on the line  $\ell(u_0, e_i)$ , each polynomial given by a list of  $d + 1$  coefficients. For questions with type DLINE, the question content is a  $(2m \log q + \log q)$ -bit string, which is parsed as a triple  $(u_0, s, v) \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$  that specifies a diagonal line  $\ell(u_0, v')$  where  $v' = \pi_{i-1}(v)$  as described in Lemma 7.5. The answer is interpreted as the description of  $k$  degree- $md$  univariate polynomials defined on the line  $\ell(u_0, v')$ . If the answers returned by the players do not fit this format the decision procedure rejects.

We define a special class of measurements that are relevant to the soundness properties of the low-degree test.

**Definition 7.7** (Low-degree polynomial measurements). Define  $\text{PolyMeas}(m, d, q)$  to be the set of POVM measurements whose outcomes correspond to (individual) degree- $d$  polynomials of  $m$  variables over  $\mathbb{F}_q$ . More generally, for an integer  $k$  and tuples  $m = (m_1, m_2, \dots, m_k)$ ,  $d = (d_1, d_2, \dots, d_k)$  and  $q = (q_1, q_2, \dots, q_k)$ , we let  $\text{PolyMeas}(m, d, q, k)$  be the set of measurements  $G = \{G_{g_1, g_2, \dots, g_k}\}$  such that for  $i \in \{1, 2, \dots, k\}$ ,  $g_i$  is a polynomial  $g_i : \mathbb{F}_{q_i}^{m_i} \rightarrow \mathbb{F}_{q_i}$  with individual degree  $d_i$  (see Section 3.4 for a definition of individual degree).

Type	Question Content	Answer Format
POINT	$u \in \mathbb{F}_q^m$	Element of $\mathbb{F}_q^k$
ALINE	$(u_0, s) \in \mathbb{F}_q^m \times \mathbb{F}_q$	$k$ degree- $d$ polynomials $f_j : \mathbb{F}_q \rightarrow \mathbb{F}_q$
DLINE	$(u_0, s, v) \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$	$k$ degree- $md$ polynomials $f_j : \mathbb{F}_q \rightarrow \mathbb{F}_q$

Input to  $\mathcal{D}^{\text{LD}}$ :  $(t_A, x_A, t_B, x_B, a_A, a_B)$ . In all cases where no action is indicated, accept. For  $w \in \{A, B\}$ ,

1. **(Consistency test)** If  $t_A = t_B$ , accept iff  $a_A = a_B$ .
2. **(Axis-parallel line-versus-point test)** If  $t_w = \text{ALINE}$  and  $t_{\bar{w}} = \text{POINT}$ , accept iff  $f_j(t) = (a_{\bar{w}})_j$  for all  $j \in \{1, 2, \dots, k\}$  where  $t \in \mathbb{F}_q$  is such that  $x_{\bar{w}} = u_0 + te_i$  where  $i = \chi(s)$ .
3. **(Diagonal line-versus-point test)** If  $t_w = \text{DLINE}$  and  $t_{\bar{w}} = \text{POINT}$ , accept iff  $f_j(t) = (a_{\bar{w}})_j$  for all  $j \in \{1, 2, \dots, k\}$  where  $t \in \mathbb{F}_q$  is such that  $x_{\bar{w}} = u_0 + tv'$ , where  $v' = \pi_{i-1}(v)$  with  $i = \chi(s)$ .

Figure 3: The decision procedure  $\mathcal{D}^{\text{LD}}$  for the simultaneous low-degree test, parameterized by  $\text{ldparams} = (q, m, d, k)$ . The function  $\chi(s)$  is defined in Equation (50), and  $\pi_{i-1}(v)$  zeroes out the first  $i - 1$  coordinates of  $v$ .

Quantum soundness of a version of the classical low-degree test for polynomials of low *total* degree was claimed in [NV18a], building on an analysis of a three-prover version of the test in [Vid16]. Unfortunately there is a gap in the soundness analysis of this test. For our construction we use the low individual degree test, whose quantum soundness is shown in [JNV<sup>+</sup>20] and generalized in [JNV<sup>+</sup>21]. We state the result in the form that is most directly useful for us, and show how the stated result follows from [JNV<sup>+</sup>21].

**Theorem 7.8** (Quantum soundness of the simultaneous classical low-degree test). *There exists a function  $\delta_{\text{LD}}(\varepsilon, q, m, d, k) = a(dm k)^a(\varepsilon^b + q^{-b} + 2^{-bmd})$  for universal constants  $a \geq 1$  and  $0 < b \leq 1$  such that the following holds. For all  $\varepsilon > 0$  and parameter tuple  $\text{ldparams} = (q, m, d, k)$ , for all projective strategies  $(\psi, A, B)$  that succeed with probability at least  $1 - \varepsilon$  in the game  $\mathfrak{G}_{\text{ldparams}}^{\text{LD}}$ , there exists measurements*

$$G^w \in \text{PolyMeas}(m, d, q, k)$$

on  $\mathcal{H}_w$ , for  $w \in \{A, B\}$ , such that

$$\begin{aligned} A_{a_1, a_2, \dots, a_k}^{\text{POINT}, u} \otimes I_B &\simeq_{\delta_{\text{LD}}} I_A \otimes G_{[(g_1(u), g_2(u), \dots, g_k(u)) = (a_1, a_2, \dots, a_k)]}^B, \\ G_{[(g_1(u), g_2(u), \dots, g_k(u)) = (a_1, a_2, \dots, a_k)]}^A \otimes I_B &\simeq_{\delta_{\text{LD}}} I_A \otimes B_{a_1, a_2, \dots, a_k}^{\text{POINT}, u}, \\ G_{g_1, g_2, \dots, g_k}^A \otimes I_B &\simeq_{\delta_{\text{LD}}} I_A \otimes G_{g_1, g_2, \dots, g_k}^B, \end{aligned}$$

where  $\delta_{\text{LD}} = \delta_{\text{LD}}(\varepsilon, q, m, d, k)$ , and all three statements are with respect to the state  $|\psi\rangle$ . In addition, the first two approximations hold under the uniform distribution over  $u \in \mathbb{F}_q^m$ , whereas there is no question distribution associated with the third approximation.

*Proof.* Let  $\text{ldparams} = (q, m, d, k)$  where  $m, d, k \in \mathbb{N}$  are integers,  $q \in \mathbb{N}$  is an admissible field size, and  $m$  divides  $q$ . Let  $\Sigma = \mathbb{F}_q$ . Let  $\mathcal{C} \subseteq \Sigma^q$  be the Reed-Solomon code with degree  $d$ . Explicitly,  $\mathcal{C}$  is the set of all  $(p(x_1), \dots, p(x_n))$  where  $p : \mathbb{F}_q \rightarrow \mathbb{F}_q$  is a univariate polynomial of degree at most  $d$  and  $x_1, \dots, x_n$  are an enumeration of  $\mathbb{F}_q$ . Then in the (standard) notation from [JNV<sup>+</sup>21]  $\mathcal{C}$  is a linear  $[q, d + 1, n - d + 1]_\Sigma$  code.

Observe that the two-prover tensor code test, for the code  $\mathcal{C}^{\otimes m}$ , as described in [JNV<sup>+</sup>21, Figure 1 and Figure 2], is identical to the game  $\mathcal{G}^{\text{LD}}$  with parameters  $\text{ldparams}$  and  $k = 1$ , whose decider is presented in Figure 3. This is because

1. The *axis-parallel line-versus-point test* from Figure 3 is identical to the *axis-parallel lines test* from [JNV<sup>+</sup>21, Figure 1], with the only difference that in Figure 3 the roles of A and B are chosen uniformly at random. This is consistent with the two-prover variant of the tensor code test from [JNV<sup>+</sup>21, Section 4.1].
2. The *diagonal line-versus-point test* from Figure 3 is identical to the *subcube commutation test* from [JNV<sup>+</sup>21, Figure 1], with again the added symmetrization between players.
3. The *consistency test* from Figure 3 is identical to the *synchronicity test* from [JNV<sup>+</sup>21, Figure 2].

As a result, [JNV<sup>+</sup>21, Theorem 4.7], in which we choose the parameter  $k$  as  $k = m^3 d$ , immediately yields Theorem 7.8.

To conclude it remains to extend the result to the case of general  $k$ . This is done via a standard reduction, following exactly the same steps as the derivation of Theorem 4.43 from Theorem 4.40 in [NW19].  $\square$

### 7.1.2 Complexity of the classical low-degree test.

The CL functions and decision procedure of the low-degree test are incorporated as subroutines in some of the normal form verifiers constructed in subsequent sections. The next lemma establishes the time complexity of these procedures as a function of the parameter tuple  $\text{ldparams} = (q, m, d, k)$ . The lemma also establishes the time complexity of computing the description of the decision procedure  $\mathcal{D}^{\text{LD}}$  as a Turing machine, given the parameter tuple  $\text{ldparams}$  as input.

**Lemma 7.9** (Complexity of the classical low-degree test). *Let  $\text{ldparams} = (q, m, d, k)$  denote a parameter tuple.*

1. *The time complexity of the decision procedure  $\mathcal{D}^{\text{LD}}$  parametrized by  $\text{ldparams}$  is  $\text{poly}(m, d, k, \log q)$ .*
2. *The time complexity of evaluating marginals of the CL functions  $L_{\text{POINT}}$ ,  $L_{\text{ALINE}}$ , and  $L_{\text{DLINE}}$  at a given input point is  $\text{poly}(m, \log q)$ .*
3. *The Turing machine description of the decision procedure  $\mathcal{D}^{\text{LD}}$  parametrized by  $\text{ldparams}$  can be computed from  $\text{ldparams}$  in  $\text{polylog}(q, m, d, k)$  time.*

*Proof.* Finite field arithmetic over  $\mathbb{F}_q$  can be performed in time  $\text{polylog } q$ , by Lemma 3.18. The most expensive step in  $\mathcal{D}^{\text{LD}}$  is to evaluate a univariate polynomial  $f : \mathbb{F}_q \rightarrow \mathbb{F}_q^k$  at some point  $t_u \in \mathbb{F}_q$  that corresponds to a point  $u \in \mathbb{F}_q^m$ , which takes time  $\text{poly}(m, d, k, \log q)$ . The function  $L_{\text{POINT}}$  is a projection onto  $V_X$ , which takes time  $\text{poly}(m, \log q)$  to compute (it just involves “zeroing out” the registers outside of  $V_X$ ). The functions  $L_{\text{ALINE}}$  and  $L_{\text{DLINE}}$  require computing a canonical linear map, which requires performing Gaussian elimination and can be done in time  $\text{poly}(m, \log q)$ .

The Turing-machine description of the decision procedure  $\mathcal{D}^{\text{LD}}$  can be uniformly computed from the integers  $(q, m, d, k)$  expressed in binary; the complexity of computing the description comes from describing the parameter tuple  $\text{ldparams}$ , which takes time that is at most polynomial in the bit length of  $(q, m, d, k)$ .  $\square$

## 7.2 The Magic Square game

We recall the *Magic Square game* of Mermin and Peres [Mer90, Per90, Ara02]. The Magic Square game is a simple self-test for EPR pairs (it tests for two of them). In addition, it allows one to test that a pair of observables *anticommutes*. Here we use it as a building block to construct the quantum low-degree test.

There are several formulations of the Magic Square game; here we present it as a *binary constraint satisfaction game* [CM14]. In this formulation of the game (denoted by  $\mathfrak{G}_{\text{MS}}$ ) there are 6 linear equations defined over 9 variables that take values in  $\mathbb{F}_2$ . The variables correspond to the cells of a  $3 \times 3$  grid, as depicted in Figure 4. Five of the equations correspond to the constraint that the sum of the variables in each row and the first two columns must be equal to 0, and the last equation requires that the sum of the variables in the last column must be equal to 1.

$x_1$	$x_2$	$x_3$
$x_4$	$x_5$	$x_6$
$x_7$	$x_8$	$x_9$

Figure 4: The Magic Square game

The question set  $\mathcal{T}^{\text{MS}}$  of the Magic Square game is the following:

$$\begin{aligned} \mathcal{T}^{\text{MC}} &= \{\text{CONSTRAINT}_i : i = 1, 2, \dots, 6\}, \\ \mathcal{T}^{\text{MV}} &= \{\text{VARIABLE}_j : j = 1, 2, \dots, 9\}, \\ \mathcal{T}^{\text{MS}} &= \mathcal{T}^{\text{MC}} \cup \mathcal{T}^{\text{MV}}. \end{aligned}$$

The questions  $\text{CONSTRAINT}_i$  for  $i \in \{1, 2, 3\}$  correspond to the three row constraints, the questions  $\text{CONSTRAINT}_4, \text{CONSTRAINT}_5$  correspond to the first two column constraints, and question  $\text{CONSTRAINT}_6$  corresponds to the third column constraint.

In the Magic Square game, the verifier first samples a constraint  $\text{CONSTRAINT}_i \in \mathcal{T}^{\text{MC}}$  uniformly at random, and then samples  $\text{VARIABLE}_j$ , one of the three variables in the row or column corresponding to  $\text{CONSTRAINT}_i$ , uniformly at random. One player is randomly assigned to be the  $\text{CONSTRAINT}$  player, and the other is assigned to be the  $\text{VARIABLE}$  player. The  $\text{CONSTRAINT}$  player is sent the question  $\text{CONSTRAINT}_i$  and is expected to respond with three bits  $(\beta_{v_1}, \beta_{v_2}, \beta_{v_3}) \in \mathbb{F}_2^3$ , where  $(v_1, v_2, v_3)$  are the indices of the three variables corresponding to  $\text{CONSTRAINT}_i$ . The  $\text{VARIABLE}$  player is given question  $\text{VARIABLE}_j$  and is expected to respond with a single bit  $\gamma \in \mathbb{F}_2$ . The players win if the  $\text{CONSTRAINT}$  player's answers satisfy the equation associated with  $\text{CONSTRAINT}_i$ , and  $\gamma = \beta_j$ . More precisely, the verifier samples an edge of the type graph (see Section 6)  $G^{\text{MS}}$  in Fig. 5, sends one endpoint to a random player, and the other endpoint to the other player.

The following theorem records the self-testing (also known as *rigidity*) properties of the Magic Square game. Its self-testing properties are crucial to the Pauli basis test. In particular, it is used to enforce anti-






Figure 5: Type graph  $G^{\text{MS}}$  for the Magic Square game.

commutation relations between certain pairs of operators. We will refer to this theorem in the proof of the soundness of the Pauli basis test (Theorem 7.14 below) in Appendix A.

**Theorem 7.10** (Rigidity of the Magic Square game). *Let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a strategy that succeeds in the Magic Square game  $\mathfrak{G}^{\text{MS}}$  with probability  $1 - \varepsilon$ , where  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  is a state. Then there exist local isometries  $\phi_A : \mathcal{H}_A \rightarrow \mathcal{H}_{A'} \otimes \mathcal{H}_{A''}$ ,  $\phi_B : \mathcal{H}_B \rightarrow \mathcal{H}_{B'} \otimes \mathcal{H}_{B''}$  (where  $\mathcal{H}_{A'}, \mathcal{H}_{B'} \cong (\mathbb{C}^2)^{\otimes 2}$  and  $\mathcal{H}_{A''}, \mathcal{H}_{B''}$  are finite dimensional) and a state  $|\text{AUX}\rangle \in \mathcal{H}_{A''} \otimes \mathcal{H}_{B''}$  such that*

1.  $\|\phi_A \otimes \phi_B |\psi\rangle - |\text{EPR}_2\rangle^{\otimes 2} \otimes |\text{AUX}\rangle\| \leq O(\sqrt{\varepsilon})$ ,
2. Letting  $\tilde{A}_a^x = \phi_A A_a^x \phi_A^\dagger$  and  $\tilde{B}_b^y = \phi_B B_b^y \phi_B^\dagger$ , we have that

$$\begin{aligned} \tilde{A}_b^{\text{VARIABLE}_1} \otimes I_{B'B''} &\approx_{\sqrt{\varepsilon}} (\sigma_b^X)_{A'} \otimes I_{A''B''} \\ \tilde{A}_b^{\text{VARIABLE}_5} \otimes I_{B'B''} &\approx_{\sqrt{\varepsilon}} (\sigma_b^Z)_{A'} \otimes I_{A''B''} \\ I_{A'A''} \otimes \tilde{B}_b^{\text{VARIABLE}_1} &\approx_{\sqrt{\varepsilon}} I_{A'A''B''} \otimes (\sigma_b^X)_{B'} \\ I_{A'A''} \otimes \tilde{B}_b^{\text{VARIABLE}_5} &\approx_{\sqrt{\varepsilon}} I_{A'A''B''} \otimes (\sigma_b^Z)_{B'} \end{aligned}$$

where the  $\approx_{\sqrt{\varepsilon}}$  statement holds with respect to the state  $|\text{EPR}_2\rangle_{A'B'}^{\otimes 2} \otimes |\text{AUX}\rangle_{A''B''}$  and the answer summation is over  $b \in \{0, 1\}$ . As a consequence, letting

$$\begin{aligned} \tilde{A}^{\text{VARIABLE}_1} &= \tilde{A}_0^{\text{VARIABLE}_1} - \tilde{A}_1^{\text{VARIABLE}_1} \\ \tilde{A}^{\text{VARIABLE}_5} &= \tilde{A}_0^{\text{VARIABLE}_5} - \tilde{A}_1^{\text{VARIABLE}_5} \end{aligned}$$

it holds that

$$\begin{aligned} \tilde{A}^{\text{VARIABLE}_1} \tilde{A}^{\text{VARIABLE}_5} \otimes I_{B'B''} &\approx_{\sqrt{\varepsilon}} -\tilde{A}^{\text{VARIABLE}_5} \tilde{A}^{\text{VARIABLE}_1} \otimes I_{B'B''} \\ I_{A'A''} \otimes \tilde{B}^{\text{VARIABLE}_1} \tilde{B}^{\text{VARIABLE}_5} &\approx_{\sqrt{\varepsilon}} -I_{A'A''} \otimes \tilde{B}^{\text{VARIABLE}_5} \tilde{B}^{\text{VARIABLE}_1}, \end{aligned}$$

where the notation  $M \approx_\delta N$  for  $M, N$  that are not POVM elements means

$$\langle \psi | (M - N)^\dagger (M - N) | \psi \rangle \leq O(\delta).$$

*Proof.* A proof of the rigidity of the Magic Square game can be found in [CS17, Theorem 6.9]. There are a couple minor differences between their rigidity statement and the one stated here. First, they state that the robustness of the Magic Square self-test is  $O(\varepsilon)$ ; however they specify the closeness between the actual and ideal states in terms of the trace norm, whereas we specify the closeness between  $\phi_A \otimes \phi_B | \psi \rangle$  and  $|\text{EPR}_2\rangle^{\otimes 2} \otimes |\text{AUX}\rangle$  in terms of the Euclidean distance, which translates to  $O(\sqrt{\varepsilon})$  instead of  $O(\varepsilon)$ . Second, their choice of ideal strategy specifies that the observables corresponding to VARIABLE<sub>1</sub> and VARIABLE<sub>5</sub> questions are  $I \otimes \sigma^Z$  and  $\sigma^Z \sigma^X \otimes \sigma^Z \sigma^X$ ; however under a local change of basis these are equivalent to  $\sigma^X \otimes I$  and  $\sigma^Z \otimes I$  respectively.  $\square$

We will need the following theorem, which shows that any pair of anticommuting observables can be used to form a value-1 strategy for the Magic Square game.

**Theorem 7.11.** *Let  $A = \{A_b\}_{b \in \mathbb{F}_2}$  and  $B = \{B_b\}_{b \in \mathbb{F}_2}$  be two-outcome projective measurements acting on  $(\mathbb{C}^q)^{\otimes n}$  which are consistent on  $|\text{EPR}_q\rangle^{\otimes n}$ , and let  $\mathcal{O}_A = A_0 - A_1$  and  $\mathcal{O}_B = B_0 - B_1$  be the corresponding observables. Suppose that  $\mathcal{O}_A \mathcal{O}_B = -\mathcal{O}_B \mathcal{O}_A$ . Then there exists a symmetric strategy  $\mathcal{S} = (\psi, M)$  for the Magic Square game with the following properties.*

1.  $\mathcal{S}$  is an SPCC strategy of value 1.
2. The state  $|\psi\rangle$  has the form  $|\psi\rangle = |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle$ .
3. For  $b \in \{0, 1\}$ , we have  $M_b^{\text{VARIABLE}_1} = A_b \otimes I$  and  $M_b^{\text{VARIABLE}_5} = B_b \otimes I$ .

*Proof.* The strategy  $\mathcal{S}$  is based on the canonical two-qubit strategy for the Magic Square game as described in, for example, [Ara02]. The state is  $|\psi\rangle = |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle$ . We specify the measurements of  $\mathcal{S}$  in Figure 6 as an operator solution for the Magic Square game, meant to be read as follows: each cell contains a two-outcome projective measurement  $\{E_0, E_1\}$  on  $(\mathbb{C}^q)^{\otimes n} \otimes \mathbb{C}^2$  written as its  $\pm 1$ -valued observable  $E_0 - E_1$ . When Player A or B receives the question VARIABLE <sub>$j$</sub>  for  $j \in \{1, \dots, 9\}$ , they measure their share of  $|\psi\rangle$  using the measurement specified by the cell corresponding to VARIABLE <sub>$j$</sub>  and receive a single-bit measurement. When they receive the question CONSTRAINT <sub>$i$</sub>  for  $i \in \{1, 2, \dots, 6\}$ , they simultaneously perform the three measurements in the corresponding row or column on  $|\psi\rangle$  to obtain three bits. For example, if Player A receives question VARIABLE<sub>1</sub>, they measure  $|\psi\rangle$  using the measurement  $\{A_0 \otimes I, A_1 \otimes I\}$  corresponding to the observable  $\mathcal{O}_A \otimes I$  (where the first operator acts on  $|\text{EPR}_q\rangle^{\otimes n}$  and the second acts on  $|\text{EPR}_2\rangle$ ). Similarly, on question VARIABLE<sub>5</sub>, they use the measurement  $\{B_0 \otimes I, B_1 \otimes I\}$ . This establishes Item 3 of the theorem.

First, we show that this gives a well-defined strategy. The VARIABLE <sub>$j$</sub>  measurements are well-defined because each cell contains a  $\pm 1$ -valued observable. This is obvious for all  $j \neq 9$ ; when  $j = 9$ , the bottom-right cell contains  $\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X$ . Because  $\mathcal{O}_A$  and  $\mathcal{O}_B$  anti-commute,

$$\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X = -\mathcal{O}_A \mathcal{O}_B \otimes \sigma^X \sigma^Z = \mathcal{O}_B \mathcal{O}_A \otimes \sigma^X \sigma^Z = (\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X)^\dagger. \quad (52)$$

As a result, this matrix is Hermitian. In addition,

$$(\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X)^2 = (\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X) \cdot (\mathcal{O}_B \mathcal{O}_A \otimes \sigma^X \sigma^Z) = (\mathcal{O}_A \mathcal{O}_B \cdot \mathcal{O}_B \mathcal{O}_A) \otimes (\sigma^Z \sigma^X \cdot \sigma^X \sigma^Z) = I,$$

$\mathcal{O}_A \otimes I$	$I \otimes \sigma^X$	$\mathcal{O}_A \otimes \sigma^X$
$I \otimes \sigma^Z$	$\mathcal{O}_B \otimes I$	$\mathcal{O}_B \otimes \sigma^Z$
$\mathcal{O}_A \otimes \sigma^Z$	$\mathcal{O}_B \otimes \sigma^X$	$\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X$

Figure 6: Observables for Magic Square strategy

where the first step uses Equation (52) and the final step uses the fact that  $\mathcal{O}_A, \mathcal{O}_B, \sigma^X, \sigma^Z$  are  $\pm 1$ -valued observables and hence square to the identity. As a result, this matrix is Hermitian and squares to the identity. Therefore, it is a  $\pm 1$ -valued observable.

As for the  $\text{CONSTRAINT}_i$  measurements, we must show that the three measurements in each row and column are simultaneously measurable. This is equivalent to the three  $\pm 1$ -valued observables being simultaneously diagonalizable, which is equivalent to them being pairwise commuting. This can be easily verified for the cases of  $i = 1, 2, 4, 5$  (i.e. the first two rows and columns). In the case of  $i = 3$ , commutativity of  $\mathcal{O}_A \otimes \sigma^Z$  and  $\mathcal{O}_B \otimes \sigma^X$  follows from Equation (52). Since these two matrices commute, they also commute with their product  $(\mathcal{O}_A \otimes \sigma^Z)(\mathcal{O}_B \otimes \sigma^X) = \mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X$ . The case of  $i = 6$  is similar.

By construction,  $\mathcal{S}$  is symmetric, and we have already shown that it is projective. It remains to show that it is commuting, consistent, and value 1. To show that it is commuting, it suffices to show that the measurement for each cell is simultaneously measurable with all three measurements in its row or column, which was already proved above. Now we show consistency of the measurements. We will show this by instead showing consistency of their observables. By this we mean the following: if  $E = \{E_0, E_1\}$  is a two-outcome projective measurement on  $\mathcal{H}$  then its corresponding  $\pm 1$ -valued observable  $\mathcal{O} = E_0 - E_1$  is consistent on a state  $|\phi\rangle \in \mathcal{H} \otimes \mathcal{H}$  if

$$\mathcal{O} \otimes I_B \cdot |\phi\rangle = I_A \otimes \mathcal{O} \cdot |\phi\rangle.$$

This is in fact equivalent to the notion of  $E$  being consistent on  $|\phi\rangle$  from Definition 5.9. To see this, if  $E$  is consistent on  $|\phi\rangle$ , then so is  $\mathcal{O}$ , because

$$\mathcal{O} \otimes I_B \cdot |\phi\rangle = (E_0 - E_1) \otimes I_B \cdot |\phi\rangle = I_A \otimes (E_0 - E_1) \cdot |\phi\rangle = I_A \otimes \mathcal{O} \cdot |\phi\rangle,$$

where the second equality is by consistency of  $E$ . On the other hand, suppose  $\mathcal{O}$  is consistent on  $|\phi\rangle$ . Then because  $E_0 + E_1 = I$ , we can write  $E_0 = \frac{1}{2} \cdot (I + \mathcal{O})$  and  $E_1 = \frac{1}{2} \cdot (I - \mathcal{O})$ . As a result,

$$E_0 \otimes I_B \cdot |\phi\rangle = \frac{1}{2} \cdot (I + \mathcal{O}) \otimes I_B \cdot |\phi\rangle = I_A \otimes \frac{1}{2} \cdot (I + \mathcal{O}) \cdot |\phi\rangle = I_A \otimes E_0 \cdot |\phi\rangle,$$

and similarly for  $E_1$ . Thus,  $E$  is consistent on  $|\phi\rangle$ . This proves the equivalence.

Now we use this to prove consistency of  $\mathcal{S}$ . To begin, we note that because  $A$  and  $B$  are consistent on  $|\text{EPR}_q\rangle^{\otimes n}$  by assumption, so are their  $\pm 1$ -valued observables  $\mathcal{O}_A$  and  $\mathcal{O}_B$ . Next, we note that  $\sigma^X$  and  $\sigma^Z$  are consistent on  $|\text{EPR}_2\rangle$ ; to verify this, we recall that  $\sigma^X|0\rangle = |1\rangle$  and  $\sigma^X|1\rangle = |0\rangle$ , and so

$$\begin{aligned} \sigma^X \otimes I_B \cdot |\text{EPR}_2\rangle &= \sigma^X \otimes I_B \cdot \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle) \\ &= \frac{1}{\sqrt{2}}(|10\rangle + |01\rangle) = I_A \otimes \sigma^X \cdot \frac{1}{\sqrt{2}}(|11\rangle + |00\rangle) = I_A \otimes \sigma^X \cdot |\text{EPR}_2\rangle. \end{aligned}$$

This shows  $\sigma^X$  is consistent on  $|\text{EPR}_2\rangle$ , and a similar argument shows the same for  $\sigma^Z$ . We now use these two facts to show that the observable in each cell of Figure 6 is consistent on  $|\psi\rangle$ . To see why, consider the  $j = 9$  case:

$$\begin{aligned}
& (\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X)_A \otimes I_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle \\
&= (\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z)_A \otimes (I \otimes \sigma^X)_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle && \text{(consistency of } \sigma^X) \\
&= (\mathcal{O}_A \mathcal{O}_B \otimes I)_A \otimes (I \otimes \sigma^X \sigma^Z)_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle && \text{(consistency of } \sigma^Z) \\
&= (\mathcal{O}_A \otimes I)_A \otimes (\mathcal{O}_B \otimes \sigma^X \sigma^Z)_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle && \text{(consistency of } \mathcal{O}_B) \\
&= I_A \otimes (\mathcal{O}_B \mathcal{O}_A \otimes \sigma^X \sigma^Z)_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle && \text{(consistency of } \mathcal{O}_A) \\
&= I_A \otimes (\mathcal{O}_A \mathcal{O}_B \otimes \sigma^Z \sigma^X)_B \cdot |\text{EPR}_q\rangle^{\otimes n} \otimes |\text{EPR}_2\rangle. && \text{(by Equation (52))}
\end{aligned}$$

The remaining cases of  $j \in \{1, \dots, 8\}$  are similar, and we omit them. This shows that the observable in each cell is consistent on  $|\psi\rangle$ , and as a result the corresponding two-outcome projective measurements are consistent as well. As a result, the  $\text{VARIABLE}_j$  measurements are consistent.

As for the  $\text{CONSTRAINT}_i$  measurements, each such measurement  $\{F_{a,b,c}\}_{a,b,c \in \{0,1\}}$  is of the form  $F_{a,b,c} = E_a^1 \cdot E_b^2 \cdot E_c^3$ , where  $E^1$ ,  $E^2$ , and  $E^3$  are  $\text{VARIABLE}$  measurements. But then consistency of  $F$  follows from the  $\text{VARIABLE}$  consistencies:

$$\begin{aligned}
F_{a,b,c} \otimes I_B |\psi\rangle &= (E_a^1 \cdot E_b^2 \cdot E_c^3)_A \otimes I_B |\psi\rangle \\
&= (E_a^1 \cdot E_b^2)_A \otimes (E_c^3)_B |\psi\rangle && \text{(consistency of } E^3) \\
&= (E_a^1)_A \otimes (E_c^3 \cdot E_b^2)_B |\psi\rangle && \text{(consistency of } E^2) \\
&= I_A \otimes (E_c^3 \cdot E_b^2 \cdot E_a^1)_B |\psi\rangle && \text{(consistency of } E^1) \\
&= I_A \otimes (E_a^1 \cdot E_b^2 \cdot E_c^3)_B |\psi\rangle && \text{(commutativity of } E^1, E^2, \text{ and } E^3) \\
&= I_A \otimes F_{a,b,c} |\psi\rangle.
\end{aligned}$$

Hence, all measurements are consistent.

Since all measurements are consistent, this implies that the answer bit of the player receiving a  $\text{VARIABLE}$  question is always consistent with the corresponding answer bit of the player receiving the  $\text{CONSTRAINT}$  question. Similarly, the answers of the player receiving the  $\text{CONSTRAINT}$  question always satisfy the given constraint; observe that in all rows and the first two columns, the observables multiply to  $I$ , whereas the observables in the last column multiply to  $-I$ . This implies that the strategy is value-1. To see why, let us again consider a  $\text{CONSTRAINT}_i$  measurement, which we write as  $F = \{F_{a,b,c}\}_{a,b,c \in \{0,1\}}$ . As above, it can be written as  $F_{a,b,c} = E_a^1 \cdot E_b^2 \cdot E_c^3$ , where  $E^1$ ,  $E^2$ , and  $E^3$  are  $\text{VARIABLE}$  measurements. Consider the measurement  $S = \{S_0, S_1\}$  that corresponds to measuring  $\{F_{a,b,c}\}$  and then outputting the sum of measured values of  $a$ ,  $b$ , and  $c$ . In other words,

$$S_0 = F_{0,0,0} + F_{0,1,1} + F_{1,0,1} + F_{1,1,0}, \quad S_1 = F_{0,0,1} + F_{0,1,0} + F_{1,0,0} + F_{1,1,1}.$$

The following expression gives a convenient formula for the  $\pm 1$ -valued observable  $S_0 - S_1$ :

$$(E_0^1 - E_1^1) \cdot (E_0^2 - E_1^2) \cdot (E_0^3 - E_1^3) = E_0^1 E_0^2 E_0^3 - E_0^1 E_0^2 E_1^3 - E_0^1 E_1^2 E_0^3 + E_0^1 E_1^2 E_1^3 + \dots = S_0 - S_1.$$

When we are looking at  $\text{CONSTRAINT}_i$  for  $i \in \{1, 2, 3, 4, 5\}$ , the left-hand side of this equation is equal to  $I$ , and so  $I = S_0 - S_1$ . Because  $S_0$  and  $S_1$  are both positive semidefinite, this is true only if  $S_0 = I$  and  $S_1 = 0$ ,

which implies that the three bits output by  $F$  always sum to 0. As a result, this strategy always satisfies the  $\text{CONSTRAINT}_i$  question. On the other hand, when  $i = 6$ , then the left-hand side of this equation is equal to  $-I$ , and so  $S_1 = I$ , and the three bits sum to 1. Thus, the strategy always satisfies the  $\text{CONSTRAINT}_6$  question as well. This concludes the proof.  $\square$

### 7.3 The Pauli basis test

We introduce the quantum low-degree test of [NV18a] in the form of a slight modification to it by [NW19] known as the *Pauli basis test*. Informally, the quantum low-degree test asks the players to measure a large number of qubits and return a highly compressed version of the measurement outcome. The Pauli basis test simply asks that the players return their uncompressed measurement outcomes, and it is designed by direct reduction to the quantum low-degree test. In Section 7.3.1 we describe the Pauli basis test as a nonlocal game  $\mathfrak{G}^{\text{PAULI}}$  and show that its question distribution is implementable via a CL distribution, as we did with the classical low-degree test in Section 7.1. In Section 7.3.2 we exhibit canonical parameters for the Pauli basis test and give bounds on the time complexity of executing the test.

#### 7.3.1 The game

We start by discussing parameter settings. The game  $\mathfrak{G}^{\text{PAULI}}$  is parametrized by a tuple

$$\text{qldparams} = (q, m, d),$$

where  $q, m, d \in \mathbb{N}$  are integers. We sometimes write  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  to emphasize the dependence of the Pauli basis test on the parameters.

Informally, the test is meant to certify that the players share a state of the form  $|\text{EPR}_q\rangle^{\otimes M}$ , where  $M = 2^m$ . Its question set includes questions that are axis-parallel/diagonal lines and points in  $\mathbb{F}_q^m$ , which are meant to correspond to questions in the classical low-degree test, and questions of the form  $(\text{PAULI}, W)$ , for  $W \in \{X, Z\}$ . Upon receipt of a question of the latter form, the players are expected to perform the POVM  $\{\tau_h^W\}_{h \in \mathbb{F}_q^m}$  and report the outcome  $h$  as their answer.

The key idea behind the test is for the provers to encode their outcomes using the *low-degree encoding* from Section 3.4. Given an outcome  $h \in \mathbb{F}_q^m$ , the low-degree encoding of  $h$  is the polynomial  $g_h : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ . Rather than asking the provers to always return the entire string  $h$ , many of the subtests in the game  $\mathfrak{G}^{\text{PAULI}}$  will “probe” limited information about  $h$  as follows: the verifier provides the provers with a string  $u_W \in \mathbb{F}_q^m$ , and using this they are expected to evaluate  $g_h(u_W) \in \mathbb{F}_q$  and return it to the verifier. For a fixed value of  $W$ , then, the prover’s responses to these “probes” should be consistent with some low-degree polynomial in the input  $u_W$ . To check this, the verifier can perform the classical low-degree test from Section 7.1. This ensures that the prover’s responses for the  $W = X$  basis are consistent with each other, as well as their responses for the  $W = Z$  basis, although it does not test consistency between the  $X$  and  $Z$  responses.

The next set of subtests the verifier performs ensure that the provers’  $X$  and  $Z$  measurements are consistent with each other. On the  $X$  side, consider the measurement in which the prover performs the  $\{\tau_h^X\}$  POVM, receives outcome  $h_X$ , and outputs  $g_{h_X}(u_X) \in \mathbb{F}_q$ . For technical reasons that will become clear shortly, it will be convenient for the prover’s measurement to have two outcomes rather than  $q$  outcomes. To accomplish this, the verifier also provides the prover with an element  $r_X \in \mathbb{F}_q$ ; given this, the prover should output not  $g_{h_X}(u_X) \in \mathbb{F}_q$  but  $\text{tr}(g_{h_X}(u_X) \cdot r_X)$ , which is an element of  $\mathbb{F}_2$ . Similarly, on the  $Z$  side, suppose the prover is provided  $r_Z \in \mathbb{F}_q$ , performs  $\{\tau_h^Z\}$  to receive  $h_Z$ , and outputs  $\text{tr}(g_{h_Z}(u_Z) \cdot r_Z) \in \mathbb{F}_2$ . Together, these two are a pair of two-outcome measurements, one in the  $X$  basis and one in the  $Z$  basis, and as it turns

out, this pair of measurements either commutes with each other or anticommutes with each other (this is why we modified the measurements to have only two outcomes). The key quantity to determine which of these is the case is the following complicated-looking expression:

$$\gamma = \text{tr}((\text{ind}_m(u_X) \cdot r_X) \cdot (\text{ind}_m(u_Z) \cdot r_Z)) .$$

If  $\gamma = 0$ , then these two measurements commute, and if  $\gamma = 1$ , then these two measurements anticommute (this fact is established in the proof of Lemma 7.13 below). In the  $\gamma = 0$  case, the verifier can check whether these two measurements commute by performing the ‘‘commutation check’’, which asks the provers to simultaneously measure both  $\text{tr}(g_{h_X}(u_X) \cdot r_X)$  and  $\text{tr}(g_{h_Z}(u_Z) \cdot r_Z)$  and report their values. In the  $\gamma = 1$  case, on the other hand, the measurements anticommute, which implies that they cannot simultaneously be measured. In spite of this, the verifier can still check that the measurements anticommute by using the Magic Square game from Section 7.2 above. As was shown in Theorem 7.11, any pair of anticommuting measurements can be used to form a perfect strategy for the Magic Square game (and indeed the reverse is true as well: any perfect strategy for the Magic Square game entails a pair of anticommuting measurements).

Together, these give the main subtests for the game  $\mathfrak{G}^{\text{PAULI}}$ .

**Definition 7.12** (Admissible parameters). We say that the tuple  $\text{qldparams} = (q, m, d)$  is *admissible* if  $q$  is an admissible field size (Definition 3.15) and  $m|q$ .

**Question distribution.** We now describe the question distribution of the game  $\mathfrak{G}^{\text{PAULI}}$ , and show that it is a (typed) CL distribution. The question types in the game  $\mathfrak{G}^{\text{PAULI}}$  are

$$\mathcal{T}^{\text{PAULI}} = (\{\text{POINT}, \text{ALINE}, \text{DLINE}, \text{PAULI}, \text{PAIR}\} \times \{X, Z\}) \cup \mathcal{T}^{\text{MS}} \cup \{\text{PAIR}\} , \quad (53)$$

where  $\mathcal{T}^{\text{MS}}$  is the question type set of the Magic Square game defined in Section 7.2.

Before presenting the CL functions of the Pauli basis test, we first give an intuitive description of the question distribution of the Pauli basis test: a pair of questions  $((t_A, x_A), (t_B, x_B))$  in  $\mathfrak{G}^{\text{PAULI}}$  can be sampled via the following procedure:

1. Sample a pair of types by sampling an edge  $(t_A, t_B)$  of the graph  $G^{\text{PAULI}}$  given in Figure 7 uniformly at random (including the self-loops).
2. Sample the following uniformly at random:
  - (a) (*Points*)  $u_X, u_Z \in \mathbb{F}_q^m$ ,
  - (b) (*Directions*)  $s \in \mathbb{F}_q, v \in \mathbb{F}_q^m$ ,
  - (c) (*Qubit basis for (anti-)commutation*)  $r_X, r_Z \in \mathbb{F}_q$ .
3. For  $w \in \{A, B\}$  and  $W \in \{X, Z\}$ ,
  - (a) If  $t_w = (\text{POINT}, W)$ , then set  $x_w = u_W$ ,
  - (b) If  $t_w = (\text{ALINE}, W)$ , then set  $x_w = (u_0, s)$ , where  $u_0 = L_{e_i}^{\text{LN}}(u_W)$ , with  $i = \chi(s)$  (see Section 7.1.1 for definition of  $L^{\text{LN}}$  and  $\chi(s)$ ),
  - (c) If  $t_w = (\text{DLINE}, W)$ , then set  $x_w = (u_0, s, v')$ , where  $i = \chi(s)$ ,  $v' = \pi_{i-1}(v)$ , and  $u_0 = L_{v'}^{\text{LN}}(u_W)$ ,
  - (d) If  $t_w = \text{CONSTRAINT}_i$  for some  $i \in \{1, \dots, 6\}$ , then set  $x_w = (u_X, u_Z, r_X, r_Z)$ ,

- (e) If  $t_w = \text{VARIABLE}_j$  for some  $j \in \{1, \dots, 9\}$ , then set  $x_w = (u_X, u_Z, r_X, r_Z)$ ,
- (f) If  $t_w = \text{PAIR}$ , then set  $x_w = (u_X, u_Z, r_X, r_Z)$ ,
- (g) If  $t_w = (\text{PAIR}, W)$ , then set  $x_w = (u_X, u_Z, r_X, r_Z)$ ,
- (h) If  $t_w = (\text{PAULI}, W)$ , then set  $x_w = 0$ .




Figure 7: Graph  $G^{\text{PAULI}}$  for the Pauli basis test. Each vertex also has a self-loop which is not drawn on the figure for clarity.

We now specify the corresponding CL functions for each of the question types in the Pauli basis test.

**Conditional linear functions for the Pauli basis test.** The CL functions corresponding to the Pauli basis test question distribution are parameterized by the parameter tuple  $\text{qldparams} = (q, m, d)$ . Let  $V^{\text{PAULI}}$  denote the linear space  $(\mathbb{F}_q^m)^2 \times \mathbb{F}_q \times \mathbb{F}_q^m \times (\mathbb{F}_q)^2$ . The space  $V^{\text{PAULI}}$  is decomposed into a direct sum of the following register subspaces:  $V_X, V_Z$  (which are  $m$ -dimensional),  $V_I$  (which is 1-dimensional),  $V_V$  (which is  $m$ -dimensional), and  $V_{R_X}, V_{R_Z}$  (which are 1-dimensional). We identify elements of  $V^{\text{PAULI}}$  as tuples  $(u_X, u_Z, s, v, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q \times \mathbb{F}_q^m \times (\mathbb{F}_q)^2$ . We define CL functions  $L_t : V^{\text{PAULI}} \rightarrow V^{\text{PAULI}}$  for every type  $t \in \mathcal{T}^{\text{PAULI}}$ :

1. For  $W \in \{X, Z\}$ , define  $L_{\text{POINT}, W}(u_X, u_Z, s, v, r_X, r_Z) = L_{\text{POINT}}(u_W, s, v)$  where  $L_{\text{POINT}}$  is the 1-level CL function defined in Equation (48).<sup>29</sup>
2. For  $W \in \{X, Z\}$ , define  $L_{\text{ALINE}, W}(u_X, u_Z, s, v, r_X, r_Z) = L_{\text{ALINE}}(u_W, s, v)$  where  $L_{\text{ALINE}}$  is the 2-level CL function defined in Equation (49).

<sup>29</sup>The range  $V_X \oplus V_I \oplus V_V$  or  $V_Z \oplus V_I \oplus V_V$  of  $L_{\text{POINT}}$  is embedded in  $V^{\text{PAULI}}$  in the natural way. The same convention is used also in Item 2 and 3 for  $L_{\text{ALINE}}$  and  $L_{\text{DLINE}}$ .

3. For  $W \in \{X, Z\}$ , define  $L_{\text{DLINE},W}(u_X, u_Z, s, v, r_X, r_Z) = L_{\text{DLINE}}(u_W, s, v)$  where  $L_{\text{DLINE}}$  is the 3-level CL function defined in Equation (51).
4. For  $t \in \mathcal{T}^{\text{MS}} \cup \{\text{PAIR}, (\text{PAIR}, X), (\text{PAIR}, Z)\}$ ,  $L_t(u_X, u_Z, s, v, r_X, r_Z) = (u_X, u_Z, 0, 0, r_X, r_Z)$ , i.e., it projects onto  $V_X \oplus V_Z \oplus V_{R_X} \oplus V_{R_Z}$ . Thus  $L_t$  is a 1-level CL function.
5. For  $W \in \{X, Z\}$ , define  $L_{\text{PAULI},W} = 0$  as the identically 0 function. Thus it is a 0-level CL function.

We note that the maps  $L_{\text{POINT},W}, L_{\text{ALINE},W}, L_{\text{DLINE},W}$  act as the zero function on  $V_{\overline{W}} \oplus V_{R_X} \oplus V_{R_Z}$  where  $\overline{W} = X$  if  $W = Z$  and  $\overline{W} = Z$  if  $W = X$ .

The question distribution of  $\mathfrak{G}^{\text{PAULI}}$  is thus a typed CL distribution on  $\mathcal{T}^{\text{PAULI}} \times V^{\text{PAULI}} \times \mathcal{T}^{\text{PAULI}} \times V^{\text{PAULI}}$  where  $(t_A, x_A, t_B, x_B)$  is sampled by first uniformly sampling an edge  $(t_A, t_B)$  from the graph  $G^{\text{PAULI}}$  defined in Figure 7, sampling a uniformly random  $z \in V^{\text{PAULI}}$ , and then setting  $x_w = L_{t_w}(z)$  for  $w \in \{A, B\}$ .

**Decision procedure.** The decision procedure for  $\mathfrak{G}^{\text{PAULI}}$  is presented in Figure 8. Similarly to Figure 3, we provide a table that summarizes a parsing scheme for the questions and answers, depending on the type of question. The answers are bit strings that are interpreted as more structured objects such as elements over  $\mathbb{F}_q$ , vectors, or polynomials, depending on the question. In the “low-degree check”, the decision procedure  $\mathcal{D}^{\text{PAULI}}$  calls the classical low-degree decision procedure  $\mathcal{D}^{\text{LD}}$  parametrized by the tuple  $\text{ldparams} = (q, m, d, 1)$  (defined in Section 7.1) as a subroutine.



Type	Question Content	Answer Format
(POINT, W)	$y \in \mathbb{F}_q^m$	Element of $\mathbb{F}_q$
(ALINE, W)	$(u_W, s) \in \mathbb{F}_q^m \times \mathbb{F}_q$	Polynomial $f : \mathbb{F}_q \rightarrow \mathbb{F}_q$
(DLINE, W)	$(u_W, s, v) \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$	Polynomial $f : \mathbb{F}_q \rightarrow \mathbb{F}_q$
PAIR	$(u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q^2$	$(\beta_X, \beta_Z) \in \mathbb{F}_2^2$
(PAIR, W)	$(u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q^2$	Element of $\mathbb{F}_2$
CONSTRAINT <sub>i</sub>	$(u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q^2$	$(\alpha_{v_1}, \alpha_{v_2}, \alpha_{v_3}) \in \mathbb{F}_2^3$
VARIABLE <sub>j</sub>	$(u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q^2$	Element of $\mathbb{F}_2$
(PAULI, W)	0	Element of $\mathbb{F}_q^M$

Table: Question and answer format of the Pauli basis game.

On input  $(t_A, x_A, t_B, x_B, a_A, a_B)$ , the decision procedure  $\mathcal{D}^{\text{PAULI}}$  performs the following checks for  $w \in \{A, B\}$ :

1. **(Consistency check).** If  $t_A = t_B$ , accept iff  $a_A = a_B$ .
2. **(Low-degree).** Let  $\text{ldparams} = (q, m, d, 1)$ .
  - (a) If  $t_w = (\text{POINT}, W), t_{\bar{w}} = (\text{ALINE}, W)$ , accept if  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  accepts  $(\text{POINT}, x_w, \text{ALINE}, x_{\bar{w}}, a_w, a_{\bar{w}})$ .
  - (b) If  $t_w = (\text{POINT}, W), t_{\bar{w}} = (\text{DLINE}, W)$ , accept if  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  accepts  $(\text{POINT}, x_w, \text{DLINE}, x_{\bar{w}}, a_w, a_{\bar{w}})$ .
3. **(Consistency check).** If  $t_w = (\text{POINT}, W), t_{\bar{w}} = (\text{PAULI}, W)$ , accept if  $g_{a_{\bar{w}}}(x_w) = a_w$ , where  $g_{a_{\bar{w}}}$  is the low-degree encoding of  $a_{\bar{w}} \in \mathbb{F}_q^M$  defined in Section 3.4.

In the remaining four cases, the decision procedure first computes the number

$$\gamma = \text{tr}((\text{ind}_m(u_X) \cdot r_X) \cdot (\text{ind}_m(u_Z) \cdot r_Z)), \quad (54)$$

where we recall the  $\text{ind}_m(\cdot)$  vector from Section 3.4.

4. **(Commutation check).** If  $t_w = (\text{PAIR}, W), t_{\bar{w}} = \text{PAIR}$ , accept if  $a_w = \beta_W$  or  $\gamma \neq 0$ .
5. **(Consistency check).** If  $t_w = (\text{POINT}, W), t_{\bar{w}} = (\text{PAIR}, W)$ , accept if  $\text{tr}(a_w r_W) = a_{\bar{w}}$  or  $\gamma \neq 0$ .
6. **(Magic square check).** If  $t_w = \text{CONSTRAINT}_i, t_{\bar{w}} = \text{VARIABLE}_j$ , accept if  $\gamma = 0$ , or  $a_w$  satisfies constraint  $\text{CONSTRAINT}_i$  and  $\alpha_j = a_{\bar{w}}$ .
7. **(Consistency check).** If  $t_w = (\text{POINT}, W), t_{\bar{w}} = \text{VARIABLE}_j$ , accept if  $\gamma = 0$  or if  $j = 1, W = X$ , and  $\text{tr}(a_w r_X) = a_{\bar{w}}$  or if  $j = 5, W = Z$ , and  $\text{tr}(a_w r_Z) = a_{\bar{w}}$ .

Figure 8: Specification of the decision procedure  $\mathcal{D}_{\text{qldparams}}^{\text{PAULI}}$ .

We describe an honest, value-1 PCC strategy for the Pauli basis test.

**Lemma 7.13.** *The Pauli basis test  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  has a value-1 SPCC strategy.*

*Proof.* We begin by specifying the value-1 strategy  $\mathcal{S} = (|\psi\rangle, E)$ . The state is

$$|\psi\rangle = |\text{EPR}_q\rangle^{\otimes M} \otimes |\text{EPR}_2\rangle.$$

Now we specify the measurements. We start with measurements associated with questions of type POINT, ALINE, DLINE, and PAULI. Using notation introduced in Section 3.4, for  $W \in \{X, Z\}$ ,

$$\begin{aligned} E_a^{(\text{POINT}, W), y} &= \tau_{[g \cdot (y)=a]}^W \otimes I, \\ E_f^{(\text{ALINE}, W), \ell} &= \tau_{[(g \cdot) |_{\ell}=f]}^W \otimes I, \\ E_f^{(\text{DLINE}, W), \ell} &= \tau_{[(g \cdot) |_{\ell}=f]}^W \otimes I, \\ E_a^{(\text{PAULI}, W)} &= \tau_a^W \otimes I. \end{aligned}$$

Here, the bracket notation used to post-process measurement outcomes is defined in Definition 3.23. Thus, for example, the measurement on question  $((\text{POINT}, W), y)$  corresponds to first performing the measurement  $\{\tau_h^W\}_{h \in \mathbb{F}_q^M}$ , receiving an outcome  $h \in \mathbb{F}_q^M$ , and then outputting the value  $a = g_h(y)$ .

For ALINE and DLINE questions, the question content  $\ell$  denotes either an axis-parallel line  $\ell(u_0, e_i)$  for some  $i \in \{1, 2, \dots, m\}$ , or a diagonal line  $\ell(u_0, v')$  for some  $v' \in \mathbb{F}_q^m$ . As described at the beginning of Section 7.3.1, axis-parallel lines are specified by pairs  $(u_0, s) \in \mathbb{F}_q^m \times \mathbb{F}_q$  and diagonal lines are specified by triples  $(u_0, s, v) \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$ . The measurement on question  $((\text{ALINE}, W), \ell)$ , for example, corresponds to first performing the measurement  $\{\tau_h^W\}_{h \in \mathbb{F}_q^M}$  to get an outcome  $h \in \mathbb{F}_q^M$ , and reporting the univariate polynomial  $f_\ell(t) = g_h(u_0 + te_i)$  where  $\ell$  is specified by  $(u_0, s)$  and  $i = \chi(s)$ .

Next we specify the POVMs associated with questions of type CONSTRAINT, VARIABLE, and PAIR. Questions with these question types have a question content that is a tuple  $\omega = (u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times \mathbb{F}_q^2$ . Given such a tuple, consider the two  $\mathbb{F}_2$ -valued POVMs  $A = \{A_b\}_{b \in \mathbb{F}_2}$  and  $B = \{B_b\}_{b \in \mathbb{F}_2}$  defined as

$$A_b = \tau_{[\text{tr}(g \cdot (u_X) r_X)=b]}^X \otimes I, \quad (55)$$

$$B_b = \tau_{[\text{tr}(g \cdot (u_Z) r_Z)=b]}^Z \otimes I. \quad (56)$$

We would like to determine when the two observables  $\mathcal{O}_A = A_0 - A_1$  and  $\mathcal{O}_B = B_0 - B_1$  commute or anti-commute. Towards this we derive alternative expressions for these observables from which their commutativity becomes plain from inspection. We begin by inspecting the first matrices on the right-hand sides of Equations (55) and (56):

$$\begin{aligned} \tau_{[\text{tr}(g \cdot (u_W) r_W)=b]}^W &= \sum_{h: \text{tr}(g_h(u_W) r_W)=b} \tau_h^W \\ &= \sum_{h: \text{tr}((h \cdot \text{ind}_m(u_W)) \cdot r_W)=b} \tau_h^W, \end{aligned} \quad (57)$$

where (57) follows by Definition 13. As a result,

$$\begin{aligned}
& \tau_{[\text{tr}(g \cdot (u_W) r_W)=0]}^W - \tau_{[\text{tr}(g \cdot (u_W) r_W)=1]}^W \\
&= \sum_{h:\text{tr}((h \cdot \text{ind}_m(u_W)) \cdot r_W)=0} \tau_h^W - \sum_{h:\text{tr}((h \cdot \text{ind}_m(u_W)) \cdot r_W)=1} \tau_h^W \\
&= \sum_h (-1)^{\text{tr}((h \cdot \text{ind}_m(u_W)) \cdot r_W)} \tau_h^W \\
&= \tau^W(\text{ind}_m(u_W) r_W), \tag{58}
\end{aligned}$$

where the last step uses (20). As a result, Equation (58) and Equations (55) and (56) imply that

$$\mathcal{O}_A = \tau^X(\text{ind}_m(u_X) r_X) \otimes I, \quad \mathcal{O}_B = \tau^Z(\text{ind}_m(u_Z) r_Z) \otimes I.$$

Now, let  $\gamma = \text{tr}((\text{ind}_m(u_X) r_X) \cdot (\text{ind}_m(u_Z) r_Z)) \in \mathbb{F}_2$ , as in Equation (54). Then by Equation (16),

$$\mathcal{O}_A \mathcal{O}_B = (-1)^\gamma \mathcal{O}_B \mathcal{O}_A.$$

As a result,  $\gamma$  quantifies whether the observables  $\mathcal{O}_A$  and  $\mathcal{O}_B$  commute, and therefore whether the measurements  $A$  and  $B$  commute. If  $\gamma = 0$ , they commute. If  $\gamma = 1$ , they anti-commute. We now specify the POVMs associated with questions of type CONSTRAINT, VARIABLE, and PAIR, considering separately the cases when  $\gamma = 0$  or  $\gamma = 1$ .

1. If  $\gamma = 0$ , for each  $\beta_X, \beta_Z \in \mathbb{F}_2$  define

$$E_{\beta_X, \beta_Z}^{\text{PAIR}, \omega} = A_{\beta_X} \cdot B_{\beta_Z}, \tag{59}$$

$$E_a^{(\text{PAIR}, X), \omega} = A_a, \quad E_a^{(\text{PAIR}, Z), \omega} = B_a.$$

The POVM associated with questions of type CONSTRAINT and VARIABLE are defined to be trivial. In particular, we define

$$E_{0,0,0}^{\text{CONSTRAINT}_i, \omega} = E_0^{\text{VARIABLE}_j, \omega} = I,$$

and the remaining POVM elements in these measurements are set to be zero.

2. If  $\gamma = 1$ , then  $\mathcal{O}_A$  and  $\mathcal{O}_B$  anti-commute. In this case we define measurements  $E^{\text{CONSTRAINT}_i, \omega}$  and  $E^{\text{VARIABLE}_j, \omega}$ , for all  $i \in \{1, \dots, 6\}$  and  $j \in \{1, \dots, 9\}$  to be those guaranteed by Theorem 7.11. Measurements associated with inputs of type PAIR are defined to be trivial. In particular, we define

$$E_{0,0}^{\text{PAIR}, \omega} = E_0^{(\text{PAIR}, X), \omega} = E_0^{(\text{PAIR}, Z), \omega} = I,$$

and the remaining matrices in these measurements are set to be zero.

This completes the specification of the strategy.

Now, we show that  $\mathcal{S}$  is a value-1 SPCC strategy. It is clearly symmetric and projective. To show that it is consistent, we note that all measurements are Pauli basis measurements, which are consistent, or measurements produced by Theorem 7.11, which are also consistent. The only exception is the PAIR measurement in the  $\gamma = 0$  case, which by Equation (59) is a product of two commuting, consistent measurements, and so it is therefore also consistent. To show that it is commuting, we note that for each  $W \in \{X, Z\}$ , all (POINT,  $W$ ), (ALINE,  $W$ ), (DLINE,  $W$ ) and (PAULI,  $W$ ) measurements commute as they are all measurements in the

Pauli  $W$  basis. Next, if  $\gamma = 0$  then the CONSTRAINT and VARIABLE measurements commute trivially, and for  $W \in \{X, Z\}$ , the (POINT,  $W$ ) measurement commutes with the (PAIR,  $W$ ) measurement, as they are both  $W$  basis measurements, and the measurement  $E_{\beta_X, \beta_Z}^{\text{PAIR}, \omega} = A_{\beta_X} \cdot B_{\beta_Z}$  commutes with  $E_a^{(\text{PAIR}, W), \omega}$  because  $A$  and  $B$  commute. On the other hand, if  $\gamma \neq 0$ , then the PAIR measurements commute trivially, and the CONSTRAINT and VARIABLE measurements commute by Theorem 7.11. Finally, the (POINT,  $X$ ) measurement commutes with VARIABLE<sub>1</sub> because both are  $X$ -basis measurements, and likewise both (POINT,  $Z$ ) and VARIABLE<sub>5</sub> are  $Z$ -basis measurements.

It remains to show that  $\mathcal{S}$  is value-1. Consider first the first three tests executed by the decision procedure in Figure 8. The strategy passes the consistency checks with probability 1 because it is projective and consistent. It passes the low-degree checks because it answers those consistently with an honest strategy in the classical low-degree test.

Next, consider the remaining four tests. Fix an  $\omega = (u_X, u_Z, r_X, r_Z)$  and  $\gamma$  as in (54). If  $\gamma = 0$ , then the strategy passes the commutation check with probability 1 by construction. As for the consistency check in Item 5, we can write the (POINT,  $W$ ) measurement as follows:

$$E_{[\text{tr}(\cdot r_W) = a\overline{w}]}^{(\text{POINT}, W), y} = \tau_{[\text{tr}(g \cdot (y) r_W) = a\overline{w}]}^W \otimes I = E_{a\overline{w}}^{(\text{PAIR}, W), \omega} \otimes I.$$

As a result, due to the consistency of the (PAIR,  $W$ ) measurement, the consistency check is passed with probability 1. On the other hand, if  $\gamma \neq 0$ , then the strategy passes the Magic Square check by Theorem 7.11. As for the consistency check in Item 7, we can write the (POINT,  $X$ ) measurement as follows:

$$E_{[\text{tr}(\cdot r_X) = a\overline{w}]}^{(\text{POINT}, X), y} = \tau_{[\text{tr}(g \cdot (y) r_X) = a\overline{w}]}^X \otimes I = E_{a\overline{w}}^{\text{VARIABLE}_1, \omega} \otimes I,$$

where the last step is by Theorem 7.11. As these measurements are consistent, this test is passed with probability 1.  $\square$

**Soundness of the Pauli basis test.** We state the soundness properties of the Pauli basis test. The following is an adaptation of the self-testing statement in [NW19, Theorem 6.4].

**Theorem 7.14.** *There exists a function*

$$\delta_{\text{QLD}}(\varepsilon, m, d, q) = a(md)^a(\varepsilon^b + q^{-b} + 2^{-bmd})$$

for universal constants  $a \geq 1$  and  $0 < b < 1$  such that the following holds. For all admissible parameter tuples  $\text{qldparams} = (q, m, d)$  and for all strategies  $\mathcal{S} = (|\psi\rangle, A, B)$  for the game  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  that succeed with probability at least  $1 - \varepsilon$ , there exist local isometries  $\phi_A : \mathcal{H}_A \rightarrow \mathcal{H}_{A'} \otimes \mathcal{H}_{A''}$ ,  $\phi_B : \mathcal{H}_B \rightarrow \mathcal{H}_{B'} \otimes \mathcal{H}_{B''}$  (where  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\mathcal{H}_{A''}, \mathcal{H}_{B''} \cong (\mathbb{C}^q)^{\otimes M}$  with  $M = 2^m$ ) and a state  $|\text{AUX}\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$  such that

1.  $\|\phi_A \otimes \phi_B |\psi\rangle - |\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}\| \leq \delta_{\text{QLD}}(\varepsilon, m, d, q)$ ,
2. Letting  $\tilde{A}_a^x = \phi_A A_a^x \phi_A^\dagger$  and  $\tilde{B}_b^y = \phi_B B_b^y \phi_B^\dagger$ , we have for  $W \in \{X, Z\}$

$$\begin{aligned} \tilde{A}_u^{(\text{PAULI}, W)} \otimes I_{B'B''} &\approx_{\delta_{\text{QLD}}} (\tau_u^W)_{A''} \otimes I_{A'B'B''} \\ I_{A'A''} \otimes \tilde{B}_u^{(\text{PAULI}, W)} &\approx_{\delta_{\text{QLD}}} I_{A'A''B'} \otimes (\tau_u^W)_{B''}, \end{aligned}$$

where the  $\approx_{\delta_{\text{QLD}}}$  statement holds with respect to the state  $|\text{AUX}\rangle_{A'B'} \otimes |\text{EPR}_q\rangle_{A''B''}^{\otimes M}$  and the answer summation is over  $u \in \mathbb{F}_q^M$ .

We provide a proof of Theorem 7.14 in Appendix A.

The strategies described in the proof of Lemma 7.13 and in the conclusion of Theorem 7.14 are described in terms of qudit Pauli measurements and maximally entangled states defined over  $q$ -dimensional qudits. However, it is more convenient for our application of the Pauli basis test to instead deal with *qubits*. In particular, we use the Pauli basis test in the “introspection game” of Section 8, where the players are commanded to sample questions according to a sampler  $\mathcal{S}$  of a normal form verifier. By definition of normal form verifier,  $\mathcal{S}$  is a sampler over  $\mathbb{F}_2$ , and therefore it is natural to use a test for qubit Pauli measurements.

Since we use field sizes  $q$  that are powers of 2, Lemma 3.26 shows that the conclusions of Theorem 7.14 can also be described in terms of testing for maximally entangled states over qubits and qubit Pauli measurements.

**Corollary 7.15.** *There exists a function*

$$\delta_{\text{QLD}}(\varepsilon, q, m, d) = a(md)^a(\varepsilon^b + q^{-b} + 2^{-bmd})$$

for universal constants  $a \geq 1$  and  $0 < b < 1$  such that the following holds. For all admissible parameter tuples  $\text{qldparams} = (q, m, d)$  and for all strategies  $\mathcal{S} = (|\psi\rangle, A, B)$  for the game  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  that succeeds with probability at least  $1 - \varepsilon$ , there exist local isometries  $\phi_A : \mathcal{H}_A \rightarrow \mathcal{H}_{A'} \otimes \mathcal{H}_{A''}$ ,  $\phi_B : \mathcal{H}_B \rightarrow \mathcal{H}_{B'} \otimes \mathcal{H}_{B''}$  (where  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\mathcal{H}_{A''}, \mathcal{H}_{B''} \cong (\mathbb{C}^2)^{\otimes M \log q}$  with  $M = 2^m$ ) and a state  $|\text{AUX}\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$  such that

1.  $\|\phi_A \otimes \phi_B |\psi\rangle - |\text{AUX}\rangle \otimes |\text{EPR}_2\rangle^{\otimes M \log q}\| \leq \delta_{\text{QLD}}(\varepsilon, m, d, q)$ ,
2. Letting  $\tilde{A}_a^x = \phi_A A_a^x \phi_A^\dagger$  and  $\tilde{B}_b^y = \phi_B B_b^y \phi_B^\dagger$ , we have for  $W \in \{X, Z\}$

$$\begin{aligned} \tilde{A}_u^{(\text{PAULI}, W)} \otimes I_{B'B''} &\approx_{\delta_{\text{QLD}}} (\sigma_u^W)_{A''} \otimes I_{A'B'B''} \\ I_{A'A''} \otimes \tilde{B}_u^{(\text{PAULI}, W)} &\approx_{\delta_{\text{QLD}}} I_{A'A''B'} \otimes (\sigma_u^W)_{B''} \end{aligned}$$

where the  $\approx_{\delta_{\text{QLD}}}$  statement holds with respect to the state  $|\text{AUX}\rangle_{A'B'} \otimes |\text{EPR}_2\rangle_{A''B''}^{\otimes M \log q}$  and the answer summation is over  $u \in \mathbb{F}_q^M$ . Here,  $\sigma_u^W$  denotes the tensor product of single-qubit Pauli measurements

$$\bigotimes_{i=1}^M \bigotimes_{j=1}^{\log q} \sigma_{u_{ij}}^W$$

where  $\kappa(u_i) = (u_{ij})_{1 \leq j \leq \log q}$  (using the bijection  $\kappa : \mathbb{F}_q \rightarrow \mathbb{F}_2^{\log q}$  from Section 3.3).

*Proof.* Lemma 3.26 implies that there are local isometries that map  $|\text{EPR}_q\rangle^{\otimes M}$  to  $|\text{EPR}_2\rangle^{\otimes M \log q}$ , and map the generalized Pauli measurements  $\tau_u^W$  to  $\bigotimes_{i=1}^M \bigotimes_{j=1}^{\log q} \sigma_{u_{ij}}^W$  where  $\kappa(u_i) = (u_{i1}, \dots, u_{i \log q})$ . Combining this with the isometries guaranteed by Theorem 7.14, we obtain the statement of the Corollary.  $\square$

### 7.3.2 Canonical parameters and complexity of the Pauli basis test

We specify a canonical setting of the parameter tuple  $\text{qldparams}$  as a function of an integer  $R$ . We call this canonical setting  $\text{introparams}$  because we use these in the “introspection” section (Section 8). We then give bounds on the complexity of computing the decision procedure and CL functions of the Pauli basis test corresponding to the parameter tuple  $\text{qldparams}$ , as a function of  $R$ .

**Definition 7.16** (Canonical parameters of the Pauli basis test). Let  $a, b$  be the universal constants specified in Theorem 7.14. Let  $c$  denote the smallest even integer that is at least  $(b + a)/b$ . For all integers  $R \geq 4$ , define the tuple  $\text{introparams}(R) = (q, m, d)$  where

- $q = 2^k$  for  $k = c \lceil \log \log R \rceil + 1$ .
- $m = 2^j$  where  $2^j \leq c \lceil \log R \rceil + 1 < 2^{j+1}$ .
- $d = 1$ .

Intuitively, this choice of parameters is such that the Pauli basis test certifies the presence of an  $M = 2^m$ -qudit EPR state (of local dimension  $q$ ), or equivalently an  $M \log q$ -qubit EPR state.

**Lemma 7.17.** *For all integers  $R \geq 4$ , the parameter tuple  $\text{introparams}(R) = (q, m, d)$  is admissible (see Definition 7.12), satisfies  $2^m \geq R$ , and furthermore there exist universal constants  $a' \geq 1$  and  $0 < b' \leq 1$  such that the function  $\delta_{\text{QLD}}(\varepsilon, m, d, q)$  from Theorem 7.14 is at most*

$$a' (\log(R))^{a'} \cdot \varepsilon^{b'} + \log(R)^{-b'}.$$

*Proof.* We verify the admissibility of  $\text{introparams}(R)$  first: the field size  $q = 2^k$  is admissible because  $k$  is odd. We have  $m|q$  because for all  $R \geq 4$  we have

$$2^j \leq 2c \lceil \log R \rceil \leq 2 \log^c R \leq 2^k.$$

Next, we show that  $R \leq 2^m$ . Since  $m = 2^j \geq (c/2) \log R$ , we have that

$$2^m \geq R^{c/2} \geq R$$

where we used that  $c \geq 2$ .

Finally, we show the desired bound on the error function  $\delta_{\text{QLD}}(\varepsilon, m, d, q)$ . First, observe that  $2^{-bmd} \leq q^{-b}$  because  $k \leq m$ . Note that for  $R \geq 2$ , we have  $m \leq 2c \log R$ , and furthermore  $q = 2^k \geq \log^c R$ . Then, we get

$$\begin{aligned} \delta_{\text{QLD}}(\varepsilon, m, d, q) &= a(md)^a (\varepsilon^b + q^{-b} + 2^{-bmd}) && \text{(by Theorem 7.14)} \\ &\leq a(md)^a (\varepsilon^b + 2q^{-b}) && \text{(because } 2^{-bmd} \leq q^{-b}\text{)} \\ &= a(m)^a (\varepsilon^b + 2q^{-b}) && \text{(because } d = 1\text{)} \\ &\leq a(2c)^a \log^a(R) (\varepsilon^b + 2q^{-b}) && \text{(because } m \leq 2c \log R\text{)} \\ &\leq a(2c)^a \log^a(R) (\varepsilon^b + 2(\log R)^{-cb}) && \text{(because } q \geq \log^c R\text{)} \\ &\leq 2a(2c)^a \log^a(R) (\varepsilon^b + (\log R)^{-cb}) \\ &= 2a(2c)^a ((\log R)^a \varepsilon^b + (\log R)^{a-cb}) \\ &\leq 2a(2c)^a \left( (\log R)^a \varepsilon^b + (\log R)^{-b} \right) && \text{(because } cb - a \geq b \geq 0\text{)} \\ &\leq a' ((\log R))^{a'} \varepsilon^{b'} + (\log R)^{-b}, \end{aligned}$$

where  $a'$  is defined as the quantity  $2a(2c)^a$ . This completes the proof of the lemma.  $\square$

**Lemma 7.18.** *Fix an integer  $R \geq 4$ , let  $\text{introparams}(R) = (q, m, d)$ . The integers  $(q, m, d) = \text{introparams}(R)$  can be computed in time  $\text{polylog } R$  when given the binary representation of  $R$  as input.*

*Proof.* This follows from the definitions of the parameters in Definition 7.16. □

**Lemma 7.19.** *Let  $R \geq 4$  be an integer, and let  $\text{introparams}(R)$  denote the parameter tuple specified by Definition 7.16.*

1. *The time complexity of the decision procedure  $\mathcal{D}^{\text{PAULI}}$  parameterized by  $\text{introparams}(R)$  is  $\text{poly}(R)$ .*
2. *The time complexity of computing marginals of the CL functions  $L_t$  as well as the associated factor spaces, for  $t \in \mathcal{T}^{\text{PAULI}}$ , is  $\text{polylog } R$ .*
3. *The Turing machine description of the decision procedure  $\mathcal{D}^{\text{PAULI}}$  parameterized by  $\text{introparams}$  can be computed from the binary presentation of  $R$  in  $\text{polylog}(R)$  time.*

*Proof.* Finite field arithmetic over  $\mathbb{F}_q$  can be performed in time  $\text{polylog } q$ , by Lemma 3.18. The parameters of  $\text{introparams}(R)$ , which the decision procedure  $\mathcal{D}^{\text{PAULI}}$  implicitly computes given  $R$ , can be computed in time  $\text{polylog}(R)$ , by Lemma 7.18. The most expensive step in  $\mathcal{D}^{\text{PAULI}}$  is to evaluate the low-degree encoding  $g_a(y)$  where  $a \in \mathbb{F}_q^M$  and  $y \in \mathbb{F}_q^m$ , which takes time  $\text{poly}(M, \log q) = \text{poly}(R)$  by Lemma 3.21.

The complexity of computing the CL functions  $L_t$  for types  $t \in \mathcal{T}^{\text{PAULI}}$  is dominated by the complexity of computing the CL functions  $L_{\text{ALINE}}$ ,  $L_{\text{DLINE}}$ , and  $L_{\text{POINT}}$  from the classical low-degree test, which takes time  $\text{poly}(m, \log q) = \text{polylog}(R)$ .

The factor spaces of  $L_t$  depend only on the question type  $t$  (of which there are only constantly many), and outputting the length- $m$  indicator vectors of the factor spaces takes  $O(m) = \text{polylog}(R)$  time.

Finally, the time complexity of computing the description of  $\mathcal{D}^{\text{PAULI}}$  from the binary representation of  $R$  requires  $\text{polylog } R$  time, because the checks performed in the decision procedure  $\mathcal{D}^{\text{PAULI}}$  depend on  $\text{introparams}$  which ultimately depends on  $R$ ; we assume that the decision procedure computes the parameter tuple  $\text{introparams}$  based on  $R$ . Thus the time complexity is dominated by the time to write  $R$  in binary. □

## 8 Introspection Games

### 8.1 Overview

Consider a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  (see Definition 5.33). In this section we design a normal form verifier  $\mathcal{V}^{\text{INTRO}} = (\mathcal{S}^{\text{INTRO}}, \mathcal{D}^{\text{INTRO}})$  (called the *introspective verifier*) such that in the  $n$ -th game  $\mathcal{V}_n^{\text{INTRO}}$  (called the *introspection game*; see Definition 5.33 for the definition of the  $n$ -th game associated with a normal form verifier) the verifier expects the players to sample for themselves questions  $x$  and  $y$  distributed as their questions in the game  $\mathcal{V}_N$  for index  $N = 2^n$ —this is the “introspection” step. Note the exponential separation of the indices of the introspection game versus the original game! Our construction of the introspection game generalizes the introspection technique of [NW19].

Recall the execution of the  $N$ -th game corresponding to the “original” verifier  $\mathcal{V}$  (see Definition 5.33). Let  $n \geq 1$ ,  $N = 2^n$ , and suppose that the CL functions of  $\mathcal{S}$  on index  $N$  are  $L^A, L^B$  acting on an ambient space  $V$ . In the game  $\mathcal{V}_N$ , the verifier first samples  $z \in V$  uniformly at random and gives each player  $w \in \{A, B\}$  the question  $x_w = L^w(z)$ . In this context, the string  $z$  is referred to as the “seed”. The players respond with answers  $a_A$  and  $a_B$ , respectively, and the verifier accepts or rejects according to the output of  $\mathcal{D}(N, x_A, x_B, a_A, a_B)$ .

In the introspection game, with some constant probability independent of  $n$ , the verifier  $\mathcal{V}_n^{\text{INTRO}}$  sends the question pair (INTRO, A) to player  $w$  and (INTRO, B) to the other player  $\bar{w}$ , where  $w \in \{A, B\}$  and recall that  $\bar{w} = B$  if  $w = A$  and  $\bar{w} = A$  otherwise. The verifier expects player  $w$  to measure their share of the state  $|\text{EPR}\rangle_V$  using a coarse-grained  $Z$ -basis measurement whose outcomes range over  $L^A(V)$ , and similarly player  $\bar{w}$  measures the state  $|\text{EPR}\rangle_V$  using a coarse-grained  $Z$ -basis measurement with outcomes that range over  $L^B(V)$ . If the players perform these measurements honestly, then the outcomes  $(x_A, x_B)$  are distributed exactly according to  $\mu_{\mathcal{S}, N}$ , the question distribution of the game  $\mathcal{V}_N$ . Players  $w$  and  $\bar{w}$  are then expected to respond with the question  $x_w$  and  $x_{\bar{w}}$  that they each sampled, together with strings  $a_w$  and  $a_{\bar{w}}$ , respectively, which are intended to be the answers for the question pair  $(x_A, x_B)$  in the game  $\mathcal{V}_N$ . In other words, the players introspectively sample the question pair  $(x_A, x_B)$  and then respond with the question itself and an answer for it.

To facilitate comprehension, we call the players that interact with the introspective verifier  $\mathcal{V}^{\text{INTRO}}$  the “introspecting players”, and the players that interact with the “original” verifier  $\mathcal{V}$  the “original players”. To ensure that the introspecting players follow the above procedure honestly, the introspective verifier  $\mathcal{V}_n^{\text{INTRO}}$  first uses the (binary) Pauli basis test described in Section 7.3 to force the introspecting players to share the state  $|\text{EPR}\rangle_V$ . The Pauli basis test also ensures that the players measure  $\sigma^W$  and report the outcome honestly when they receive questions (PAULI,  $W$ ) for  $W \in \{X, Z\}$ . For  $v \in \{A, B\}$  the verifier cross-checks the question pairs (INTRO,  $v$ ) and (PAULI,  $Z$ ) to enforce that the honest measurement is performed for question (INTRO,  $v$ ).

The main difficulty in the soundness analysis is to ensure that the answer of player  $w$  who received question (INTRO,  $v$ ) depends only on  $L^v(z)$  and not on any other information about the string  $z$ . First assume for simplicity that  $L^v$  is a linear function. As shown below (based on Lemma 8.4),  $L^v(z)$  can be obtained by measuring a specific collection of  $\sigma^Z$  observables; namely, the set

$$\{\sigma^Z(u) : u \in \ker(L^v)^\perp\}. \quad (60)$$

To prevent the player from obtaining any additional information the verifier needs to enforce that the player does *not* additionally measure any  $\sigma^Z(u)$  for  $u \notin \ker(L^v)^\perp$ . (We refer to any such  $\sigma^Z$  as a “prohibited”  $\sigma^Z$ .)



The introspective verifier achieves this by sometimes sending the READ type question  $(\text{READ}, v)$  or  $\text{HIDE}_k$  type questions  $(\text{HIDE}_k, v)$  to the players. When receiving the READ question, the players are required to also measure observables from the set

$$\{\sigma^X(r) : r \in \ker(L^v)\}, \quad (61)$$

which (as shown in Lemma 8.5 below) commute with every Z-basis measurement in (60). On the other hand, any prohibited  $\sigma^Z(u)$  must anticommute with at least one of the  $\sigma^X(r)$ , as otherwise  $u$  would be in  $\ker(L^v)^\perp$ . As a result, honestly measuring the  $\sigma^X$  observables of (61) has the effect of preventing the player from measuring any of the prohibited  $\sigma^Z$  observables, so that the answer  $a$  can effectively only depend on the question  $L^v(z)$ . (In the protocol, the verifier asks the player to measure the function  $(L^v)^\perp(z)$  in the X basis, rather than all of the  $\sigma^X$  observables. By Lemma 8.4 the two are equivalent.) Similarly to how questions  $(\text{PAULI}, Z)$  and  $(\text{INTRO}, v)$  are cross-checked, the questions  $(\text{PAULI}, X)$ ,  $(\text{HIDE}_k, v)$  and  $(\text{READ}, v)$  are cross-checked in order to ensure that the honest X-basis measurements are performed for the  $\text{HIDE}_k$  questions and the READ questions.

When the CL functions  $L^v$  are  $\ell$ -level for  $\ell > 1$ , the introspective verifier sends one of  $O(\ell)$  different hiding questions to the players, chosen at random; together these hiding checks ensure that each of the constituent linear maps of  $L^v$  are honestly measured. Intuitively, these hiding questions “interpolate” between questions  $(\text{PAULI}, Z)$ ,  $(\text{INTRO}, v)$  and  $(\text{PAULI}, X)$  in a way that, for all pairs of questions asked by the verifier, the honest measurements commute. (See Figure 11 for an overview of the honest measurements.) This strong commuting property of the strategy is essential for the oracularization transformation in Section 9 to be possible.

A key property of the introspection game is that the distribution of questions (which include the Pauli basis test questions as well as the introspection questions and the hiding questions) is also conditionally linear. This means that the introspection game can be ultimately specified by a normal form verifier  $\mathcal{V}^{\text{INTRO}} = (\mathcal{S}^{\text{INTRO}}, \mathcal{D}^{\text{INTRO}})$ , which is crucial for recursive compression of games. Moreover, while the time complexity of the introspective verifier’s decider  $\mathcal{D}^{\text{INTRO}}$  remains polynomially related to that of  $\mathcal{D}$  on index  $N$ , the time complexity of the sampler  $\mathcal{S}^{\text{INTRO}}$  is polylog( $N$ ) (exponentially smaller), due to the efficiency of the Pauli basis test. Finally,  $\mathcal{S}^{\text{INTRO}}$  only depends on  $\mathcal{V}$  through the number of levels  $\ell$  of  $\mathcal{S}$  as well as upper bounds on the time complexities of  $\mathcal{S}$  and  $\mathcal{D}$ .

## 8.2 The introspective verifier

Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be an  $\ell$ -level normal form verifier. We call  $\mathcal{V}$ ,  $\mathcal{S}$ , and  $\mathcal{D}$  the “original” verifier, sampler, and decider, respectively. For all original verifier  $\mathcal{V}$ , we define the introspective verifier  $\mathcal{V}^{\text{INTRO}}$  in this section. We use  $N = 2^n$  to denote the index of the original verifier  $\mathcal{V}$  that is simulated by the introspective verifier  $\mathcal{V}^{\text{INTRO}}$  on index  $n$ .

The *introspective verifier corresponding to  $\mathcal{V}$  and parameters  $(\lambda, \ell)$*  is a *typed* normal form verifier  $\hat{\mathcal{V}}^{\text{INTRO}} = (\hat{\mathcal{S}}^{\text{INTRO}}, \hat{\mathcal{D}}^{\text{INTRO}})$ , sketched in Section 8.1 and specified in detail in the present section (see Section 6 for the definition of typed normal form verifiers). In the following descriptions of the sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  and decider  $\hat{\mathcal{D}}^{\text{INTRO}}$ , all parameters are functions of the index  $n$ , the number of levels  $\ell$  of the sampler  $\mathcal{S}$ , and the parameter  $\lambda$ ; we often leave this dependence implicit.

Let  $R = N^\lambda$ , and let  $\text{introparams}(R) = (q, m, d)$  denote the parameter tuple specified in Section 7.3.2. Note that  $\text{introparams}$  is implicitly a function of  $n$  (since  $R$  is a function of  $n$ ). The associated parameter tuple  $\text{introparams}$  is intended to parametrize a Pauli basis test that certifies the presence of  $Q = 2^m \log q$ -qubit EPR states; intuitively, these qubits are meant to serve as the source of randomness for the CL functions

of the original sampler  $\mathcal{S}$ . Since we will assume that the original verifier  $\mathcal{V}$  is  $\lambda$ -bounded, the original sampler  $\mathcal{S}$  on index  $N$  has time complexity at most  $R$ , and therefore the amount of randomness needed is at most  $R$ . By Lemma 7.17, the number of qubits  $Q$  in the EPR states is at least  $R$ , and thus the EPR qubits can be used to properly introspect the original game.

Recall that the players in the introspection game are referred to as “introspecting players” and the players in the original game are referred to as “original players”. We use the following notation in order to distinguish between questions and answers meant for the introspecting players versus the original players. The questions and answers of the introspecting players are denoted by hatted variables (e.g.,  $\hat{x}$  and  $\hat{a}$ ). Similarly, the associated question types are denoted using hatted variables  $\hat{f}$ . The questions and answers of the original players in the original game  $\mathcal{V}_N$  are denoted using non-hatted variables (e.g.  $x$ ,  $a$ , and so on).

**Types and type graph.** The type set  $\mathcal{T}^{\text{INTRO}}$  for the introspective verifier  $\hat{\mathcal{V}}^{\text{INTRO}}$  is

$$\mathcal{T}^{\text{INTRO}} = \mathcal{T}^{\text{PAULI}} \cup \left( \left( \{\text{INTRO, SAMPLE, READ}\} \cup \left( \bigcup_{k=1}^{\ell} \{\text{HIDE}_k\} \right) \right) \times \{A, B\} \right),$$

where  $\mathcal{T}^{\text{PAULI}}$  is the type set of the Pauli basis test, defined in Section 7.3. The type graph  $G^{\text{INTRO}}$  is specified in Figure 9.




Figure 9: Type graph  $G^{\text{INTRO}}$  for the introspection game. Each vertex also has a self-loop which is not drawn in the figure for clarity.

**Sampler.** We first define a 3-level  $(\mathcal{T}^{\text{INTRO}}, G^{\text{INTRO}})$ -typed sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$ , which has field size  $q(n)$  and dimension  $3m(n) + 3$ , where  $q(n)$  and  $m(n)$  are specified by  $\text{introparams}(R)$ . Note that the dimension matches that of the space  $V^{\text{PAULI}}$  of the CL functions of the Pauli basis test parameterized by  $\text{introparams}(R)$ , specified in Section 7.3.1.

Fix  $n \in \mathbb{N}$ . We specify the CL functions of  $\hat{\mathcal{S}}^{\text{INTRO}}$  on index  $n$ . Since the functions  $L_{\hat{t}}^{A,n}$  and  $L_{\hat{t}}^{B,n}$  are identical for all  $n$  and  $\hat{t}$ , we omit the superscripts A and B. For types  $\hat{t} \in \mathcal{T}^{\text{PAULI}}$ , the CL functions  $L_{\hat{t}}^n$  are given by those specified in Section 7.3.1, parameterized by  $\text{introparams}(R)$ . For types  $\hat{t} \in \mathcal{T}^{\text{INTRO}} \setminus \mathcal{T}^{\text{PAULI}}$ , the associated CL functions are defined to be 0-level CL functions (i.e., they are the 0 map). This means that for question types such as  $\hat{t} = (\text{INTRO}, v)$  or  $\hat{t} = (\text{SAMPLE}, v)$  for  $v \in \{A, B\}$  the associated question is solely comprised of the question type label.

Finally, we define the typed sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  to be  $\kappa(\hat{\mathcal{S}}^{\text{INTRO}})$ , the downsized sampler (Definition 6.6) corresponding to  $\hat{\mathcal{S}}^{\text{INTRO}}$ . The typed CL functions associated with the Pauli basis test are 3-level; using Remark 4.4 and Lemma 4.11 it follows that  $\hat{\mathcal{S}}^{\text{INTRO}}$  is a 3-level typed sampler.

The following lemma establishes the complexity of the sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  as well as the complexity of computing a description of it from the parameters  $(\lambda, \ell)$ .

**Lemma 8.1.** *There exists a 2-input Turing machine `ComputeIntroSampler` that on input  $(\lambda, \ell)$  outputs a description of the sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  (corresponding to parameters  $\lambda, \ell$ ) in time  $\text{polylog}(\lambda, \ell)$ . Furthermore,*

1.  $\text{TIME}_{\hat{\mathcal{S}}^{\text{INTRO}}}(n) = \text{poly}(n, \lambda, \ell)$ , and
2.  $\hat{\mathcal{S}}^{\text{INTRO}}$  is a 3-level typed sampler.

*Proof.* Define the following 9-input Turing machine `RawIntroSampler`, that does not depend on any parameters (so that its description length is constant). On input  $(\lambda', \ell', n', x'_1, \dots, x'_6)$ , the Turing machine `RawIntroSampler` computes the output of the typed sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  (parameterized by  $(\lambda, \ell)$ ) with input tapes set to  $(n', x'_1, \dots, x'_6)$ . In more detail, the Turing machine `RawIntroSampler` first computes  $\text{introparams}(R)$  for  $R = 2^{\lambda'n'}$ . Using Lemma 7.18, this computation takes time  $\text{polylog}(R)$ . Next, depending on the contents of the last 7 input tapes of `RawIntroSampler`, the Turing machine evaluates the dimension of  $\hat{\mathcal{S}}^{\text{INTRO}}$  (which can easily be computed from  $\text{introparams}(R)$ ), or one of the CL functions, or returns a representation of a factor space of  $\hat{\mathcal{S}}^{\text{INTRO}}$ . If the type passed as input is  $\hat{t} \in \mathcal{T}^{\text{PAULI}}$  then by Lemma 7.19 this takes time  $\text{polylog}(R)$ . If  $\hat{t} \in \mathcal{T}^{\text{INTRO}} \setminus \mathcal{T}^{\text{PAULI}}$  then this can be done in  $O(\log \ell')$  time (to read the input type). Overall, `RawIntroSampler` runs in time  $\text{polylog}(R, \ell')$ .

We now define the Turing machine `ComputeIntroSampler` as follows. On input  $(\lambda, \ell)$ , `ComputeIntroSampler` outputs the description of `RawIntroSampler` with the first two input tapes hardwired to  $\lambda$  and  $\ell$ , respectively, yielding the sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  corresponding to parameters  $(\lambda, \ell)$ . Computing this description takes  $O(\log \lambda + \log \ell)$  time.

The time complexity of  $\hat{\mathcal{S}}^{\text{INTRO}}$  follows from the time complexity of `RawIntroSampler` and the number of levels is by construction.  $\square$

**Decider.** The typed decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  is specified in Figure 10. We explain how to interpret the figure, including the notation. (It may also be helpful to review the description of the intended strategy for the players in the game, described in Section 8.3.)

*Question and answer format.* The decider takes as input a tuple  $(n, \hat{t}_A, \hat{x}_A, \hat{t}_B, \hat{x}_B, \hat{a}_A, \hat{a}_B)$  where  $(\hat{t}_w, \hat{x}_w)$  denotes the question for introspecting player  $w \in \{A, B\}$ , and  $\hat{a}_w$  denotes their answer. As in the specification of the Pauli basis test, in Figure 10 we include an “answer key” indicating how the players’ answers are parsed, depending on the question type. When the question type is from  $\mathcal{T}^{\text{PAULI}}$ , the question and answer format are as described in Figure 8. When the question type is in  $\mathcal{T}^{\text{INTRO}} \setminus \mathcal{T}^{\text{PAULI}}$ , the answer format is described in the table at the top of Figure 10.<sup>30</sup> For each such question, there is an associated variable

<sup>30</sup>There is no question format specification for the question types in  $\mathcal{T}^{\text{INTRO}} \setminus \mathcal{T}^{\text{PAULI}}$ , because the question is solely comprised of the question type label.

Type	Answer format
(INTRO, $v$ )	$(y, a) \in V \times \{0, 1\}^*$
(SAMPLE, $v$ )	$(z, a) \in V \times \{0, 1\}^*$
(READ, $v$ )	$(y, y^\perp, a) \in V \times V \times \{0, 1\}^*$
(HIDE $_k$ , $v$ )	$(y, y^\perp, x) \in V \times V \times V$

In the following, whenever  $\mathcal{S}$  or  $\mathcal{D}$  is called as a subroutine,  $\hat{\mathcal{D}}^{\text{INTRO}}$  aborts and rejects if the subroutine takes more than  $N^\lambda$  time steps. On input  $(n, \hat{t}_A, \hat{x}_A, \hat{t}_B, \hat{x}_B, \hat{a}_A, \hat{a}_B)$ , the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  first computes the dimension  $s(N)$  of  $V$  by calling the original sampler  $\mathcal{S}$  on input  $(N, \text{DIMENSION})$ . The decider rejects if  $s(N) > N^\lambda$ , or if  $\max\{|\hat{a}_A|, |\hat{a}_B|\} \geq 3 \cdot 2^m \cdot \log q$  where  $\text{introparams}(N^\lambda) = (q, m, d)$ . Otherwise, it performs the following tests for all  $w, v \in \{A, B\}$ . (If no test applies, the decider accepts.)

1. **(Pauli test).** If  $\hat{t}_A, \hat{t}_B \in \mathcal{T}^{\text{PAULI}}$ , the decider accepts if  $\mathcal{D}_{\text{introparams}}^{\text{PAULI}}$  accepts the input  $(\hat{t}_A, \hat{x}_A, \hat{t}_B, \hat{x}_B, \hat{a}_A, \hat{a}_B)$ .
2. **(Sampling tests).**
  - (a) If  $\hat{t}_w = (\text{PAULI}, Z)$  and  $\hat{t}_{\bar{w}} = (\text{SAMPLE}, v)$ , accept if  $\hat{a}_w^V = z_{\bar{w}}$ .
  - (b) If  $\hat{t}_w = (\text{INTRO}, v)$ ,  $\hat{t}_{\bar{w}} = (\text{SAMPLE}, v)$ , accept if  $y_w = L^v(z_{\bar{w}})$  and  $a_w = a_{\bar{w}}$ .
3. **(Hiding tests).**
  - (a) If  $\hat{t}_w = (\text{INTRO}, v)$ ,  $\hat{t}_{\bar{w}} = (\text{READ}, v)$ , accept if  $y_w = y_{\bar{w}}$  and  $a_w = a_{\bar{w}}$ .
  - (b) If  $\hat{t}_w = (\text{HIDE}_\ell, v)$ ,  $\hat{t}_{\bar{w}} = (\text{READ}, v)$ , accept if  $y_{w, < \ell} = y_{\bar{w}, < \ell}$ , and  $y_w^\perp = y_{\bar{w}}^\perp$ .
  - (c) If  $\hat{t}_w = (\text{HIDE}_k, v)$ ,  $\hat{t}_{\bar{w}} = (\text{HIDE}_{k+1}, v)$  for some  $k \in \{1, 2, \dots, \ell - 1\}$ , accept if
$$y_{w, < k} = y_{\bar{w}, < k}, \quad y_{w, \leq k}^\perp = y_{\bar{w}, \leq k}^\perp, \quad x_{w, > k+1} = x_{\bar{w}, > k+1},$$
and  $y_{\bar{w}, k+1}^\perp = (L_{k+1, u}^v)^\perp(x_{w, k+1})$  where  $u = y_{\bar{w}, \leq k}$ .
  - (d) If  $\hat{t}_w = (\text{PAULI}, X)$ ,  $\hat{t}_{\bar{w}} = (\text{HIDE}_1, v)$ , accept if  $y_{\bar{w}, 1}^\perp = (L_1^v)^\perp(\hat{a}_w^{V_1})$  and  $\hat{a}_w^{V_{>1}} = x_{\bar{w}, > 1}$ .
4. **(Game test).** If  $\hat{t}_w = (\text{INTRO}, A)$  and  $\hat{t}_{\bar{w}} = (\text{INTRO}, B)$ , accept if  $\mathcal{D}$  accepts  $(N, y_w, y_{\bar{w}}, a_w, a_{\bar{w}})$ .
5. **(Consistency test).** If  $\hat{t}_A = \hat{t}_B$ , accept if and only if  $\hat{a}_A = \hat{a}_B$ .

Figure 10: The typed decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  for the introspective verifier, parameterized by integers  $\lambda, \ell$ , on index  $n$ .  $N$  denotes  $2^n$ ,  $V$  is the ambient space for  $\mathcal{S}$ , and  $\{L^v\}_{v \in \{A, B\}}$  the associated CL functions on index  $N$ .

$v \in \{A, B\}$  that indicates to the introspecting player which original player it is supposed to impersonate in the introspection game.

In the figure,  $V$  denotes the ambient space of the original sampler  $\mathcal{S}$  on index  $N = 2^n$ . We assume that the original verifier  $\mathcal{V}$  is  $\lambda$ -bounded and in particular the running time of the original sampler  $\mathcal{S}$  (and therefore its dimension) is at most  $R = N^\lambda$ . If the dimension were larger than  $N^\lambda$ , the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  would always reject in the beginning.

Since  $V$  is isomorphic to  $\mathbb{F}_2^{s(N)}$ , where  $s(N)$  is the dimension of  $\mathcal{S}$ , the space  $V$  is identified in a canonical way as the register subspace of  $\mathbb{F}_2^Q$  spanned by  $e_1, \dots, e_{s(N)}$  where  $e_i$  is the  $i$ -th elementary basis vector. For example, if  $\hat{t}_w = (\text{READ}, v)$ , then syntactically the player's answer is a triple  $(y, y^\perp, a)$  in  $\mathbb{F}_2^Q \times \mathbb{F}_2^Q \times \{0, 1\}^*$ . We assume that the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  computes the dimension  $s(N)$  of the subspace  $V$  by calling  $\mathcal{S}$  on input  $(N, \text{DIMENSION})$ , and if  $y, y^\perp$  are not presented as vectors in the subspace  $V$ , then the decider rejects. Thus in the analysis we directly consider  $y, y^\perp$  as vectors in  $V$ . In Figure 10, the components of the players' answers are subscripted by the player index. For example, if player  $w$  receives question  $(\text{HIDE}_k, v)$ , then their answers are denoted by  $(y_w, y_w^\perp, x_w)$ .

The notation used in the “answer key” is meant to give an indication of the intended meaning of the players' answers. We use  $y$  to denote a vector that is supposed to be the result of measuring a CL function  $L^v$ ;  $y^\perp$  is supposed to be the result of measuring “dual” linear maps  $L^\perp$  (as used in Step 3 in Figure 10);  $x$  is supposed to be the result of  $\sigma^X$  measurements, and  $z$  is supposed to be the result of  $\sigma^Z$  measurements. We use  $a$  to denote the introspected answers meant for the original decider  $\mathcal{D}$ .

*CL functions and factor spaces.* For  $v \in \{A, B\}$ , let  $L^v = L^{v, N}$  denote the CL function for original player  $v$  specified by  $\mathcal{S}$  on index  $N = 2^n$ . For  $z \in V$ , the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  computes  $L^v(z)$  by calling  $\mathcal{S}$  on input  $(N, v, \text{MARGINAL}, \ell, z)$ .

For  $y \in V$  and  $k \in \{1, \dots, \ell\}$  we define register subspaces  $V_k^v(y)$  by induction on  $k$ . For  $k = 1$ ,  $V_1^v(y)$  is the first factor space<sup>31</sup> of  $L^v$  and is independent of  $y$ . Suppose  $V_j^v(y)$  has been defined for all  $j < k$ . Then we define the marginal space  $V_{<k}^v(y) = \bigoplus_{j=1}^{k-1} V_j^v(y)$ , and define  $V_k^v(y)$  as the  $k$ -th factor space  $V_{k,u}^v$  of  $L^v$  with prefix  $u = L_{<k}^v(y)$ . We also define  $V_{\leq k}^v(y) = V_{<k+1}^v(y)$ , and  $V_{>k}^v(y)$  to be the complementary register subspace to  $V_{\leq k}^v(y)$  within  $V$ .

The decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  computes factor spaces  $V_j^v(y)$  from  $y \in V$  in the following sequential manner: first, the indicator vector for the factor space  $V_1^v$  is computed by calling the original sampler  $\mathcal{S}$  on input  $(N, v, \text{FACTOR}, 1, 0)$ . Let  $y_1$  denote  $L_1^v(y)$ . Then, for  $j \in \{2, \dots, \ell\}$ , the factor space  $V_j^v(y)$  is computed by calling  $\mathcal{S}$  on input  $(N, v, \text{FACTOR}, j, y_{\leq j-1})$ , where  $y_{\leq j-1} = L_{\leq j-1}^v(y)$ .

Finally, Lemma 4.6 implies that the CL function  $L^v$  gives rise to functions  $\{L_{k,u}^v\}$  where for all  $k$  and prefixes  $u$ ,  $L_{k,u}^v$  is a map from  $V_{k,u}^v$  to  $V_{k,u}^v$ .

*Detailed explanation of the steps of  $\hat{\mathcal{D}}^{\text{INTRO}}$ .* We give more details on the implementation of decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  specified in Figure 10.

1. The decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  first checks that the answers are not too long; the maximum-length answer should be a tuple  $(y, y^\perp, a)$  in response to question  $(\text{READ}, v)$  where  $y, y^\perp \in V$  and  $a$  is an answer intended for the original decider  $\mathcal{D}$  on index  $N$  (which we assume runs in time at most  $N^\lambda$ ), or a tuple  $(y, y^\perp, x) \in V \times V \times V$  in response to question  $(\text{HIDE}_k, v)$ . Either way, the maximum answer length should be  $3Q = 3 \cdot 2^m \cdot \log q = \text{poly}(R)$  bits long. This check is necessary in order to ensure that the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  halts in time that is a fixed polynomial of  $R = N^\lambda$ .

<sup>31</sup>See Lemma 4.6 for the definitions of factor spaces of a CL function.

2. If the question types for the players are both in  $\mathcal{T}^{\text{PAULI}}$ ,  $\hat{\mathcal{D}}^{\text{INTRO}}$  executes the decision procedure  $\mathcal{D}^{\text{PAULI}}$  for the Pauli basis test parametrized by  $\text{introparams}(R) = (q, m, d)$  (see Section 7.3.1 for the definition of  $\mathcal{D}^{\text{PAULI}}$ ). The Pauli basis test is intended to ensure that the players share at least  $Q \geq R$  EPR pairs, where  $Q = 2^m \log q$ . This number of EPR pairs is chosen so that, under the assumption that the original verifier  $\mathcal{V}$  is  $\lambda$ -bounded (see Definition 5.32), the time complexity of  $\mathcal{S}$  on index  $N$  (and therefore its dimension) is at most  $R = N^\lambda$ . When analyzing the Completeness, Soundness, and Entanglement properties of the introspective verifier  $\mathcal{V}^{\text{INTRO}}$ , we assume that the original verifier  $\mathcal{V}$  is  $\lambda$ -bounded (see Theorem 8.3).
3. In Step 2a of  $\hat{\mathcal{D}}^{\text{INTRO}}$ , player  $w \in \{A, B\}$  receives question (PAULI,  $Z$ ) and player  $\bar{w}$  receives question (SAMPLE,  $v$ ), for some  $v \in \{A, B\}$ . According to the answer key, player  $w$  is expected to return an answer  $\hat{a}_w \in \mathbb{F}_2^Q$  and player  $\bar{w}$  is expected to respond with a pair  $(y_{\bar{w}}, a_{\bar{w}}) \in V \times \{0, 1\}^*$ . Thus, the dimension of answer  $\hat{a}_w$  may be larger than that of  $y_{\bar{w}}$ ; this is the reason that Step 2a checks consistency between  $y_{\bar{w}}$  and the projection of  $\hat{a}_w$  to  $V$ .
4. In Step 2b of  $\hat{\mathcal{D}}^{\text{INTRO}}$ , player  $w \in \{A, B\}$  receives question (INTRO,  $v$ ) and player  $\bar{w}$  receives question (SAMPLE,  $v$ ). As specified by the “answer key”, player  $w$  responds with  $(y_w, a_w) \in V \times \{0, 1\}^*$  and player  $\bar{w}$  responds with  $(z_{\bar{w}}, a_{\bar{w}}) \in V \times \{0, 1\}^*$ . The decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  checks that  $a_w = a_{\bar{w}}$  and  $y_w = L^v(z_{\bar{w}})$  where  $L^v$  denotes the CL function for player  $v$  specified by  $\mathcal{S}$  for index  $N = 2^n$ . The CL function is computed by calling  $\mathcal{S}$  on input  $(N, v, \text{MARGINAL}, \ell, z)$ .
5. In Step 3b,  $\hat{\mathcal{D}}^{\text{INTRO}}$  checks that the answer of player  $w$  who received (HIDE $_{\ell}$ ,  $v$ ) is consistent with the answer of player  $\bar{w}$  who received (READ,  $v$ ). One of the checks is that  $y_{w, < \ell} = y_{\bar{w}, < \ell}$ ; these are, respectively,  $L^v_{< \ell}(y_w)$  and  $L^v_{< \ell}(y_{\bar{w}})$ .
6. In Steps 3c, the vectors  $x_{w, > k+1}$  and  $x_{\bar{w}, > k+1}$  denote the projections of  $x_w$  and  $x_{\bar{w}}$  to the factor spaces  $V^v_{> k+1}(y_w)$  and  $V^v_{> k+1}(y_{\bar{w}})$ , respectively. Similarly,  $y_{\bar{w}, k+1}^\perp$  denotes the projection of  $y_{\bar{w}}^\perp$  to the factor space  $V^v_{k+1}(y_{\bar{w}})$  and  $x_{w, k+1}$  denotes the projection of  $x_w$  to the factor space  $V^v_{k+1}(y_{\bar{w}})$ . We emphasize that the latter factor space depends on  $y_{\bar{w}}$  and not  $y_w$ .<sup>32</sup>

The decider also has to compute  $(L^v_{k+1, y_{\bar{w}, \leq k}})^\perp(x_{w, k+1})$ . According to Definition 3.11, this requires specifying a basis for  $\ker(L^v_{k+1, y_{\bar{w}, \leq k}})^\perp$ . To compute the value, the decider performs the following steps:

- (a) Call  $\mathcal{S}$  on input  $(N, v, \text{FACTOR}, k+1, y_{\bar{w}, \leq k})$  to obtain a subset  $H = \{h_1, \dots, h_m\}$  of the canonical basis for  $\mathbb{F}_2^{s(N)}$  that is a basis of the register subspace  $V^v_{k+1}(y_{\bar{w}})$ .
- (b) For  $i \in \{1, 2, \dots, m\}$  compute  $c_i = \mathcal{S}(N, v, \text{LINEAR}, k+1, y_{\bar{w}, \leq k}, h_i)$ . Compute a matrix representation  $M$  for  $L^v_{k+1, y_{\bar{w}, \leq k}}$  in the basis  $H$ , whose columns are the vectors  $c_1, \dots, c_m$  as elements of  $V^v_{k+1}(y_{\bar{w}})$ .
- (c) Using a canonical algorithm for Gaussian elimination, compute a basis  $F$  for  $\ker(M)$ .
- (d) Compute the canonical complement  $S$  of  $F$ , as in Definition 3.6.  $S$  is a basis for  $\ker(L^v_{k+1, y_{\bar{w}, \leq k}})^\perp$ .
- (e) To compute  $(L^v_{k+1, y_{\bar{w}, \leq k}})^\perp$  on input  $x_{w, k+1}$ , compute the canonical linear map with kernel basis  $S$  (see Definition 3.10) on input  $x_{w, k+1}$ .

---

<sup>32</sup>This is because it is the  $\bar{w}$  player who receives the (HIDE $_{k+1}$ ,  $v$ ) question, and in the “honest” strategy (described in Section 8.3) they are supposed to measure  $k$  registers to obtain a sequence of values  $(y_{\bar{w}, 1}, y_{\bar{w}, 2}, \dots, y_{\bar{w}, k})$  which specify the  $(k+1)$ -st factor subspace, whereas the  $w$  player is only supposed to measure  $k-1$  registers.

7. In Step 3d, the player  $w$  that receives  $(\text{PAULI}, X)$  is expected to return an answer  $\hat{a}_w$  in  $\mathbb{F}_2^Q$ . Part of this step checks that the projection of  $\hat{a}_w$  to  $V_{>1}$  is equal to  $x_{\bar{w}, >1}$  (which is the projection to  $V_{>1}$  of the third component of the answer triple of player  $\bar{w}$  that receives question  $(\text{HIDE}_1, v)$ ).

### 8.2.1 Complexity of the introspective verifier

In this section we determine the complexity of the introspective verifier. The following lemma establishes the complexity of the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  as well as the complexity of computing a description of it from the parameters  $(\lambda, \ell)$  and the description of the original verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ .

**Lemma 8.2.** *There exists a polynomial time Turing machine `ComputeIntroDecider` that on input  $(\mathcal{V}, \lambda, \ell)$  outputs a description of a Turing machine where:*

1. *Its description length is at most  $\text{poly}(\lambda, \ell)$ .*
2. *Its time complexity is at most  $\text{TIME}_{\hat{\mathcal{D}}^{\text{INTRO}}}(n) = \text{poly}(2^{\lambda n}, \ell)$ .*

*Furthermore, if  $|\mathcal{V}| \leq \lambda$ , then the output of `ComputeIntroDecider` is the introspective decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  corresponding to verifier  $\mathcal{V}$  and parameters  $(\lambda, \ell)$ .*

*Proof.* Define the following 10-input Turing machine `RawIntroDecider`, which is a universal Turing machine that computes the same function as the introspection decider  $\hat{\mathcal{D}}^{\text{INTRO}}$ , except it also takes as input extra inputs to specify the sampler, decider, and the parameters  $\lambda, \ell$  that are used in  $\hat{\mathcal{D}}^{\text{INTRO}}$ . Since `RawIntroDecider` is a universal Turing machine, its description length is constant. More formally, on input

$$(\mathcal{V}', \lambda', \ell', n', x'_1, x'_2, \dots, x'_6),$$

the Turing machine `RawIntroDecider` computes the output of the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  corresponding to  $\mathcal{V}', \lambda', \ell'$  with input tapes set to  $(n', x'_1, \dots, x'_6)$ . In more detail, the Turing machine  $\hat{\mathcal{D}}^{\text{INTRO}}$  first computes  $\text{introparams}(R)$  for  $R = 2^{\lambda' n'}$ . Using Lemma 7.18, this computation takes time  $\text{polylog}(R)$ . It then executes the tests described in Figure 10. Write  $\mathcal{V}' = (\mathcal{S}', \mathcal{D}')$ . The complexity of performing the entire procedure is subsumed by the complexity of the following steps:

1. Running the decider  $\mathcal{D}^{\text{PAULI}}$ , which takes time  $\text{poly}(R)$  by Lemma 7.19;
2. Running the decider  $\mathcal{D}'$  (on index  $N' = 2^{n'}$ ) for at most  $2^{\lambda' n'}$  steps;
3. Running the sampler  $\mathcal{S}'$  (on index  $N'$ ) in order to compute the dimension  $s(N')$  and the marginal and factor spaces, and the CL functions as described in Section 8.2.  $\mathcal{S}'$  is called at most  $\text{poly}(s(N'), \ell')$  times; due to the timeout, each call takes time at most  $2^{\lambda' n'}$ .
4. Computing  $(L_{k,u}^v)^\perp(x_{\bar{w},k})$  in Step 3c. This only requires to perform Gaussian elimination and other simple finite field manipulations that can be implemented in  $\text{poly}(s(N'))$  time.

All other tests are elementary. Thus the time complexity of `RawIntroDecider` is  $\text{poly}(R, \ell')$ . Note that the bound is independent of  $\mathcal{V}'$ : due to the abort condition in the definition of  $\hat{\mathcal{D}}^{\text{INTRO}}$ , the Turing machine  $\hat{\mathcal{D}}^{\text{INTRO}}$  aborts if the runtime of  $\mathcal{S}'$  or  $\mathcal{D}'$  is larger than  $2^{\lambda' n'}$ .

We now define the Turing machine `ComputeIntroDecider`: on input  $(\mathcal{V}, \lambda, \ell)$ , it first determines whether the description length  $|\mathcal{V}|$  of the verifier is at most  $\lambda$ .<sup>33</sup> If it isn't, then it sets  $\mathcal{V}' = (\mathcal{S}', \mathcal{D}')$  where  $\mathcal{S}', \mathcal{D}'$

<sup>33</sup>This can be done by keeping a counter of  $O(\log \lambda)$  bits, and incrementing it as the input  $\mathcal{V}$  is being scanned through. If at any point the counter exceeds  $\lambda$ , the scanning stops.

are the trivial Turing machine whose description is 0. Otherwise, `ComputeIntroDecider` sets  $\mathcal{V}' = \mathcal{V}$ . Note that by construction, the description length of  $\mathcal{V}'$  is at most  $\lambda$ , and that the time complexity of computing the description of  $\mathcal{V}'$  is polynomial in  $|\mathcal{V}|$  and  $\log \lambda$ .

Then `ComputeIntroDecider` outputs the description of `RawIntroDecider` with the first three input tapes hardwired to  $\mathcal{V}'$ ,  $\lambda$ , and  $\ell$ , respectively, yielding the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  corresponding to the verifier  $\mathcal{V}'$  and parameters  $(\lambda, \ell)$ . Computing this description takes  $\text{poly}(|\mathcal{V}|, \log \lambda, \log \ell)$  time. Thus overall the complexity of `ComputeIntroDecider` is  $\text{poly}(|\mathcal{V}|, \log \lambda, \log \ell)$ , which is polynomial in its input length.

The description length of  $\hat{\mathcal{D}}^{\text{INTRO}}$  is simply the description length of `RawIntroDecider` (which is a universal constant), plus the description length of  $\mathcal{V}'$  (which is at most  $\lambda$ ), plus the description lengths of  $\lambda$  and  $\ell$ , which altogether totals  $\text{poly}(\lambda, \ell)$  as desired. The time complexity of  $\hat{\mathcal{D}}^{\text{INTRO}}$  follows from the time complexity of `RawIntroDecider`, which we established is  $\text{poly}(R, \ell) = \text{poly}(2^{\lambda n}, \ell)$ .

Finally, we note that if  $|\mathcal{V}| \leq \lambda$ , then by construction the output of `ComputeIntroDecider` is the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  corresponding to the verifier  $\mathcal{V}$  and parameters  $(\lambda, \ell)$ .  $\square$

### 8.2.2 Introspection theorem

We are now ready to state the introspection theorem.

**Theorem 8.3** (Introspection theorem). *There is a polynomial time Turing machine `ComputeIntroVerifier` which takes as input a tuple  $(\mathcal{V}, \lambda, \ell)$  for  $\lambda, \ell \in \mathbb{N}$  and returns the description of the introspective verifier  $\mathcal{V}^{\text{INTRO}}$ . For all  $\ell \in \mathbb{N}$ ,  $\mathcal{V}^{\text{INTRO}}$  is a 5-level normal-form verifier with complexity measures*

$$\begin{aligned} \text{TIME}_{\mathcal{S}^{\text{INTRO}}}(n) &= \text{poly}(n, \lambda, \ell), \\ \text{TIME}_{\mathcal{D}^{\text{INTRO}}}(n) &= \text{poly}(2^{\lambda n}, \ell), \\ |\mathcal{D}^{\text{INTRO}}| &= \text{poly}(\lambda, \ell). \end{aligned}$$

Moreover, for all  $\ell$ , there exist constants  $a > 0$ ,  $0 < b < 1$  (depending on  $\ell$ ), and a function  $\delta(\varepsilon, n) = a((\lambda n)^{a\varepsilon^b} + (\lambda n)^{-b})$  such that for all  $n \in \mathbb{N}$  and  $\varepsilon \geq 0$  the following statements hold if  $\mathcal{V}$  is a  $\lambda$ -bounded  $\ell$ -level verifier.

1. (**Completeness**) If  $\mathcal{V}_{2^n}$  has a projective, consistent, and commuting (PCC) strategy with value 1 then  $\mathcal{V}_n^{\text{INTRO}}$  also has a PCC strategy with value 1.
2. (**Soundness**) If  $\text{val}^*(\mathcal{V}_n^{\text{INTRO}}) > 1 - \varepsilon$ , then  $\text{val}^*(\mathcal{V}_{2^n}) \geq 1 - \delta(\varepsilon, n)$ .
3. (**Entanglement**) The entanglement bound  $\mathcal{E}$  defined in Definition 5.12 satisfies

$$\mathcal{E}(\mathcal{V}_n^{\text{INTRO}}, 1 - \varepsilon) \geq \max \left\{ \mathcal{E}(\mathcal{V}_{2^n}, 1 - \delta(\varepsilon, n)), (1 - \delta(\varepsilon, n)) 2^{2^{\lambda n}} \right\}.$$

*Proof.* The Turing machine `ComputeIntroVerifier` does the following on input  $(\mathcal{V}, \ell)$ : it first computes

$$\begin{aligned} \hat{\mathcal{S}}^{\text{INTRO}} &= \text{ComputeIntroSampler}(\lambda, \ell), \\ \hat{\mathcal{D}}^{\text{INTRO}} &= \text{ComputeIntroDecider}(\mathcal{V}, \lambda, \ell), \end{aligned}$$

using Lemmas 8.1 and 8.2, runs the detyping procedure from Definition 6.17 to compute

$$\mathcal{V}^{\text{INTRO}} = \text{Detype}(\hat{\mathcal{V}}^{\text{INTRO}})$$



where  $\hat{\mathcal{V}}^{\text{INTRO}} = (\hat{\mathcal{S}}^{\text{INTRO}}, \hat{\mathcal{D}}^{\text{INTRO}})$  and then outputs the resulting detyped verifier. This takes time  $\text{poly}(|\mathcal{V}|, \log \lambda, \log \ell)$ , which is polynomial in the input length of `ComputeIntroVerifier`. The complexity parameters of the typed sampler  $\hat{\mathcal{S}}^{\text{INTRO}}$  and typed decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  follow from Lemmas 8.1 and 8.2. The complexity parameters of the detyped verifier  $\mathcal{V}^{\text{INTRO}}$  then follow from Lemma 6.18. As  $\hat{\mathcal{V}}^{\text{INTRO}}$  is a 3-level verifier,  $\mathcal{V}^{\text{INTRO}}$  is 5-level as detyping will increase the number of levels by at most 2.

The completeness part of the theorem is proven in Section 8.3 and the soundness and entanglement properties are proven in Section 8.4.  $\square$

### 8.3 Completeness of the introspective verifier

In this section we establish the completeness property of the introspection game: if for  $N = 2^n$ ,  $\mathcal{V}_N$  has a PCC strategy (see Definition 5.11) with value 1, then so does  $\mathcal{V}_n^{\text{INTRO}}$ . For this we describe the actions that are expected of the players in the introspection game (i.e. the “honest strategy”). We first prove several preliminary lemmas that will be used in both the completeness and soundness analysis.

#### 8.3.1 Preliminary lemmas

The lemmas in this section are stated for general fields  $\mathbb{F}$  and generalized Pauli observables and projectors, although in the application to introspection games we use  $\mathbb{F} = \mathbb{F}_2$ ,  $\omega = -1$ , and qubit Pauli observables and projectors. Furthermore, the Pauli observables  $\tau^W(v)$  and projectors  $\tau_u^W$  act on  $\mathbb{C}^{\mathbb{F}^k}$  for some integer  $k$  (in our application, we set  $k = R$ ).

The following lemma generalizes Eq. (21).

**Lemma 8.4.** *Let  $L : \mathbb{F}^k \rightarrow \mathbb{F}^k$  be a linear map, and let  $W \in \{X, Z\}$ . For each  $a$  in the range of  $L$ , let  $u_a \in \mathbb{F}^k$  be such that  $L(u_a) = a$ .*

1. *For each  $v \in \ker(L)^\perp$ ,*

$$\tau^W(v) = \sum_{a \in \mathbb{F}^k} \omega^{\text{tr}(u_a \cdot v)} \tau_{[L(\cdot)=a]}^W.$$

2. *For all  $a$  in the range of  $L$ ,*

$$\tau_{[L(\cdot)=a]}^W = \mathbb{E}_{v \sim \ker(L)^\perp} \omega^{-\text{tr}(v \cdot u_a)} \tau^W(v).$$

*Proof.* Let  $V$  denote the image of  $\mathbb{F}^k$  under  $L$ . Let  $a \in V$ ,  $v \in \ker(L)^\perp$ . For all  $u, u' \in L^{-1}(a)$ , we have that  $u - u' \in \ker(L)$  and thus  $\text{tr}(u \cdot v) = \text{tr}(u' \cdot v)$ . As a result, using (20), for any  $v \in \ker(L)^\perp$  it holds that

$$\tau^W(v) = \sum_{u \in \mathbb{F}^k} \omega^{\text{tr}(u \cdot v)} \tau_u^W = \sum_{a \in V} \sum_{u \in L^{-1}(a)} \omega^{\text{tr}(u \cdot v)} \tau_u^W = \sum_{a \in V} \omega^{\text{tr}(u_a \cdot v)} \tau_{[L(\cdot)=a]}^W = \sum_{a \in \mathbb{F}^k} \omega^{\text{tr}(u_a \cdot v)} \tau_{[L(\cdot)=a]}^W,$$

where in the last equality we used that for  $a \notin V$ , the projector  $\tau_{[L(\cdot)=a]}^W$  vanishes. This shows the first item.

For the second item,

$$\begin{aligned}
\tau_{[L(\cdot)=a]}^W &= \sum_{u \in L^{-1}(a)} \tau_u^W \\
&= \sum_{u \in L^{-1}(a)} \mathbb{E}_{v \sim \mathbb{F}^k} \left( \omega^{-\text{tr}(u \cdot v)} \tau^W(v) \right) \\
&= \sum_{u \in \ker(L)} \mathbb{E}_{v \sim \mathbb{F}^k} \left( \omega^{-\text{tr}((u_a+u) \cdot v)} \tau^W(v) \right) \\
&= \frac{|\ker(L)|}{|\mathbb{F}^k|} \sum_{v \in \mathbb{F}^k} \left( \left( \mathbb{E}_{u \sim \ker(L)} \omega^{-\text{tr}(u \cdot v)} \right) \omega^{-\text{tr}(u_a \cdot v)} \tau^W(v) \right) \\
&= \mathbb{E}_{v \in \ker(L)^\perp} \omega^{-\text{tr}(v \cdot u_a)} \tau^W(v),
\end{aligned}$$

where the second equality follows from (21) and the last uses the fact that  $|\mathbb{F}^k| = |\ker(L)| |\ker(L)^\perp|$ , as shown in Lemma 3.4, and Lemma 3.25 applied to the expectation over  $u$ .  $\square$

**Lemma 8.5** (Commuting  $X$  and  $Z$  measurements). *For all linear maps  $L, R : \mathbb{F}^k \rightarrow \mathbb{F}^k$  such that*

$$\ker(R)^\perp \subseteq \ker(L),$$

*the measurements  $\{\tau_{[L(\cdot)=b]}^Z\}_{b \in \mathbb{F}^k}$  and  $\{\tau_{[R(\cdot)=d]}^X\}_{d \in \mathbb{F}^k}$  commute.*

*Proof.* Let  $b, d \in \mathbb{F}^k$ . If either  $b$  is not in the range of  $L$ , or  $d$  is not in the range of  $R$ , then at least one of  $\tau_{[L(\cdot)=b]}^Z$  or  $\tau_{[R(\cdot)=d]}^X$  is 0, and thus the operators trivially commute. Otherwise, both  $b$  and  $d$  are in the range of  $L$  and  $R$ , respectively. Let  $a_0 \in L^{-1}(b)$ ,  $c_0 \in R^{-1}(d)$ . By Lemma 8.4,

$$\begin{aligned}
\tau_{[L(\cdot)=b]}^Z &= \mathbb{E}_{u \in \ker(L)^\perp} \omega^{-\text{tr}(u \cdot a_0)} \tau^Z(u), \\
\tau_{[R(\cdot)=d]}^X &= \mathbb{E}_{v \in \ker(R)^\perp} \omega^{-\text{tr}(v \cdot c_0)} \tau^X(v).
\end{aligned}$$

For any  $v \in \ker(R)^\perp$ , by assumption  $v \in \ker(L)$ , so for  $u \in \ker(L)^\perp$  it holds that  $u \cdot v = 0$ . Thus  $\tau^Z(u)$  and  $\tau^X(v)$  commute, and it follows that  $\tau_{[L(\cdot)=b]}^Z$  and  $\tau_{[R(\cdot)=d]}^X$  commute as well.  $\square$

### 8.3.2 Completeness of the introspective verifier

*Proof of the completeness part of Theorem 8.3.* We analyze the completeness of the *typed* verifier  $\hat{\mathcal{V}}^{\text{INTRO}}$ ; the corresponding completeness of the *detyped* verifier  $\mathcal{V}^{\text{INTRO}} = \text{Detype}(\hat{\mathcal{V}}^{\text{INTRO}})$  follows from Lemma 6.18, and the fact that the type set  $\mathcal{T}^{\text{INTRO}}$  has size  $O(\ell)$ .

**Completeness.** We first show completeness. Let  $n \geq 1$  be an index for  $\hat{\mathcal{V}}^{\text{INTRO}}$  and  $N = 2^n$  be the corresponding index for  $\mathcal{V}$ . Since we assume that the verifier  $\mathcal{V}$  is  $\lambda$ -bounded, we have that

$$\max \{ \text{TIME}_{\mathcal{S}}(N), \text{TIME}_{\mathcal{D}}(N) \} \leq N^\lambda = R. \quad (62)$$

This assumption on the time complexity of  $\mathcal{V}$  ensures that  $\hat{\mathcal{D}}^{\text{INTRO}}$  never aborts due to a timeout. Let  $L^v = L^{v, N}$  denote the CL function of the original sampler  $\mathcal{S}$  on index  $N$  corresponding to player  $v \in \{A, B\}$ . Let

$R = N^\lambda$ , and let  $\text{introparams}(R) = (q, m, d)$ , as in Section 8.2. Set  $Q = M \log q$  where  $M = 2^m$ ; this represents the number of qubits that are certified by the Pauli basis test parameterized by  $\text{introparams}(R)$ .

Let  $\mathcal{S} = (|AUX\rangle, A, B)$  be a PCC strategy for  $\mathcal{V}_N$  with value 1. We first construct a PCC strategy  $\mathcal{S}_n^{\text{INTRO}}$  for the typed verifier  $\hat{\mathcal{V}}_n^{\text{INTRO}}$  with value 1. We then conclude using Lemma 6.18.

	$V_1^v$	$V_2^v$	$V_3^v$	AUX
(INTRO, $v$ )	$\sigma_{L_1}^Z$	$\sigma_{L_2}^Z$	$\sigma_{L_3}^Z$	$A^x / B^x$
(SAMPLE, $v$ )	$\sigma^Z$	$\sigma^Z$	$\sigma^Z$	$A^x / B^x$
(READ, $v$ )	$\sigma_{L_1}^Z \sigma_{L_1^\perp}^X$	$\sigma_{L_2}^Z \sigma_{L_2^\perp}^X$	$\sigma_{L_3}^Z \sigma_{L_3^\perp}^X$	$A^x / B^x$
(HIDE <sub>3</sub> , $v$ )	$\sigma_{L_1}^Z \sigma_{L_1^\perp}^X$	$\sigma_{L_2}^Z \sigma_{L_2^\perp}^X$	$\sigma_{L_3^\perp}^X$	$I$
(HIDE <sub>2</sub> , $v$ )	$\sigma_{L_1}^Z \sigma_{L_1^\perp}^X$	$\sigma_{L_2^\perp}^X$	$\sigma^X$	$I$
(HIDE <sub>1</sub> , $v$ )	$\sigma_{L_1^\perp}^X$	$\sigma^X$	$\sigma^X$	$I$

The left-most column denotes the introspection/hiding questions that a player may receive. The top row denotes the registers corresponding to the factor spaces of a CL function  $L^v$  (we note that the partition of the registers depends on the prefixes), as well as the register corresponding to the state  $|AUX\rangle$  coming from the original PCC strategy  $\mathcal{S}$ . We use  $\sigma_{L_j}^Z$  as shorthand for  $\sigma_{[(L_{j,x_{<j}^v})^\perp(\cdot)=x_j]}^Z$  and similarly  $\sigma_{L_j^\perp}^X$  for  $\sigma_{[(L_{j,x_{<j}^v})^\perp(\cdot)=x_j^\perp]}^X$ . A symbol  $I$  means that the register is left unmeasured.

Figure 11: Summary of the honest strategy  $\mathcal{S}_n^{\text{INTRO}}$  for  $\hat{\mathcal{V}}_n^{\text{INTRO}}$  for a 3-level sampler.

**Remark 8.6.** Note that by definition in  $\mathcal{V}_n^{\text{INTRO}}$  the players receive questions  $(x, y)$  that are sampled according to the distribution  $\mu_{\hat{\mathcal{S}}_{\text{INTRO}, n}^{\text{INTRO}}}^{\text{GINTRO}}$  associated with the downsized typed sampler  $\kappa(\hat{\mathcal{S}}^{\text{INTRO}})$ . Using the definition of the downsized typed sampler; Definition 6.6, and Lemma 4.12 the distribution is identical to the distribution  $\mu_{\hat{\mathcal{S}}_{\text{INTRO}, n}^{\text{GINTRO}}}$ , up to the bijective mapping  $\kappa$ . This mapping can be computed by the players themselves. Therefore, we construct a strategy for players that receive questions from  $\mu_{\hat{\mathcal{S}}_{\text{INTRO}, n}^{\text{GINTRO}}}$ , and a strategy for questions from  $\mu_{\hat{\mathcal{S}}_{\text{INTRO}, n}^{\text{GINTRO}}}$  follows immediately.

The strategy  $\mathcal{S}_n^{\text{INTRO}}$  uses the state  $|\text{EPR}_2\rangle^{\otimes(Q+1)} \otimes |AUX\rangle$ , where recall that  $|\text{EPR}_2\rangle = \frac{1}{\sqrt{2}}(|00\rangle + |11\rangle)$ . For all register subspaces  $K \subseteq \mathbb{F}_2^Q$  (see Definition 4.1 for the definition of register subspace), whenever we refer to “register  $K$ ”, we mean the qubits of  $|\text{EPR}_2\rangle^{\otimes Q}$  corresponding to  $K$  (see Section 3.5). The most frequent register subspace we consider is  $V$ , spanned by  $e_1, \dots, e_{s(N)}$ . We write  $\bar{V}$  for the complement of  $V$ , i.e. the register subspace spanned by  $e_{s(N)+1}, \dots, e_Q$ . (Note that  $s(N) \leq R$  by assumption (62), and by Lemma 7.17 we have  $R \leq M \leq Q$ .) Then  $|\text{EPR}_2\rangle^{\otimes Q} = |\text{EPR}\rangle_V \otimes |\text{EPR}\rangle_{\bar{V}}$ .

Let  $\mathcal{S}_n^{\text{PAULI}}$  be the honest Pauli strategy with respect to  $\text{introparams}$  defined in the proof of Lemma 7.13. Using the isomorphism between  $\mathbb{C}^q$  and  $(\mathbb{C}^2)^{\otimes \log q}$  specified by Lemma 3.26, the measurements of  $\mathcal{S}_n^{\text{PAULI}}$

can be treated as qubit Pauli measurements acting on the maximally entangled state over qubits. Furthermore the questions and answer labels of the measurements can be viewed as binary strings using the bijection  $\kappa : \mathbb{F}_q \rightarrow \mathbb{F}_2^{\log q}$  specified in Section 3.3. For a question with a type from  $\mathcal{T}^{\text{PAULI}}$  the player measures the shared state  $|\text{EPR}_2\rangle^{\otimes(Q+1)}$  using the measurements specified by  $\mathcal{S}_n^{\text{PAULI}}$ , and reports the measurement outcomes. When a player receives questions of type  $\hat{t} \in \mathcal{T}^{\text{INTRO}} \setminus \mathcal{T}^{\text{PAULI}}$ , they perform measurements described as follows. (Below, whenever we write a Pauli operator  $\sigma_a^W$  the register on which the operator acts should always be clear from context, and is implicit from the space in which the outcome  $a$  ranges.) The reader may find it helpful to consult Figure 11 to see a summary of the honest strategy  $\mathcal{S}_n^{\text{INTRO}}$  for the special case when  $\ell = 3$ .

(INTRO,  $v$ ): The player performs the measurement

$$\left\{ \sigma_{[L^v(\cdot)=y]}^Z \right\} \quad (63)$$

to obtain an  $y \in V$ . Intuitively, the player has now introspectively sampled the question  $y$  for original player  $v$  in game  $\mathcal{V}_N$ . The player then measures  $|\text{AUX}\rangle$  using player A's measurement  $\{A_a^y\}$  from  $\mathcal{S}$  if  $v = \text{A}$  and using player B's measurement  $\{B_a^y\}$  if  $v = \text{B}$  to obtain an answer  $a$ . The player replies with  $(y, a)$ .

(SAMPLE,  $v$ ): The player measures their share of  $|\text{EPR}\rangle_V$  in the  $Z$  basis to obtain  $z \in V$ . Using this, they compute the question  $y = L^v(z)$ . The player then uses player  $v$ 's strategy and question  $y$  to measure  $|\text{AUX}\rangle$  and obtain outcome  $a$ . The player replies with  $(z, a)$ .

(READ,  $v$ ): The player first performs all measurements as in the (INTRO,  $v$ ) question for player  $v$  and records the outcomes as  $y \in L(V)$  and  $a \in \{0, 1\}^*$ . For  $j \in \{1, 2, \dots, \ell\}$ , the player measures their share of the state  $|\text{EPR}\rangle_V$  with the measurement

$$\left\{ \sigma_{[L_j^\perp(\cdot)=y_j^\perp]}^X \right\}_{y_j^\perp} \quad (64)$$

to obtain  $y^\perp = y_1^\perp + \dots + y_\ell^\perp$ . Here  $L_j^\perp$  is shorthand for the function  $(L_{j, y_{<j}}^v)^\perp$  defined in Item 6 of the decider description in Section 8.2. (That these operators are simultaneously measurable with the measurement in (63) follows from Lemma 8.5 and the fact that  $\ker(L_j^\perp)^\perp = \ker(L_j)$  by Lemma 3.4 and the definition of  $L_j^\perp$  in Section 8.2.)

The player measures  $|\text{AUX}\rangle$  with player  $v$ 's measurement strategy in  $\mathcal{S}$  for question  $y$  to obtain  $a$  and replies with  $(y, y^\perp, a)$ .

(HIDE $_k$ ,  $v$ ): The player performs the following sequence of measurements: first measure  $\left\{ \sigma_{[L_1^v(\cdot)=y_1]}^Z \right\}$  on register  $V_1^v$  to obtain  $y_1$ . Then, use  $y_1$  to specify the second linear function  $L_{2, y_1}^v(\cdot)$  and measure register  $V_{2, y_1}^v$  using  $\left\{ \sigma_{[L_{2, y_1}^v(\cdot)=y_2]}^Z \right\}$  to obtain  $y_2$ . This process continues until the  $(k-1)$ -th linear map  $L_{k-1, y_{<k-1}}^v(\cdot)$  has been measured to obtain  $y_{k-1}$  in factor space  $V_{k-1, y_{<k-1}}^v$ . Let  $y = y_1 + y_2 + \dots + y_{k-1}$ .

Next, for  $j \in \{1, 2, \dots, k\}$  the player measures

$$\left\{ \sigma_{[L_j^\perp(\cdot)=y_j^\perp]}^X \right\}_{y_j^\perp},$$

where  $L_j^\perp$  denotes the linear map  $(L_{j,y_{<j}}^v)^\perp$  as in the case  $(\text{READ}, v)$ . Let  $y^\perp = y_1^\perp + y_2^\perp + \dots + y_k^\perp$ , where each  $y_j^\perp$  is a vector in the factor space  $V_{j,y_{<j}}^v$ . Finally, the player measures register  $V_{>k}^v(y)$  using  $\{\sigma_{x_{>k}}^X\}$  to obtain outcome  $x_{>k}$ . Let  $x = x_{>k}$ . The player replies with  $(y, y^\perp, x)$ .

By definition, when player  $w$  performs the honest measurement for question  $(\text{INTRO}, A)$  and player  $\bar{w}$  performs the honest measurement for question  $(\text{INTRO}, B)$ , the joint outcome  $(y, y')$  has distribution  $\mu_{S,N}$ . In this case, the players play according to strategy  $\mathcal{S}$  and succeed with probability 1. In all other cases, it is straightforward to verify that the players succeed in all tests performed by  $\hat{\mathcal{D}}^{\text{INTRO}}$  (Figure 10) with probability 1. As a result, the value of this strategy is 1.

The strategy  $\mathcal{S}^{\text{INTRO}}$  is projective by construction. It is also consistent because of the assumed consistency of the strategy  $\mathcal{S}$  as well as consistency of the honest Pauli strategy  $\mathcal{S}^{\text{PAULI}}$ . Furthermore, note that  $\mathcal{S}^{\text{INTRO}}$  only calls  $\mathcal{S}$  for both players on question pairs such that both types are in  $\{\text{INTRO}, \text{SAMPLE}, \text{READ}\}$ . In all these cases,  $\mathcal{S}$  is called on a pair of questions  $(y, y')$  distributed as  $(L^v(z), L^{v'}(z))$  for  $v, v' \in \{A, B\}$  and  $z$  uniform in  $V$ . When  $v \neq v'$ , any such pair by definition has positive probability under  $\mu_{S,n}$ , and so by assumption the associated measurements from  $\mathcal{S}$  commute. On the other hand, when  $v = v'$ , then the players apply the same measurements from  $\mathcal{S}$ , and because  $\mathcal{S}$  only uses projective measurements, their measurements commute as well. Examining all other cases, it follows by direct inspection that the strategy commutes on all question pairs whose corresponding types appear as an edge in the graph  $G^{\text{INTRO}}$ . Thus the strategy commutes with respect to the support of the distribution  $\mu_{\hat{\mathcal{S}}^{\text{INTRO}}, n}$ .  $\square$

**Remark 8.7.** For future reference, we note that on any input  $(\mathcal{V}, \lambda, \ell)$ , the Turing machine `ComputeIntroVerifier` always returns a normal form verifier  $\hat{\mathcal{V}}^{\text{INTRO}} = (\hat{\mathcal{S}}^{\text{INTRO}}, \hat{\mathcal{D}}^{\text{INTRO}})$  — even if  $\mathcal{V}$  itself is not the description of a normal form verifier. This is because, for any two integer  $\lambda, \ell$ , `ComputeIntroSampler` $(\lambda, \ell)$  returns a sampler with field size 2, and for any  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ ,  $\lambda$  and  $\ell$  the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  specified in Figure 10 takes 7 inputs and always halts with a single-bit output, even if  $\mathcal{S}$  or  $\mathcal{D}$  themselves do not halt.

## 8.4 Soundness of the introspective verifier

The main result of this section is the soundness part of Theorem 8.3.

### 8.4.1 The Pauli twirl

A key tool in the proof of soundness is the *Pauli twirl*. In this section we introduce the Pauli twirl and establish several of its properties. The section closely follows Sections 8 and 10 of [NW19].

To begin, we define the twirl with respect to an arbitrary distribution over unitaries.

**Definition 8.8** (Twirl). Let  $\mu$  be a probability distribution over a finite set of unitary matrices. Then for any matrix  $A$ , the *twirl of  $A$  with respect to  $\mu$* , denoted  $\mathcal{T}_\mu(A)$ , is defined as

$$\mathcal{T}_\mu(A) = \mathbb{E}_{U \sim \mu} (UAU^\dagger).$$

In the next two lemmas we consider the Pauli twirl, in which the distribution  $\mu$  is over subsets of Pauli observables. First, we show how the Pauli twirl acts on Pauli matrices. Then, using this, we derive an expression for the Pauli twirl applied to general matrices.

**Lemma 8.9** (Pauli twirl of Pauli matrices). *Let  $V$  be a subspace of  $\mathbb{F}^k$ , let  $W \in \{X, Z\}$ , and let  $\mu$  be the uniform distribution over  $\{\tau^W(w) : w \in V\}$ . Let  $W' \neq W$  and  $u \in \mathbb{F}^k$ . Then*

$$\mathcal{F}_\mu(\tau^{W'}(u)) = \begin{cases} \tau^{W'}(u) & \text{if } u \in V^\perp, \\ 0 & \text{if } u \notin V^\perp. \end{cases}$$

*Proof.* Let  $u \in \mathbb{F}^k$ . Let  $c = 1$  if  $W = Z$  and let  $c = -1$  if  $W = X$ . Then

$$\begin{aligned} \mathcal{F}_\mu(\tau^{W'}(u)) &= \mathbb{E}_{z \sim V} (\tau^W(z) \tau^{W'}(u) \tau^W(z)^\dagger) \\ &= \mathbb{E}_{z \sim V} (\omega^{c \cdot \text{tr}(u \cdot z)} \tau^{W'}(u) \tau^W(z) \tau^W(z)^\dagger) \\ &= \left( \mathbb{E}_{z \sim V} \omega^{c \cdot \text{tr}(u \cdot z)} \right) \tau^{W'}(u). \end{aligned}$$

The lemma now follows from Lemma 3.25 and the fact that  $c \cdot u \in V^\perp$  if and only if  $u \in V^\perp$ .  $\square$

**Lemma 8.10** (Pauli twirl of general matrices). *Let  $V = \mathbb{F}^k$ , and let  $L : V \rightarrow V$  be a linear map. Let  $\zeta$  be the uniform distribution over  $\{\tau^Z(z) \mid z \in V\}$  and  $\chi$  the uniform distribution over  $\{\tau^X(x) \mid x \in \ker(L)\}$ . Let  $M$  be a matrix acting on  $\mathbb{C}^V \otimes \mathcal{H}_A$ , where  $\mathcal{H}_A$  is a finite dimensional Hilbert space. Then there exist matrices  $\{M^y\}_{y \in L(V)}$  acting on  $\mathcal{H}_A$  such that the twirl of  $M$  with respect to  $\zeta$  and  $\chi$  is given by*

$$(\mathcal{F}_\chi \circ \mathcal{F}_\zeta \otimes I_A)(M) = \sum_{y \in V} \tau_{[L(\cdot)=y]}^Z \otimes M^y. \quad (65)$$

Moreover, if we apply (65) to each element of a POVM measurement  $\{M_a\}$ , then for each  $y \in V$ , the set  $\{M_a^y\}$  also forms a POVM measurement.

*Proof.* The collection  $\{\tau^X(x) \tau^Z(z)\}_{x,z \in V}$  forms a basis for the complex linear space of matrices acting on  $\mathbb{C}^V$ . As a result, we can write

$$M = \sum_{x,z \in V} \tau^X(x) \tau^Z(z) \otimes M_{x,z},$$

for matrices  $M_{x,z}$  on  $\mathcal{H}_A$ . We now use Lemma 8.9 to compute the twirl first with respect to  $\zeta$  and then with respect to  $\chi$ :

$$\begin{aligned} (\mathcal{F}_\zeta \otimes I_A)(M) &= \sum_{x,z \in V} \mathcal{F}_\zeta(\tau^X(x) \tau^Z(z)) \otimes M_{x,z} = \sum_{z \in V} \tau^Z(z) \otimes M_{0,z}, \\ (\mathcal{F}_\chi \circ \mathcal{F}_\zeta \otimes I_A)(M) &= \sum_{z \in V} \mathcal{F}_\chi(\tau^Z(z)) \otimes M_{0,z} = \sum_{z \in \ker(L)^\perp} \tau^Z(z) \otimes M_{0,z}. \end{aligned}$$

For all  $y \in V$ , let  $u_y$  denote an arbitrary element of  $L^{-1}(y)$  if  $y$  is in the image of  $L$ ; otherwise, set  $u_y = 0$ . Expanding  $\tau^Z(z)$  using the first part of Lemma 8.4,

$$\begin{aligned} \sum_{z \in \ker(L)^\perp} \tau^Z(z) \otimes M_{0,z} &= \sum_{z \in \ker(L)^\perp} \sum_y \omega^{\text{tr}(u_y \cdot z)} \tau_{[L(\cdot)=y]}^Z \otimes M_{0,z} \\ &= \sum_y \tau_{[L(\cdot)=y]}^Z \otimes \left( \sum_{z \in \ker(L)^\perp} \omega^{\text{tr}(u_y \cdot z)} M_{0,z} \right). \end{aligned} \quad (66)$$

Equation (65) follows by setting  $M^y = \sum_{z \in \ker(L)^\perp} \omega^{\text{tr}(u_y \cdot z)} M_{0,z}$ .

For the “moreover” part, note first that whenever  $M \geq 0$  it holds that any twirl satisfies  $0 \leq \mathcal{T}_\mu(M)$ . As a result, each matrix  $M^y$  must be positive semi-definite due to Equation (66) and the fact that the  $\{\tau_{[L(\cdot)=y]}^Z\}_y$  matrices are orthogonal projections. Next, suppose  $\{M_a\}$  is a POVM measurement, and write  $N = \sum_a M_a$  for the identity matrix. Then by linearity, for each  $y \in V$ ,  $\sum_a M_a^y = N^y$ . In addition,  $N_{0,0} = I$ , and  $N_{x,z} = 0$  otherwise. As a result,  $N^y = N_{0,0} = I$ , and so  $\{M_a^y\}$  also forms a POVM.  $\square$

In the next few lemmas we derive a sufficient condition for a measurement to be close to its own Pauli twirl, namely that it satisfies certain commutation relations with the Pauli basis measurements.

**Lemma 8.11** (Commuting with Pauli basis implies commuting with Pauli observables). *Let  $\mathcal{X}, \mathcal{A}$  be finite sets and  $D$  be a distribution over  $\mathcal{X}$ . For each  $x \in \mathcal{X}$ , let  $V_x$  be a register subspace of  $V = \mathbb{F}^k$ , and let  $L_x : V_x \rightarrow V_x$  be a linear map.*

*Consider a state  $|\psi\rangle = |\text{EPR}\rangle_V \otimes |\text{AUX}\rangle$ , where  $|\text{EPR}\rangle_V \in \mathcal{H}_A \otimes \mathcal{H}_B$ , for  $\mathcal{H}_A, \mathcal{H}_B \cong \mathbb{C}^V$ , is defined in Definition 3.22 and  $|\text{AUX}\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$  is arbitrary. For each  $x \in \mathcal{X}$ , let  $\{M_a^x\}_{a \in \mathcal{A}}$  be a measurement on  $\mathcal{H}_A \otimes \mathcal{H}_{A'}$ . Let  $W \in \{X, Z\}$ . Then the following are equivalent on the state  $|\psi\rangle$ :*

- On average over  $x \sim D$ ,

$$[M_a^x, (\tau_{[L_x(\cdot)=y]}^W \otimes I_{\overline{V}_x} \otimes I_{A'})] \otimes I_B \approx_\varepsilon 0.$$

- On average over  $x \sim D$  and  $v$  drawn uniformly from  $\ker(L_x)^\perp$ ,

$$[M_a^x, (\tau^W(v) \otimes I_{\overline{V}_x} \otimes I_{A'})] \otimes I_B \approx_\varepsilon 0.$$

In the equations above, for each  $x \in \mathcal{X}$  we decompose  $\mathcal{H}_A = \mathbb{C}^{V_x} \otimes \mathbb{C}^{\overline{V}_x}$  and interpret  $\tau_{[L_x(\cdot)=y]}^W$  and  $\tau^W(v)$  for  $v \in \ker(L_x)^\perp$  as acting on  $\mathbb{C}^{V_x}$ .

*Proof.* For  $x \in \mathcal{X}$ ,  $a \in \mathcal{A}$ ,  $y$  in the range of  $L_x$ , and  $v \in \mathbb{F}^k$  define

$$\Delta_{a,y}^x = [M_a^x, (\tau_{[L_x(\cdot)=y]}^W \otimes I_{\overline{V}_x} \otimes I_{A'})] \otimes I_B,$$

$$\Delta_a^x(v) = [M_a^x, (\tau^W(v) \otimes I_{\overline{V}_x} \otimes I_{A'})] \otimes I_B.$$

By the first item of Lemma 8.4, for each  $v \in \ker(L_x)^\perp$ ,

$$\Delta_a^x(v) = \sum_{y \in V_x} \omega^{\text{tr}(u_y \cdot v)} \Delta_{a,y}^x,$$

where for every  $y$  in the range of  $L_x$ ,  $u_y$  is a fixed element in  $L_x^{-1}(y)$ . The expression for the closeness of  $\Delta_a^x$  to 0 on average over  $x \sim D$  and  $v \sim \ker(L_x)^\perp$  (i.e. the second quantity of the Lemma statement), is equal to

$$\mathbb{E}_{x \sim D} \mathbb{E}_{v \sim \ker(L_x)^\perp} \sum_a \|\Delta_a^x(v)|\psi\rangle\|^2,$$

and can be expanded as

$$\mathbb{E}_{x \sim D} \mathbb{E}_{v \sim \ker(L_x)^\perp} \sum_a \langle \psi | \sum_{y,y'} \omega^{\text{tr}((u'_y - u_y) \cdot v)} (\Delta_{a,y}^x)^\dagger \Delta_{a,y'}^x | \psi \rangle. \quad (67)$$

If  $y \neq y'$ , then by definition  $L_x(u_y) \neq L_x(u_{y'})$ , and so  $u'_y - u_y$  is not in  $\ker(L_x)$ . Lemma 3.25 (with  $V = \ker(L_x)^\perp$ ) thus implies that (67) equals

$$\mathbb{E}_{x \sim D} \sum_a \langle \psi | \sum_y ((\Delta_{a,y}^x)^\dagger \Delta_{a,y}^x) | \psi \rangle = \mathbb{E}_{x \sim D} \sum_{a,y} \|\Delta_{a,y}^x | \psi \rangle\|^2$$

which is the closeness of  $\Delta_{a,y}^x$  to 0 on average over  $x \sim D$ . Thus,  $\Delta_a^x(v) \approx_\varepsilon 0$  on average over  $x$  and  $v$  if and only if  $\Delta_{a,y}^x \approx_\varepsilon 0$  on average over  $x$ .  $\square$

**Lemma 8.12** (Commuting implies twirl). *Let  $\mathcal{X}$  be a finite set and  $D$  a distribution on  $\mathcal{X}$ . For each  $x \in \mathcal{X}$ , let  $\{M_a^x\}$  be a POVM on  $\mathcal{H}_A$ , and let  $\mu_x$  be a distribution over unitary matrices acting on  $\mathcal{H}_A$ . Suppose that on average over  $x \sim D$  and  $U \sim \mu_x$  it holds that*

$$[M_a^x, U^\dagger] \otimes I_B \approx_\varepsilon 0,$$

where the commutator is evaluated on a state  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ . Then on average over  $x \sim D$ ,

$$M_a^x \otimes I_B \approx_\varepsilon \mathcal{T}_{\mu_x}(M_a^x) \otimes I_B.$$

*Proof.* Observe that

$$\mathbb{E}_{x \sim D} \sum_a \left\| (\mathcal{T}_{\mu_x}(M_a^x) - M_a^x) \otimes I_B | \psi \rangle \right\|^2 = \mathbb{E}_{x \sim D} \sum_a \left\| \mathbb{E}_{U \sim \mu_x} (U [M_a^x, U^\dagger]) \otimes I_B | \psi \rangle \right\|^2. \quad (68)$$

Applying Jensen's inequality, the right-hand side of (68) is at most

$$\mathbb{E}_{x \sim D} \mathbb{E}_{U \sim \mu_x} \sum_a \left\| (U [M_a^x, U^\dagger]) \otimes I_B | \psi \rangle \right\|^2 = \mathbb{E}_{x \sim D} \mathbb{E}_{U \sim \mu_x} \sum_a \left\| [M_a^x, U^\dagger] \otimes I_B | \psi \rangle \right\|^2,$$

using the unitary invariance of the Euclidean norm. This last quantity is  $O(\varepsilon)$ , by assumption.  $\square$

**Lemma 8.13** (Commuting with each implies commuting with both). *Let  $\mathcal{X}$  be a finite set and  $D$  a distribution on  $\mathcal{X}$ . For each  $x \in \mathcal{X}$ , let  $\{M_a^x\}$  be a POVM on  $\mathcal{H}_A$  and let  $\mu_{x,1}, \mu_{x,2}$  be two distributions over unitary matrices acting on  $\mathcal{H}_A$ . Suppose that for each  $i \in \{1, 2\}$ , on average over  $x \sim D$  and  $U_i \sim \mu_{x,i}$ ,*

$$[M_a^x, U_i^\dagger] \otimes I_B \approx_\varepsilon 0, \quad (69)$$

where the expression is evaluated on some state  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$ , where  $\mathcal{H}_B \cong \mathcal{H}_A$ . Suppose further that on average over  $x \sim D$  and  $U_2 \sim \mu_{x,2}$ ,

$$U_2^\dagger \otimes I_B \approx_\varepsilon I_A \otimes U_2. \quad (70)$$

(The corresponding statement for  $i = 1$  is not needed.) Then on average over  $x \sim D$ ,  $U_1 \sim \mu_{x,1}$ , and  $U_2 \sim \mu_{x,2}$ ,

$$[M_a^x, U_1^\dagger U_2^\dagger] \otimes I_B \approx_\varepsilon 0.$$

*Proof.* The claim follows from the following sequence of approximations:

$$\begin{aligned} M_a^x U_1^\dagger U_2^\dagger \otimes I_B &\approx_\varepsilon M_a^x U_1^\dagger \otimes U_2 && \text{(by (70))} \\ &\approx_\varepsilon U_1^\dagger M_a^x \otimes U_2 && \text{(by (69) for } i = 1) \\ &\approx_\varepsilon U_1^\dagger M_a^x U_2^\dagger \otimes I_B && \text{(by (70))} \\ &\approx_\varepsilon U_1^\dagger U_2^\dagger M_a^x \otimes I_B, && \text{(by (69) for } i = 2) \end{aligned}$$

where each step also uses Fact 5.19. This is equivalent to  $[M_a^x, U_1^\dagger U_2^\dagger] \otimes I_B \approx_\varepsilon 0$ , completing the proof.  $\square$



The following is a slight generalization of [NW19, Fact 4.25], and we give a similar proof.

**Lemma 8.14** (Close to sub-measurement implies close to measurement). *Let  $\mathcal{X}, \mathcal{A}$  be finite sets and  $D$  be a distribution on  $\mathcal{X}$ . Suppose that for each  $x \in \mathcal{X}$ ,  $\{A_a^x\}_{a \in \mathcal{A}}$  is a projective measurement and  $\{B_a^x\}_{a \in \mathcal{A}}$  is a set of matrices such that each  $B_a^x$  is positive semidefinite and  $\sum_a B_a^x \leq I$ . Suppose  $\{C_a^x\}_{a \in \mathcal{A}}$  is a POVM such that  $C_a^x \geq B_a^x$  for all  $x$  and  $a$ . Then if, on average over  $x \sim D$ ,  $A_a^x \approx_\varepsilon B_a^x$ , then, on average over  $x \sim D$ ,  $A_a^x \approx_{\varepsilon^{1/2}} C_a^x$ .*

*Proof.* By the triangle inequality,

$$\mathbb{E}_x \sum_a \|(A_a^x - C_a^x)|\psi\rangle\|^2 \leq 2 \mathbb{E}_x \sum_a \|(A_a^x - B_a^x)|\psi\rangle\|^2 + 2 \mathbb{E}_x \sum_a \|(C_a^x - B_a^x)|\psi\rangle\|^2.$$

The first term on the right-hand side is  $O(\varepsilon)$  by assumption. For the second,

$$\begin{aligned} \mathbb{E}_x \sum_a \|(C_a^x - B_a^x)|\psi\rangle\|^2 &= \mathbb{E}_x \sum_a \langle \psi | (C_a^x - B_a^x)^2 | \psi \rangle \leq \mathbb{E}_x \sum_a \langle \psi | (C_a^x - B_a^x) | \psi \rangle \\ &= 1 - \mathbb{E}_x \sum_a \langle \psi | B_a^x | \psi \rangle \leq 1 - \mathbb{E}_x \sum_a \langle \psi | (B_a^x)^2 | \psi \rangle, \end{aligned}$$

where the middle inequality uses  $0 \leq C_a^x - B_a^x \leq I$  for all  $x, a$ . Write  $1 = \mathbb{E}_x \sum_a \langle \psi | (A_a^x)^2 | \psi \rangle$ , which holds because  $A$  is a projective measurement. Then

$$\begin{aligned} \mathbb{E}_x \sum_a \langle \psi | ((A_a^x)^2 - (B_a^x)^2) | \psi \rangle &= \Re \left( \mathbb{E}_x \sum_a \langle \psi | (A_a^x + B_a^x)(A_a^x - B_a^x) | \psi \rangle \right) \\ &\leq \mathbb{E}_x \sqrt{\sum_a \|(A_a^x + B_a^x)|\psi\rangle\|^2} \cdot \sqrt{\sum_a \|(A_a^x - B_a^x)|\psi\rangle\|^2} \end{aligned}$$

where the first equality follows from the fact that  $A_a^x$  and  $B_a^x$  are Hermitian. For each  $x \in X$  the first square root is  $O(1)$ . This allows us to move the expectation into the second square root by Jensen's inequality. The result is  $O(\varepsilon^{1/2})$  by assumption.  $\square$

Now we put everything together to show the main result of this section.

**Lemma 8.15.** *Let  $\mathcal{X}, \mathcal{A}$  be finite sets and  $D$  be a distribution over  $\mathcal{X}$ . For each  $x \in \mathcal{X}$ , let  $V_x$  be a register subspace of  $V = \mathbb{F}^k$ , let  $U_x$  be a register subspace of  $V_x$ , and let  $L_x : U_x \rightarrow U_x$  be a linear map.*

*Consider a state  $|\psi\rangle = |\text{EPR}\rangle_V \otimes |\text{AUX}\rangle$ , where  $|\text{EPR}\rangle_V \in \mathcal{H}_A \otimes \mathcal{H}_B$ , for  $\mathcal{H}_A, \mathcal{H}_B \cong \mathbb{C}^V$ , is defined in Definition 3.22 and  $|\text{AUX}\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$  is arbitrary. For each  $x \in \mathcal{X}$ , let  $\{M_{y,a}^x\}_{y \in U_x, a \in \mathcal{A}}$  be a projective measurement on  $\mathcal{H}_{V_x} \otimes \mathcal{H}_{A'}$ . Suppose that on average over  $x \sim D$  the following conditions hold.*

$$\begin{aligned} (\text{Consistency}): \quad & (M_y^x \otimes I_{\overline{V_x}} - \tau_{[L_x(\cdot)=y]}^Z \otimes I_{\overline{U_x}} \otimes I_{A'}) \otimes I_B \approx_\varepsilon 0, \\ (\text{Commutation}): \quad & [M_{y,a}^x \otimes I_{\overline{V_x}}, \tau_z^Z \otimes I_{\overline{U_x}} \otimes I_{A'}] \otimes I_B \approx_\varepsilon 0, \\ & [M_{y,a}^x \otimes I_{\overline{V_x}}, \tau_{[L_x^\perp(\cdot)=y^\perp]}^X \otimes I_{\overline{U_x}} \otimes I_{A'}] \otimes I_B \approx_\varepsilon 0. \end{aligned}$$

Here, the projector  $\tau_{[L_x(\cdot)=y]}^Z$  acts on the register subspace  $\mathcal{H}_{U_x}$ , and  $\overline{U_x}$  and  $\overline{V_x}$  denote the complementary register subspaces of  $U_x$  and  $V_x$ , respectively, within  $V$ . Furthermore, the answer summations are over all  $y, z \in U_x$  and  $a \in \mathcal{A}$ .

Then for each  $x \in \mathcal{X}$  and  $y \in U_x$ , there exists a POVM measurement  $\{M_a^{x,y}\}_{a \in \mathcal{A}}$  on  $\mathcal{H}_{V_x \setminus U_x} \otimes \mathcal{H}_{A'}$  such that on average over  $x \sim D$ ,

$$(M_{y,a}^x \otimes I_{\overline{V_x}}) \otimes I_B \approx_{\varepsilon^{1/2}} \left( \tau_{[L_x(\cdot)=y]}^Z \otimes M_a^{x,y} \otimes I_{\overline{V_x}} \right) \otimes I_B .$$

*Proof.* For each  $x \in X$ , let  $\zeta_x$  be the uniform distribution over  $U_x$  and  $\chi_x$  be the uniform distribution over  $\ker(L_x)$ . Applying Lemma 8.11 to the first commutation assumption (letting “ $L_x$ ” in Lemma 8.11 be the identity map on  $U_x$ , and letting “ $D$ ” be the uniform distribution over  $\ker(L_x)^\perp = U_x$ ), we get that on average over  $x \sim \mathcal{X}$  and  $v \sim \zeta_x$ , we have

$$[M_{y,a}^x \otimes I_{\overline{V_x}}, \tau^Z(v) \otimes I_{\overline{U_x}} \otimes I_{A'}] \otimes I_B \approx_\varepsilon 0 .$$

Applying Lemma 8.11 to the second commutation assumption (letting “ $L_x$ ” in Lemma 8.11 be the  $L_x^\perp$  in the statement of Lemma 8.15 and letting  $D$  be the uniform distribution over  $\ker(L_x^\perp)^\perp$ , which is  $\ker(L_x)$  by Lemma 3.12), we get on average over  $x \sim \mathcal{X}$  and  $v \sim \zeta_x$ , we have

$$[M_{y,a}^x \otimes I_{\overline{V_x}}, \tau^X(v) \otimes I_{\overline{U_x}} \otimes I_{A'}] \otimes I_B \approx_\varepsilon 0 .$$

By Lemma 8.13, this implies that on average over  $x \sim D$ ,  $u \sim \zeta_x$ , and  $v \sim \chi_x$ ,

$$[M_{y,a}^x \otimes I_{\overline{V_x}}, \tau^Z(u) \tau^X(v) \otimes I_{\overline{U_x}} \otimes I_{A'}] \otimes I_B \approx_\varepsilon 0 .$$

By Lemma 8.12 and Lemma 8.10, this implies that on average over  $x \sim D$ ,

$$\begin{aligned} M_{y,a}^x \otimes I_{\overline{V_x}} \otimes I_B &\approx_\varepsilon (\mathcal{F}_{\chi_x} \circ \mathcal{F}_{\zeta_x}(M_{y,a}^x)) \otimes I_{\overline{V_x}} \otimes I_B \\ &= \left( \sum_{y' \in V_x} \tau_{[L_x(\cdot)=y']}^Z \otimes M_{y,a}^{x,y'} \right) \otimes I_{\overline{V_x}} \otimes I_B , \end{aligned} \quad (71)$$

for some POVM measurement  $\{M_{y,a}^{x,y'}\}$  on  $\mathcal{H}_{V_x \setminus U_x} \otimes \mathcal{H}_{A'}$ .

In the following sequence of equations, whenever an operator does not act on a subsystem it should be assumed that it is appropriately tensored with the identity. For clarity, we explicitly indicate using a subscript A or B whether a Pauli operator acts on  $\mathcal{H}_A$  or  $\mathcal{H}_B$ . Then on average over  $x \sim D$  we have

$$\begin{aligned} M_{y,a}^x &= M_{y,a}^x \cdot M_y^x && (M^x \text{ is projective}) \\ &\approx_\varepsilon M_{y,a}^x \cdot (\tau_{[L_x(\cdot)=y]}^Z)_A && (\text{Consistency assumption}) \\ &\approx_0 M_{y,a}^x \otimes (\tau_{[L_x(\cdot)=y]}^Z)_B && (\text{Paulis are self-consistent}) \\ &\approx_\varepsilon \left( \sum_{y'} (\tau_{[L_x(\cdot)=y']}^Z)_A \otimes M_{y,a}^{x,y'} \right) \otimes (\tau_{[L_x(\cdot)=y]}^Z)_B && (\text{Equation (71)}) \\ &\approx_0 \sum_{y'} (\tau_{[L_x(\cdot)=y']}^Z \tau_{[L_x(\cdot)=y]}^Z)_A \otimes M_{y,a}^{x,y'} && (\text{Paulis are self-consistent}) \\ &= (\tau_{[L_x(\cdot)=y]}^Z)_A \otimes M_{y,a}^{x,y} , \end{aligned}$$

where  $\approx_0$  indicates equality with respect to the state  $|EPR\rangle_V \otimes |AUX\rangle$ , and we have repeatedly used Fact 5.19. This is essentially the statement promised by the lemma, except that  $\{M_{y,a}^{x,y}\}_a$  does not necessarily sum to identity (since we only sum over  $a$ ). To remedy this, define  $M_a^{x,y} = \sum_{y'} M_{y,a}^{x,y'}$  and note that  $M_a^{x,y} \geq M_{y,a}^{x,y}$ , so by Lemma 8.14 and the fact that  $M^x$  is projective,

$$(M_{y,a}^x \otimes I_{\overline{V_x}}) \otimes I_B \approx_{\varepsilon^{1/2}} \left( \tau_{[L_x(\cdot)=y]}^Z \otimes M_a^{x,y} \otimes I_{\overline{V_x}} \right) \otimes I_B . \quad (72)$$

$\{M_a^{x,y}\}$  is the POVM measurement guaranteed in the lemma statement, which concludes the proof.  $\square$

### 8.4.2 Preliminary lemmas

We show a few simple lemmas that allow us to argue about measurements that have a decomposition across a tensor product of two Hilbert spaces, within the space of a single player.

**Lemma 8.16.** *Let  $\mathcal{A}, \mathcal{B}$  be finite sets. Let  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  be a state. Consider the following: for all  $a \in \mathcal{A}$ ,*

1. *Let  $\mathcal{H}_{A,a}, \mathcal{H}_{B,a}, \mathcal{H}_{A',a}$ , and  $\mathcal{H}_{B',a}$  be Hilbert spaces such that*

$$\mathcal{H}_A = \mathcal{H}_{A,a} \otimes \mathcal{H}_{A',a} \quad \text{and} \quad \mathcal{H}_B = \mathcal{H}_{B,a} \otimes \mathcal{H}_{B',a},$$

*and let  $|\psi_{\text{QUE},a}\rangle \in \mathcal{H}_{A,a} \otimes \mathcal{H}_{B,a}$  be a “question state” and  $|\psi_{\text{ANS},a}\rangle \in \mathcal{H}_{A',a} \otimes \mathcal{H}_{B',a}$  be an “answer state” such that  $|\psi\rangle = |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle$ .*

2. *Let  $Q_a$  be projectors on  $\mathcal{H}_{A,a}$  such that  $\{Q_a \otimes I_{\mathcal{H}_{A',a}}\}$  forms a projective measurement on  $\mathcal{H}_A$  and let  $\{A_b^a\}_{b \in \mathcal{B}}$  and  $\{B_b^a\}_{b \in \mathcal{B}}$  be matrices acting on  $\mathcal{H}_{A',a}$ .*
3. *Let  $D$  be the distribution on  $\mathcal{A}$  obtained by measuring  $|\psi\rangle$  using  $\{Q_a \otimes I_{\mathcal{H}_{A',a}}\}_{a \in \mathcal{A}}$ .*

*Then the following are equivalent:*

- *On average over  $a \sim D$  and with respect to state  $|\psi\rangle$ ,*

$$(I_{\mathcal{H}_{A,a}} \otimes A_b^a) \otimes I_B \approx_\varepsilon (I_{\mathcal{H}_{B,a}} \otimes B_b^a) \otimes I_B.$$

- $(Q_a \otimes A_b^a) \otimes I_B \approx_\varepsilon (Q_a \otimes B_b^a) \otimes I_B$  *on state  $|\psi\rangle$ .*

*Proof.* Expand

$$\begin{aligned} & \sum_{a,b} \|(Q_a \otimes A_b^a - Q_a \otimes B_b^a) \otimes I_B \cdot |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle\|^2 \\ &= \sum_{a,b} \|Q_a \otimes (A_b^a - B_b^a) \otimes I_B \cdot |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle\|^2 \\ &= \sum_{a,b} \|Q_a \otimes I_{\mathcal{H}_{B,a}} |\psi_{\text{QUE},a}\rangle\|^2 \cdot \|(A_b^a - B_b^a) \otimes I_{\mathcal{H}_{B',a}} |\psi_{\text{ANS},a}\rangle\|^2 \\ &= \sum_{a,b} \|Q_a \otimes I_B |\psi\rangle\|^2 \cdot \|(A_b^a - B_b^a) \otimes I_{\mathcal{H}_{B',a}} |\psi_{\text{ANS},a}\rangle\|^2 \\ &= \mathbb{E}_{a \sim D} \sum_b \|(A_b^a - B_b^a) \otimes I_{\mathcal{H}_{B',a}} |\psi_{\text{ANS},a}\rangle\|^2 \\ &= \mathbb{E}_{a \sim D} \sum_b \|I_{\mathcal{H}_{A,a}} \otimes (A_b^a - B_b^a) \otimes I_B \cdot |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle\|^2. \end{aligned}$$

In going from the third to the fourth line we used the fact that

$$\|Q_a \otimes I_{\mathcal{H}_{B,a}} |\psi_{\text{QUE},a}\rangle\|^2 = \|Q_a \otimes I_{\mathcal{H}_{A',a}} \otimes I_B |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle\|^2 = \|Q_a \otimes I_{\mathcal{H}_{A',a}} \otimes I_B |\psi\rangle\|^2.$$

Hence, the first line is  $O(\varepsilon)$  if and only if the last one is. □

**Lemma 8.17.** Let  $\mathcal{A}, \mathcal{B}, \mathcal{C}$  be finite sets. Let  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  be a state, and let  $\{A_{a,b}\}_{a \in \mathcal{A}, b \in \mathcal{B}}$  and  $\{B_{a,c}\}_{a \in \mathcal{A}, c \in \mathcal{C}}$  be POVMs acting on  $\mathcal{H}_A$ . Suppose further that:

1. The measurements approximately commute, i.e.

$$[A_{a,b}, B_{a,c}] \otimes I_B \approx_\delta 0,$$

where the approximation holds with respect to the state  $|\psi\rangle$ .

2. For all  $a \in \mathcal{A}$ , there exist Hilbert spaces  $\mathcal{H}_{A,a}, \mathcal{H}_{A',a}, \mathcal{H}_{B,a}, \mathcal{H}_{B',a}$ , and states  $|\psi_{\text{QUE},a}\rangle \in \mathcal{H}_{A,a} \otimes \mathcal{H}_{B,a}, |\psi_{\text{ANS},a}\rangle \in \mathcal{H}_{A',a} \otimes \mathcal{H}_{B',a}$  such that

$$\begin{aligned} \mathcal{H}_A &= \mathcal{H}_{A,a} \otimes \mathcal{H}_{A',a}, \\ \mathcal{H}_B &= \mathcal{H}_{B,a} \otimes \mathcal{H}_{B',a}, \\ |\psi\rangle &= |\psi_{\text{QUE},a}\rangle \otimes |\psi_{\text{ANS},a}\rangle. \end{aligned}$$

3. For all  $a \in \mathcal{A}$ , there exist projectors  $Q_a$  on  $\mathcal{H}_{A,a}$  and matrices  $\{A_b^a\}_{b \in \mathcal{B}}, \{B_c^a\}_{c \in \mathcal{C}}$  acting on  $\mathcal{H}_{A',a}$  such that  $\{Q_a \otimes I_{\mathcal{H}_{A',a}}\}$  is a projective measurement on  $\mathcal{H}_A$  and

$$A_{a,b} = Q_a \otimes A_b^a, \quad B_{a,c} = Q_a \otimes B_c^a.$$

Then

$$[I_{\mathcal{H}_{A,a}} \otimes A_b^a, I_{\mathcal{H}_{A,a}} \otimes B_c^a] \otimes I_B \approx_\delta 0,$$

on average over  $a \sim D$  where  $D$  is the distribution on  $\mathcal{A}$  obtained by measuring  $|\psi\rangle$  using  $\{Q_a \otimes I_{\mathcal{H}_{A',a}}\}_{a \in \mathcal{A}}$ .

*Proof.* The assumptions of the lemma imply that

$$\begin{aligned} (Q_a \otimes A_b^a B_c^a) \otimes I_B &= ((Q_a \otimes A_b^a) \cdot (Q_a \otimes B_c^a)) \otimes I_B && (Q_a \text{ is a projector}) \\ &= (A_{a,b} \cdot B_{a,c}) \otimes I_B && (\text{Item 3}) \\ &\approx_\delta (B_{a,c} \cdot A_{a,b}) \otimes I_B && (\text{Item 1}) \\ &= ((Q_a \otimes B_c^a) \cdot (Q_a \otimes A_b^a)) \otimes I_B && (\text{Item 3}) \\ &= (Q_a \otimes B_c^a A_b^a) \otimes I_B. && (Q_a \text{ is a projector}) \end{aligned}$$

We apply Lemma 8.16 as follows. The set “ $\mathcal{A}$ ” in Lemma 8.16 is the same as  $\mathcal{A}$  here, and the set “ $\mathcal{B}$ ” is the product set  $\mathcal{B} \times \mathcal{C}$  here. The matrices “ $\{A_b^a\}$ ” are  $\{A_b^a B_c^a\}$  here and “ $\{B_b^a\}$ ” are  $\{B_c^a A_b^a\}$  here. We then obtain, on average over  $a \sim D$ ,

$$(I_{\mathcal{H}_{A,a}} \otimes A_b^a B_c^a) \otimes I_B \approx_\delta (I_{\mathcal{H}_{A,a}} \otimes B_c^a A_b^a) \otimes I_B.$$

This implies the conclusion of the lemma.  $\square$

**Lemma 8.18.** Let  $\mathcal{Y}$  be a finite set and for all  $y \in \mathcal{Y}$  let  $\{A_{x,z}^y\}$  be a POVM on  $\mathcal{H}_A$ . Let  $\{B_{x,y,z}\}$  be a projective measurement on  $\mathcal{H}_B$ . Suppose that

$$\sum_{x,y,z} \langle \psi | A_{x,z}^y \otimes B_{x,y,z} | \psi \rangle \geq 1 - \delta. \quad (73)$$

Then with respect to state  $|\psi\rangle$ ,

$$I_A \otimes B_{x,y,z} \approx_\delta A_{x,z}^y \otimes B_{x,y}.$$

*Proof.* Using the fact that  $\{B_{x,y,z}\}$  is projective, we have  $B_{x,y,z} = B_{x,y}B_z$  for all  $x, y, z$ , so that (73) implies

$$\sum_{x,y,z} \langle \psi | A_{x,z}^y \otimes B_{x,y} B_z | \psi \rangle \geq 1 - \delta .$$

Define  $\hat{A}_z = \sum_{x,y} A_{x,z}^y \otimes B_{x,y}$  and  $\hat{B}_z = I_A \otimes B_z$ . The above equation simplifies to

$$\sum_{y,z} \langle \psi | \hat{A}_z \hat{B}_z | \psi \rangle \geq 1 - \delta .$$

This implies that  $\hat{A}_z \approx_\delta \hat{B}_z$  as

$$\sum_z \left\| (\hat{A}_z - \hat{B}_z) | \psi \rangle \right\|^2 = \sum_z \langle \psi | \hat{A}_z^2 + \hat{B}_z^2 | \psi \rangle - 2 \langle \psi | \hat{A}_z \hat{B}_z | \psi \rangle \leq 2\delta ,$$

where the equality uses the fact that  $\hat{A}_z$  and  $\hat{B}_z$  commute. To conclude the proof, we have

$$I_A \otimes B_{x,y,z} = I_A \otimes B_{x,y} B_z \approx_\delta (I_A \otimes B_{x,y}) \sum_{x',y'} A_{x',z}^{y'} \otimes B_{x',y'} = A_{x,z}^y \otimes B_{x,y},$$

where the approximation follows from  $\hat{A}_z \approx_\delta \hat{B}_z$  and Fact 5.19. □

### 8.4.3 Soundness proof

We analyze the soundness of the introspective verifier.

*Proof of the soundness part of Theorem 8.3.* Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier such that  $\mathcal{S}$  is an  $\ell$ -level sampler. Recall the definition of the introspective verifier  $\hat{\mathcal{V}}^{\text{INTRO}} = (\hat{\mathcal{S}}^{\text{INTRO}}, \hat{\mathcal{D}}^{\text{INTRO}})$  corresponding to  $\mathcal{V}$  from Section 8.2. Fix an index  $n \geq 1$  and let  $N = 2^n$ . Let  $R = N^\lambda$  and  $\text{introparams}(R) = (q, m, d)$ , as in Section 8.2.

As in the proof of completeness part of Theorem 8.3, we make the following simplifications. First, we analyze the soundness of the typed introspective verifier  $\hat{\mathcal{V}}^{\text{INTRO}}$ ; the soundness of the detyped verifier  $\mathcal{V}^{\text{INTRO}} = \text{Detype}(\hat{\mathcal{V}}^{\text{INTRO}})$  follows from Lemma 6.18 and the fact that the type set  $\mathcal{T}^{\text{INTRO}}$  has size  $O(\ell)$ . Second, analogously to Remark 8.6 without loss of generality for notational simplicity we consider strategies for questions sampled from  $\hat{\mathcal{S}}^{\text{INTRO}}$  rather than the downsized sampler  $\mathcal{S}^{\text{INTRO}}$ .

Suppose that  $\text{val}^*(\mathcal{V}_n^{\text{INTRO}}) > 1 - \varepsilon$  for some  $0 < \varepsilon < 1$ , and let  $\mathcal{S} = (|\hat{\psi}\rangle, \hat{A}, \hat{B})$  be a strategy for  $\hat{\mathcal{V}}_n^{\text{INTRO}}$  with value at least  $1 - \varepsilon$ . Since  $\mathcal{S}$  has success probability that is strictly positive, the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  does not automatically reject, which means that

$$s(N) \leq N^\lambda . \tag{74}$$

We analyze each of the tests performed by  $\hat{\mathcal{D}}^{\text{INTRO}}$  (see Figure 10) in sequence, and state consequences of each test. We start with Item 1, the Pauli test.

**Lemma 8.19.** *There exists a function  $\delta_1(\varepsilon, R) = a((\log R)^a \varepsilon^b + (\log R)^{-b})$  for universal constants  $a > 1$  and  $0 < b < 1$  such that the following holds. Let  $Q = M \log q$ , where  $M = 2^m$ . There is a projective strategy  $\mathcal{S}' = (|\psi\rangle, A, B)$  for  $\hat{\mathcal{V}}_n^{\text{INTRO}}$  that succeeds with probability at least  $1 - \delta_1$  and furthermore*

$$|\psi\rangle_{AB} = |\text{EPR}_2\rangle_{A'B'}^{\otimes Q} \otimes |\text{AUX}\rangle_{A''B''} \tag{75}$$

for some bipartite state  $|AUX\rangle$ , and for all  $W \in \{X, Z\}$ ,

$$A_x^{\text{PAULI},W} = \sigma_x^W, \quad B_x^{\text{PAULI},W} = \sigma_x^W, \quad (76)$$

where  $\sigma_x^W$  acts on the first  $s(N)$  qubits of player A's share (resp. B's share) of  $|EPR_2\rangle^{\otimes Q}$ .

*Proof.* Given the definition of the type graph  $G^{\text{INTRO}}$ , for  $((\hat{t}_A, \hat{x}_A), (\hat{t}_B, \hat{x}_B))$  sampled according to  $\mu_{\mathcal{S}^{\text{INTRO}},n}$  it holds that  $(\hat{t}_A, \hat{t}_B) \in \mathcal{T}^{\text{PAULI}} \times \mathcal{T}^{\text{PAULI}}$  with constant probability. Therefore, conditioned on the Pauli test, Item 1, being executed,  $\mathcal{S}$  must succeed in the test with probability  $1 - O(\varepsilon)$ .

Observe that conditioned on the test being executed, the distribution of  $((\hat{t}_A, \hat{x}_A), (\hat{t}_B, \hat{x}_B))$  is, by definition, exactly the distribution of questions in the Pauli basis game with parameters qldparams, as described in Section 7.3.1. By Corollary 7.15 it follows that there exists a local isometry  $\phi = \phi_A \otimes \phi_B$  and a state  $|AUX\rangle \in \mathcal{H}_{A''} \otimes \mathcal{H}_{B''}$  such that

$$\|\phi(|\hat{\psi}\rangle) - |EPR_2\rangle^{\otimes Q} \otimes |AUX\rangle\|^2 \leq \delta'(\varepsilon, R), \quad (77)$$

where  $\delta'(\varepsilon, R)$  is an upper bound on  $\delta_{\text{QLD}}(O(\varepsilon), q, m, d)$  that only depends on  $\varepsilon$  and  $R$ , as stated in Lemma 7.17. In addition, defining  $A_{\hat{a}}^{\hat{x}} = \phi_A \hat{A}_{\hat{a}}^{\hat{x}} \phi_A^\dagger$  for all questions  $\hat{x}$  and answers  $\hat{a}$ , for  $W \in \{X, Z\}$  it holds that

$$A_x^{\text{PAULI},W} \otimes I_B \approx_{\delta'(\varepsilon, R)} \sigma_x^W \otimes I_B, \quad (78)$$

and a similar set of equations hold for operators associated with the second player. Using Naimark's theorem as formulated in [JNV<sup>+</sup>20, Theorem 5.1], at the cost of extending the state  $|AUX\rangle$  we may assume that the measurements are projective without loss of generality. Define the strategy  $\mathcal{S}'$  which uses the state  $|EPR_2\rangle^{\otimes Q} \otimes |AUX\rangle$  and measurement operators  $\{A_{\hat{a}}^{\hat{x}}\}$  and  $\{B_{\hat{a}}^{\hat{x}}\}$  for all questions  $\hat{x}$ , except for (PAULI,  $W$ )-type questions where instead the Pauli measurements  $\sigma_{\hat{a}}^W$  are used. Using (77) and (78) the strategy  $\mathcal{S}'$  succeeds in  $\hat{\mathcal{V}}_n^{\text{INTRO}}$  with probability at least  $1 - \delta'(\varepsilon, R)$ .

The claimed bound on  $\delta_1$  follows from the bound given in Lemma 7.17.  $\square$

In the remainder of the proof we analyze the strategy  $\mathcal{S}'$  from Lemma 8.19. We use the following notation conventions:

1. We use indices A and B to label each player's Hilbert space after application of the isometry  $\phi$  from Lemma 8.19.
2. We write  $V$  for the register subspace of  $\mathbb{F}_2^Q$  spanned by  $e_1, \dots, e_{s(N)}$  and  $\bar{V}$  for its complement. (Note that by definition of introparams in Section 7.3.2 it holds that  $s(N) \leq R \leq Q$ , where the first inequality follows from (74).)
3. Whenever we write a Pauli operator  $\sigma_a^W$  the register on which the operator acts should always be clear from context, and is implicit from the space in which the outcome  $a$  ranges.
4. For measurement operators in the introspection game, the variables for the measurement outcomes follow the specification of the "answer key" in Figure 8 (for  $\mathcal{T}^{\text{PAULI}}$ -type questions) and Figure 10 (for all other question types). For example, the measurement operators  $\{A_x^{\text{PAULI},W}\}$  corresponding to question type (PAULI,  $W$ ) are indexed by vectors  $x \in \mathbb{F}_q^Q$  where  $Q = M \log q$  where  $M = 2^m$ . The measurement operators corresponding to question type (INTRO,  $v$ ) for  $v \in \{A, B\}$  are indexed by pairs  $(y, a) \in V \times \{0, 1\}^{\leq 3Q}$ .<sup>34</sup> We often refer to marginalized measurement operators, e.g., the

<sup>34</sup>Technically the answer  $a$  may be a binary string of any length; however, if  $a$  is too long the decider rejects due to the answer length check. Thus we assume without loss of generality that the answer  $a$  is a binary string of length at most  $3Q = 3 \cdot 2^m \cdot \log q$ .

operator  $A_y^{\text{INTRO},v}$  denotes marginalizing  $A_{y,a}^{\text{INTRO},v}$  over all  $a$ . In these cases, the part of the answer that is marginalized over will be clear from context.

5. We use the notation  $\delta$  to denote a function which is polynomial in  $\delta_1$ , although the exact expression may differ from occurrence to occurrence. The polynomial itself may depend on  $\ell$ , but we leave this dependence implicit; due to the use of inductive steps that involve taking the square root of the error  $\ell$  times in sequence (e.g. Lemma 8.24) the exponent generally depends on  $\ell$ .

The next two lemma derive conditions implied by Items 2 and 3 of the checks performed by the decider  $\hat{\mathcal{D}}^{\text{INTRO}}$  described in Figure 10. As these two parts are performed independently for the two possible values of  $v \in \{A, B\}$ , we only discuss the case where  $v = A$ . For notational simplicity, whenever possible we omit  $v$  when referring to the measurement operators. For example  $L$ ,  $A_{y,a}^{\text{INTRO}}$  and  $B_{z,a}^{\text{SAMPLE}}$  are used as shorthand notation for  $L^A$ ,  $A_{y,a}^{\text{INTRO},A}$ , and  $B_{z,a}^{\text{SAMPLE},A}$  respectively.

**Lemma 8.20** (Sampling test, Item 2 of Figure 10). *For each  $k \in \{1, 2, \dots, \ell\}$ ,*

$$I_A \otimes B_z^{\text{SAMPLE}} \simeq_\delta \sigma_z^Z \otimes I_B, \quad (79)$$

$$A_{y_{\leq k}, a}^{\text{INTRO}} \otimes I_B \simeq_\delta I_A \otimes B_{[L_{\leq k}(\cdot)=y_{\leq k}], a}^{\text{SAMPLE}}, \quad (80)$$

where  $z$  ranges over  $V$  and  $y_{\leq k}$  ranges over  $L_{\leq k}(V)$ . Moreover, analogous equations hold with operators acting on the other side of the tensor product.

*Proof.* When  $((\hat{t}_A, \hat{x}_A), (\hat{t}_B, \hat{x}_B))$  is sampled according to  $\mu_{\hat{\mathcal{S}}^{\text{INTRO},n}}$ , each check in Item 2 of Figure 10 is executed with probability  $\Omega(1/\ell)$  (this is due to the number of types in  $\mathcal{T}^{\text{INTRO}}$  and the structure of the type graph  $G^{\text{INTRO}}$ ). Therefore, in each of the checks specified by Items 2a and 2b, the strategy  $\mathcal{S}'$  succeeds with probability at least  $1 - O(\ell\delta)$ , conditioned on the test being executed. Item 2a for  $w = A$  combined with (76) implies (79). Item 2b for  $w = A$ , combined with Fact 5.24, implies (80). The lemma follows from repeating the same argument with the tensor factors interchanged.  $\square$

**Lemma 8.21** (Hiding test, Item 3 of Figure 10). *For each  $k \in \{1, \dots, \ell + 1\}$ ,*

$$A_{y_{<k}, a}^{\text{INTRO}} \otimes I_B \simeq_\delta I_A \otimes B_{y_{<k}, a}^{\text{READ}}, \quad (81)$$

and if  $k \leq \ell$ ,

$$I_A \otimes B_{y_{<k}}^{\text{HIDE}_k} \approx_\delta \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes I_B. \quad (82)$$

Furthermore, for all  $j, k \in \{1, \dots, \ell - 1\}$  such that  $j \leq k$ , we have

$$A_{y_{<j}, y_{\leq j}^\perp}^{\text{HIDE}_k} \otimes I_B \approx_\delta A_{y_{<j}, y_{\leq j}^\perp}^{\text{HIDE}_{k+1}} \otimes I_B. \quad (83)$$

Analogous equations to (81), (82), and (83) hold with operators acting on the other side of the tensor product.

*Proof.* When  $((\hat{t}_A, \hat{x}_A), (\hat{t}_B, \hat{x}_B))$  is sampled according to  $\mu_{\hat{\mathcal{S}}^{\text{INTRO},n}}$ , each check in Item 3 of Figure 10 is executed with probability  $\Omega(1/\ell)$ . Therefore, in each of the checks specified by Items 3a, 3b, and 3c (conditioned on the right types) the strategy  $\mathcal{S}'$  succeeds with probability at least  $1 - O(\ell\delta)$ .

Item 3a for  $w = A$  combined with Fact 5.24 implies (81). Combining Eqs. (79) and (80) with (81) yields

$$I_A \otimes B_{y_{<k}}^{\text{READ}} \approx_\delta \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes I_B. \quad (84)$$

The fact that  $\mathcal{S}'$  is projective and succeeds in Items 3b and 3c of Figure 10 with probability  $1 - O(\ell\delta)$ , along with Item 1 of Fact 5.18, imply (82).

We now establish the ‘‘Furthermore’’ part of the lemma statement. Let  $1 \leq j \leq k \leq \ell - 1$ . Item 3c, Item 1 of Fact 5.18, and Fact 5.24 imply that

$$A_{y < j, y_{\leq j}^{\perp}}^{\text{HIDE}_k} \otimes I_B \approx_{\delta} I_A \otimes B_{y < j, y_{\leq j}^{\perp}}^{\text{HIDE}_{k+1}}. \quad (85)$$

Item 5 and Fact 5.24 imply

$$A_{y < j, y_{\leq j}^{\perp}}^{\text{HIDE}_{k+1}} \otimes I_B \simeq_{\delta} I_A \otimes B_{y < j, y_{\leq j}^{\perp}}^{\text{HIDE}_{k+1}}. \quad (86)$$

This proves (83).

The lemma follows from repeating the same arguments with the tensor factors interchanged.  $\square$

We exploit the tests performed in Item 3 further to show the following lemma.

**Lemma 8.22.** *For all  $k \in \{1, 2, \dots, \ell\}$ ,*

$$I_A \otimes B_{y < k, y_k^{\perp}, x > k}^{\text{HIDE}_k} \approx_{\delta} \left( \sigma_{[L_{<k}^{\perp}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k, y_{<k}^{\perp}}^{\perp}(\cdot)=y_k^{\perp}]}^X \otimes \sigma_{x > k}^X \right) \otimes I_B,$$

and an analogous equation holds with operators acting on the other side of the tensor product.

*Proof.* The proof is by induction on  $k$ . We first show the case  $k = 1$ . Under the distribution  $\mu_{\hat{\mathcal{S}}^{\text{INTRO}}, n}$ , the check in Item 3d of Fig. 10 is executed with probability  $\Omega(1/\ell)$ . That part of the check for  $w = A$  together with Eq. (76) implies that

$$I_A \otimes B_{y_1^{\perp}, x > 1}^{\text{HIDE}_1} \approx_{\delta} \left( \sigma_{[L_{1, y_{<1}^{\perp}}^{\perp}(\cdot)=y_1^{\perp}]}^X \otimes \sigma_{x > 1}^X \right) \otimes I_B. \quad (87)$$

This proves the case for  $k = 1$ . Next we perform the induction step. Assume that the lemma holds for some  $k \in \{1, 2, \dots, \ell - 1\}$ . The check of Item 3c is executed with probability  $\Omega(1/\ell)$ ; using that  $\mathcal{S}'$  succeeds in Item 3c (conditioned on the right types having been sampled) with probability at least  $1 - O(\ell\delta)$ , we have

$$\sum_{y \leq k, y_{k+1}^{\perp}, x > k+1} \langle \psi | A_{y < k, [L_{k+1, y_{\leq k}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}], x > k+1}^{\text{HIDE}_k} \otimes B_{y \leq k, y_{k+1}^{\perp}, x > k+1}^{\text{HIDE}_{k+1}} | \psi \rangle \geq 1 - O(\ell\delta).$$

We now apply Lemma 8.18, choosing the measurements  $A, B$  and outcomes  $x, y, z$  in the lemma as follows:

$$\begin{aligned} \text{‘‘A’’} &: A_{y < k, [L_{k+1, y_{\leq k}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}], x > k+1}^{\text{HIDE}_k}, & \text{‘‘B’’} &: B_{y \leq k, y_{k+1}^{\perp}, x > k+1}^{\text{HIDE}_{k+1}}, \\ \text{‘‘x’’} &: y < k, & \text{‘‘y’’} &: y_k, & \text{‘‘z’’} &: (y_{k+1}^{\perp}, x > k+1). \end{aligned}$$

Note that here  $A$  does not depend on  $y$ , so we use the same  $A$  for all values of  $y$ . Lemma 8.18 with the above choices of parameters implies that

$$\begin{aligned} I_A \otimes B_{y \leq k, y_{k+1}^{\perp}, x > k+1}^{\text{HIDE}_{k+1}} &\approx_{\delta} A_{y < k, [L_{k+1, y_{\leq k}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}], x > k+1}^{\text{HIDE}_k} \otimes B_{y \leq k}^{\text{HIDE}_{k+1}} \\ &\approx_{\delta} \left( \sigma_{[L_{<k}^{\perp}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k+1, y_{<k+1}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}]}^X \otimes \sigma_{x > k+1}^X \right) \otimes B_{y \leq k}^{\text{HIDE}_{k+1}} \\ &\approx_{\delta} \left( \sigma_{[L_{<k}^{\perp}(\cdot)=y_{<k}]}^Z \sigma_{[L_{\leq k}^{\perp}(\cdot)=y_{\leq k}]}^Z \otimes \sigma_{[L_{k+1, y_{<k+1}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}]}^X \otimes \sigma_{x > k+1}^X \right) \otimes I_B, \\ &\approx_0 \left( \sigma_{[L_{<k+1}^{\perp}(\cdot)=y_{<k+1}]}^Z \otimes \sigma_{[L_{k+1, y_{<k+1}^{\perp}}^{\perp}(\cdot)=y_{k+1}^{\perp}]}^X \otimes \sigma_{x > k+1}^X \right) \otimes I_B, \end{aligned}$$



where the input to  $L_{k+1, y_{<k+1}}^\perp(\cdot)$  is  $x_{k+1}$ . The second approximation uses the induction hypothesis, Fact 5.24, Fact 5.18, and Fact 5.19. The third approximation follows from Eq. (82) and Fact 5.19. The fourth approximation follows from the definition of CL functions. This completes the induction.  $\square$

**Lemma 8.23.** For all  $k \in \{1, \dots, \ell\}$ ,

$$I_A \otimes B_{y_{<k}, y_k^\perp}^{\text{READ}} \approx_\delta \left( \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k, y_{<k}}^\perp(\cdot)=y_k^\perp]}^X \right) \otimes I_B. \quad (88)$$

Moreover, analogous equations hold with operators acting on the other side of the tensor product.

*Proof.* Lemma 8.22 and Fact 5.24 imply that

$$A_{y_{<k}, y_k^\perp}^{\text{HIDE}_k} \otimes I_B \approx_\delta \left( \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k, y_{<k}}^\perp(\cdot)=y_k^\perp]}^X \right) \otimes I_B. \quad (89)$$

Since the strategy  $\mathcal{S}'$  succeeds in Item 3b with probability at least  $1 - O(\ell\delta)$  it follows from Fact 5.24 that

$$A_{y_{<\ell}, y_\ell^\perp}^{\text{HIDE}_\ell} \otimes I_B \approx_\delta I_A \otimes B_{y_{<\ell}, y_\ell^\perp}^{\text{READ}}. \quad (90)$$

An inductive argument applied to (83) of Lemma 8.21, combined with Fact 5.24, implies that for all  $1 \leq k \leq \ell$  we have

$$A_{y_{<k}, y_k^\perp}^{\text{HIDE}_k} \otimes I_B \approx_\delta A_{y_{<k}, y_k^\perp}^{\text{HIDE}_\ell} \otimes I_B. \quad (91)$$

Equations (89), (90), and (91), combined with Fact 5.24, then establishes the lemma statement.  $\square$

**Lemma 8.24.** For each  $k \in \{1, 2, \dots, \ell + 1\}$ , there exists a product state  $|\text{ANC}_k\rangle = |\text{ANC}_{k,A}\rangle \otimes |\text{ANC}_{k,B}\rangle \in \mathcal{H}_{A'_k} \otimes \mathcal{H}_{B'_k}$  and, for each  $y_{<k} \in L_{<k}(V)$ , a projective measurement  $\{A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}}\}$  that acts on  $\mathcal{H}_A \otimes \mathcal{H}_{A'_k}$  such that the following holds. First, for all  $y \in V$ , the operator  $A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}}$  acts as identity on the register subspace spanned by basis vectors for the subspace  $V_{<k}(y_{<k})$ , and as a consequence the operator

$$A_{y, a}^{\text{INTRO}, Z_{<k}} = \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} \quad (92)$$

is well-defined. Second, let  $\mathcal{S}_k''$  be the strategy defined as follows. The state is  $|\text{EPR}_2\rangle^{\otimes Q} \otimes |\text{AUX}\rangle \otimes |\text{ANC}_k\rangle$ . The measurements are identical to those in  $\mathcal{S}'$  defined in Lemma 8.19, except that  $\{A_{y, a}^{\text{INTRO}}\}$  is replaced with  $\{A_{y, a}^{\text{INTRO}, Z_{<k}}\}$ . Then  $\mathcal{S}_k''$  succeeds with probability at least  $1 - \delta$  in the game  $\mathcal{V}_n^{\text{INTRO}}$ .

*Proof.* The proof is by induction on  $k$  from 1 to  $\ell + 1$ . The case  $k = 1$  is trivial by setting  $A_{y_{\geq 1}, a}^{\text{INTRO}, y_{<1}} = A_{y, a}^{\text{INTRO}}$  for all  $y, a$ . Assume that for some  $k \in \{1, 2, \dots, \ell\}$  there exist projective measurements  $\{A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}}\}$  for every  $y_{<k}$  and a strategy  $\mathcal{S}_k''$  satisfying the conditions of the lemma statement. We show the statement of the lemma for  $k + 1$ .

**Commutation with Z-basis measurements.** We first prove that on average over  $y_{<k}$ , the measurement operator  $A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}}$ , which comes from the inductive assumption, commutes with the projective measurement  $\{\sigma_{z_k}^Z\}$  where the outcomes  $z_k$  range over the factor space  $V_k(y_{<k})$ .

To do so, we first apply Lemma 5.25 where we choose the measurements ‘‘A’’, ‘‘B’’, and ‘‘C’’ and outcomes ‘‘a’’, ‘‘b’’, and ‘‘c’’ in the lemma as follows:

$$\begin{aligned} \text{‘‘A’’} &: \{A_{y, a}^{\text{INTRO}, Z_{<k}}\}, & \text{‘‘B’’} &: \{B_{z, a}^{\text{SAMPLE}}\}, & \text{‘‘C’’} &: \{\sigma_z^Z\}, \\ \text{‘‘a’’} &: y_{<k}, & \text{‘‘b’’} &: (y_{\geq k}, a), & \text{‘‘c’’} &: z. \end{aligned}$$

To make sense of how the “B” POVM is indexed by “a”, “b”, and “c” as described above, we use the following relabelling: for all  $(z, a)$ , identify  $B_{z,a}^{\text{SAMPLE}}$  with  $B_{y,a,z}^{\text{SAMPLE}}$  where  $y = L(z)$ . Similarly, for the “C” POVM, we identify  $\sigma_z^Z$  with the operator  $\sigma_{y_{<k},z}^Z$  where  $y_{<k} = L_{<k}(z)$ . By applying Lemma 8.20 to  $\mathcal{S}_k''$  (the strategy given by the inductive hypothesis) with “k” in Lemma 8.20 set to  $\ell$ , we have that

$$A_{y,a}^{\text{INTRO},Z_{<k}} \otimes I_B \approx_\delta I_A \otimes B_{y,a}^{\text{SAMPLE}} \quad (93)$$

where  $B_{y,a}^{\text{SAMPLE}} = \sum_{z:L(z)=y} B_{z,a}^{\text{SAMPLE}}$ . Equations (79) and (93) imply that the conditions of Lemma 5.25 are satisfied, and thus we obtain

$$[A_{y,a}^{\text{INTRO},Z_{<k}}, \sigma_z^Z] \otimes I_B \approx_\delta 0 \quad (94)$$

where in the answer summation,  $y$  is a deterministic function of  $z$ . We now apply Lemma 8.17, choosing the measurements “A”, “B”, “Q” and outcomes “a”, “b”, “c” in the lemma as follows:

$$\begin{aligned} \text{“A”} : \{A_{y,a}^{\text{INTRO},Z_{<k}}\}, \quad \text{“B”} : \{\sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{z_k}^Z\}, \quad \text{“Q”} : \{\sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z\}, \\ \text{“a”} : y_{<k}, \quad \text{“b”} : (y_{\geq k}, a), \quad \text{“c”} : z_k. \end{aligned}$$

Here, we write  $\sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{z_k}^Z$  to denote

$$\sum_{\substack{z' \in V: \\ L_{<k}(z')=y_{<k} \\ (z')^V_k(y_{<k})=z_k}} \sigma_{z'}^Z.$$

We choose the Hilbert spaces “ $\mathcal{H}_{A,a}$ ” and “ $\mathcal{H}_{B,a}$ ” as the register subspace  $V_{<k}(y_{<k})$ , and “ $\mathcal{H}_{A',a}$ ” and “ $\mathcal{H}_{B',a}$ ” as the register subspace  $V_{\geq k}(y_{<k})$  tensored with  $\mathcal{H}_{A''} \otimes \mathcal{H}_{B''}$ , the Hilbert space of the state  $|AUX\rangle$ . Thus for every  $y_{<k}$ , the state  $|EPR\rangle^{\otimes Q} \otimes |AUX\rangle$  of the strategy  $\mathcal{S}_k''$  can be decomposed into a tensor product of a “question state” and an “answer state” as follows:

$$\left( |EPR_2\rangle_{V_{<k}(y_{<k})} \right)_{\mathcal{H}_{A,a} \mathcal{H}_{B,a}} \otimes \left( |EPR_2\rangle_{V_{\geq k}(y_{<k})} \otimes |AUX\rangle \right)_{\mathcal{H}_{A',a} \mathcal{H}_{B',a}}.$$

Let  $\mu_{L_{<k}}$  denote the distribution over outcomes  $y_{<k}$  generated by performing the “Q” measurement on the state  $|EPR\rangle^{\otimes Q}$ , which is equivalent to the distribution generated by the following procedure: (1) sample a uniformly random  $z \in V$ ; (2) compute  $y = L(z)$ ; (3) return  $y_{<k}$ . Then, since Equation (94) and the inductive hypothesis about the structure of  $A_{y_{\leq k},a}^{\text{INTRO},Z_{<k}}$  satisfy the conditions of Lemma 8.17, we obtain on average over  $y_{<k} \sim \mu_{L_{<k}}$

$$[A_{y_{\geq k},a}^{\text{INTRO},y_{<k}}, \sigma_{z_k}^Z] \otimes I_B \approx_\delta 0. \quad (95)$$

Here, the measurement outcomes  $z_k$  range over  $V_k(y_{<k})$ .

**Commutation with X-basis measurements.** Next, we first prove that on average over  $y_{<k}$ , the measurement operator  $A_{y_{\geq k},a}^{\text{INTRO},y_{<k}}$  commutes with the projective measurement  $\{\sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k^\perp]}^X\}$  where the outcomes  $y_k^\perp$  are elements of the factor space  $V_k(y_{<k})$ .

We again apply Lemma 5.25, choosing the measurements and outcomes in the lemma as follows:

$$\begin{aligned} \text{“A”} : \{A_{y,a}^{\text{INTRO},Z_{<k}}\}, \quad \text{“B”} : \{B_{y,y_k^\perp,a}^{\text{READ}}\}, \quad \text{“C”} : \{\sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k^\perp]}^X\}, \\ \text{“a”} : y_{<k}, \quad \text{“b”} : (y_{\geq k}, a), \quad \text{“c”} : y_k^\perp. \end{aligned}$$

Equations (81) (with “ $k$ ” in Lemma 8.21 chosen to be  $\ell + 1$ ) and (88) imply the conditions of Lemma 5.25, so we obtain

$$\left[ A_{y,a}^{\text{INTRO}, Z_{<k}} , \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k^\perp]}^X \right] \otimes I_B \approx_\delta 0 . \quad (96)$$

We then apply Lemma 8.17 with the following choice of measurements and outcomes:

$$\begin{aligned} \text{“}A\text{”} : \{ A_{y,a}^{\text{INTRO}, Z_{<k}} \} , \quad \text{“}B\text{”} : \{ \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k^\perp]}^X \} , \quad \text{“}Q\text{”} : \{ \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \} , \\ \text{“}a\text{”} : y_{<k} , \quad \text{“}b\text{”} : (y_{\geq k}, a) , \quad \text{“}c\text{”} : y_k^\perp . \end{aligned}$$

The Hilbert space and state decomposition are the same as in the previous invocation of Lemma 8.17. Equation (96) and the inductive hypothesis satisfy the conditions of Lemma 8.17, and we similarly obtain that on average over  $y_{<k} \sim \mu_{L_{<k}}$ ,

$$\left[ A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} , \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k^\perp]}^X \right] \otimes I_B \approx_\delta 0 , \quad (97)$$

**Applying the Pauli twirl.** The last step is to apply the Pauli twirl to decompose the family of measurements  $\{ A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} \}$  into a tensor product measurement, with the first part of the tensor product measuring the  $k$ -th linear map of  $L$ .

Again applying Lemma 8.20 to  $\mathcal{S}_k''$ , and using Facts 5.24 and 5.18, we obtain that

$$A_{y_{\leq k}}^{\text{INTRO}, Z_{<k}} \otimes I_B \approx_\delta \sigma_{[L_{\leq k}(\cdot)=y_{\leq k}]}^Z \otimes I_B$$

which is equivalent to, by the inductive hypothesis,

$$\sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes A_{y_k}^{\text{INTRO}, y_{<k}} \otimes I_B \approx_\delta \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k]}^Z \otimes I_B . \quad (98)$$

Applying Lemma 8.16 to Equation (98), we conclude that

$$A_{y_k}^{\text{INTRO}, y_{<k}} \otimes I_B \approx_\delta \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k]}^Z \otimes I_B , \quad (99)$$

on average over  $y_{<k} \sim \mu_{L_{<k}}$ .

Now we apply Lemma 8.15 with the following identification:

$$\begin{aligned} \text{“}x\text{”} : y_{<k} , \quad \text{“}y\text{”} : y_k , \quad \text{“}a\text{”} : (y_{>k}, a) , \quad \text{“}L_x\text{”} : L_{k,y_{<k}} , \\ \text{“}U_x\text{”} : V_k(y_{<k}) , \quad \text{“}M_{y,a}^x\text{”} : A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} , \quad \text{“}M_a^{x,y}\text{”} : A_{y_{>k}, a}^{\text{INTRO}, y_{\leq k}} . \end{aligned}$$

The “Consistency” condition is implied by Equation (99) and the “Commutation” conditions are implied by Equations (95) and (97). We obtain that for all  $y_{\leq k}$  there exists POVM measurements  $\{ A_{y_{>k}, a}^{\text{INTRO}, y_{\leq k}} \}$  that act as identity on the register subspace spanned by basis vectors for the subspace  $V_{<k}(y_{<k})$  and, on average over  $y_{<k} \sim \mu_{L_{<k}}$ , we have

$$A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} \otimes I_B \approx_\delta \left( \sigma_{[L_{k,y_{<k}}^\perp(\cdot)=y_k]}^Z \otimes A_{y_{>k}, a}^{\text{INTRO}, y_{\leq k}} \right) \otimes I_B . \quad (100)$$

Using the inductive assumption on the structure of  $A_{y,a}^{\text{INTRO}, Z_{<k}}$  from (92), we get

$$A_{y,a}^{\text{INTRO}, Z_{<k}} \otimes I_B = \left( \sigma_{[L_{<k}(\cdot)=y_{<k}]}^Z \otimes A_{y_{\geq k}, a}^{\text{INTRO}, y_{<k}} \right) \otimes I_B \quad (101)$$

$$\approx_\delta \left( \sigma_{[L_{\leq k}(\cdot)=y_{\leq k}]}^Z \otimes A_{y_{>k}, a}^{\text{INTRO}, y_{\leq k}} \right) \otimes I_B . \quad (102)$$

where the second line follows from (100) and Lemma 8.16. By (102), replacing the projective measurement  $\{A_{y,a}^{\text{INTRO}, Z_{<k}}\}$  with the POVM

$$\{\sigma_{[L_{\leq k}(\cdot)=y_{\leq k}]}^Z \otimes A_{y_{>k},a}^{\text{INTRO}, y_{\leq k}}\}$$

in the strategy  $\mathcal{S}_k''$  results in a strategy that succeeds with probability at least  $1 - \delta$ . To show that  $\{A_{y_{>k},a}^{\text{INTRO}, y_{\leq k}}\}$  can furthermore be turned into a projective measurement we use Naimark's theorem as formulated in [JNV<sup>+</sup>20, Theorem 5.1]. In particular, we see that the state  $|\text{ANC}'\rangle$  it produces is a product state with no entanglement. This yields a strategy  $\mathcal{S}_{k+1}''$  with ancilla state  $|\text{ANC}_{k+1}\rangle = |\text{ANC}_k\rangle|\text{ANC}'\rangle$ , which establishes the induction hypothesis for  $k + 1$ . This completes the proof of Lemma 8.24.  $\square$

Taking  $k = \ell + 1$  in Lemma 8.24 we obtain a strategy  $\mathcal{S}'' = \mathcal{S}_{\ell+1}''$  with value  $1 - \delta$  in which, given question (INTRO, A), player A performs the measurement

$$\{\sigma_{[L^A(\cdot)=y]}^Z \otimes A_a^{\text{INTRO}, y}\}, \quad (103)$$

for a family of measurements  $\{A_a^{\text{INTRO}, y}\}$  acting on the state  $|\text{EPR}\rangle_{\overline{V}} \otimes |\text{AUX}\rangle \otimes |\text{ANC}\rangle$  where  $|\text{ANC}\rangle = |\text{ANC}_{\ell+1}\rangle$  is unentangled. An analogous argument for Player B's measurements shows that we may additionally assume Player B responds to the question (INTRO, B) using the measurement

$$\{\sigma_{[L^B(\cdot)=y]}^Z \otimes B_a^{\text{INTRO}, y}\}, \quad (104)$$

for a family of measurements  $\{B_a^{\text{INTRO}, y}\}$  acting on the state  $|\text{EPR}\rangle_{\overline{V}} \otimes |\text{AUX}\rangle \otimes |\text{ANC}'\rangle$ , where  $|\text{ANC}'\rangle$  is unentangled. Summarizing, the strategy  $\mathcal{S}''$  uses the state

$$|\theta\rangle = |\text{EPR}\rangle^{\otimes Q} \otimes |\text{AUX}\rangle \otimes |\text{ANC}\rangle \otimes |\text{ANC}'\rangle,$$

and the measurements given by Equations (103) and (104).

To conclude the proof of the soundness part of the theorem we analyze Item 4 in Figure 10. The test in Item 4 is executed with probability  $\Omega(1/\ell)$ , so the strategy  $\mathcal{S}''$  succeeds with probability at least  $1 - \delta$  in that test, conditioned on the right types (here we absorb factors of  $O(\ell)$  into  $\delta$ ). Using (103) and (104), conditioned on the test being executed the distribution of the part  $(y_A, y_B)$  of the players' answers in the test is exactly the distribution  $\mu_{S,N}$  associated with game  $\mathcal{V}_N$ . As a result, the strategy which uses the state  $|\text{EPR}\rangle_{\overline{V}} \otimes |\text{AUX}\rangle \otimes |\text{ANC}\rangle \otimes |\text{ANC}'\rangle$  and measurements  $\{A_a^{\text{INTRO}, y}\}, \{B_a^{\text{INTRO}, y}\}$  succeeds with probability at least  $1 - \delta$  in the game  $\mathcal{V}_N$ . Thus,  $\text{val}^*(\mathcal{V}_N) \geq 1 - \delta$ . It remains to show that  $\delta$  has the required form. As  $\delta = \text{poly}(\delta')$ ,  $\delta'(\varepsilon, R) = a'((\log R)^{a'} \varepsilon^{b'} + (\log R)^{-b'})$ , and  $R = (2^n)^\lambda$ , there is a constant  $C \geq 1$  such that

$$\begin{aligned} \delta(\varepsilon, n) &\leq C \left( a' \left( (\lambda n)^{a'} \varepsilon^{b'} + (\lambda n)^{-b'} \right) \right)^{\frac{1}{C}} \\ &\leq C (a')^{1/C} \left( (\lambda n)^{a'/C} \cdot \varepsilon^{b'/C} + (\lambda n)^{-b'/C} \right) \\ &\leq a \left( (\lambda n)^a \cdot \varepsilon^b + (\lambda n)^{-b} \right), \end{aligned}$$

for  $a = \max\{C(a')^{1/C}, a'/C\}$ , and  $b = b'/C$ , where the second line uses the inequality  $(x + y)^{1/C} \leq x^{1/C} + y^{1/C}$  for  $x, y > 0$  and  $C \geq 1$ .

This establishes the soundness part of the theorem.

To show the entanglement bound, we observe that local isometries do not change the Schmidt rank of a state. Define  $|\psi'\rangle = \phi(|\psi\rangle) \otimes |\text{ANC}\rangle \otimes |\text{ANC}'\rangle$ . Since the strategy  $(|\psi'\rangle, \{A_a^{\text{INTRO}, y}\}, \{B_a^{\text{INTRO}, y}\})$  is

$\delta$ -close to  $(|\theta\rangle, \{A_a^{\text{INTRO}, y}\}, \{B_a^{\text{INTRO}, y}\})$  which has value  $1 - \delta$ , the strategy  $(|\psi'\rangle, \{A_a^{\text{INTRO}, y}\}, \{B_a^{\text{INTRO}, y}\})$  has value at least  $1 - 2\delta$  in the game  $\mathcal{V}_N$ , and therefore the Schmidt rank of  $\phi(|\psi\rangle)$  (and thus of  $|\psi\rangle$ ) must be at least  $\mathcal{E}(\mathcal{V}_N, 1 - 2\delta)$ . Here, we use the fact that ancilla states are product states and therefore have Schmidt rank 1.

Moreover, recall that  $\phi(|\psi\rangle)$  is  $\delta$ -close to  $|\text{EPR}_2\rangle^{\otimes Q} \otimes |\text{AUX}\rangle$  whose Schmidt coefficients are all at most  $2^{-Q/2}$ . For any bipartite state  $|a\rangle$  with Schmidt rank at most  $R$  and  $|b\rangle$  whose Schmidt coefficients are all at most  $\beta$ , it follows from the Cauchy-Schwarz inequality that  $|\langle a|b\rangle|^2 \leq R\beta^2$ . Therefore the Schmidt rank of  $\phi(|\psi\rangle)$  (and thus of  $|\psi\rangle$ ) is at least

$$(1 - \delta)^2 \cdot 2^Q \geq (1 - \delta)^2 \cdot 2^{N^\lambda},$$

where we used that  $Q = M \log q \geq R$  (using the canonical parameter settings of Definition 7.16) and  $\delta \geq \|\phi(|\psi\rangle) - |\theta\rangle\|^2 = 2 - 2\Re\langle\theta|\phi(|\psi\rangle)\rangle$ . Combining the two lower bounds on the Schmidt rank of  $|\psi\rangle$  shows the desired lower bound on  $\mathcal{E}(\hat{\mathcal{V}}_n^{\text{INTRO}}, 1 - \varepsilon)$ .  $\square$

## 9 Oracularization

### 9.1 Overview

In this section we introduce the *oracularization* transformation. At a high level, the oracularization  $\mathfrak{G}^{\text{ORAC}}$  of a nonlocal game  $\mathfrak{G}$  is intended to implement the following: one player (called the *oracle player*) is supposed to receive questions  $(x, y)$  meant for both players in the original game  $\mathfrak{G}$ , and the other player (called the *isolated player*) only receives either  $x$  or  $y$  (but not both), along with a label indicating which player in the original game the question is associated with (we refer to such players as the *original* players, e.g. “original A player” and “original B player”). The oracle player is supposed to respond with an answer pair  $(a, b)$ , and the other player is supposed to respond with an answer  $c$ . The oracle and isolated player win the oracularized game if  $(x, y, a, b)$  satisfies the predicate of the original game  $\mathfrak{G}$  and the isolated player’s answer is consistent with the oracle player’s answer.

The oracularization step is needed in preparation to the next section, in which we perform answer reduction on the introspection game. To implement answer reduction we need at least one player to be able to compute a proof, in the form of a PCP, that the decider of the original game would have accepted the questions  $(x, y)$  and answers  $(a, b)$ . This requires the player to have access to both questions, and be able to compute both answers.

### 9.2 Oracularizing normal form verifiers

Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier such that  $\mathcal{S}$  is an  $\ell$ -level sampler for some  $\ell \geq 0$ . We specify the *typed oracularized verifier*  $\hat{\mathcal{V}}^{\text{ORAC}} = (\hat{\mathcal{S}}^{\text{ORAC}}, \hat{\mathcal{D}}^{\text{ORAC}})$  associated with  $\mathcal{V}$  as follows.

**Sampler.** Define the type set  $\mathcal{T}^{\text{ORAC}} = \{\text{ORACLE}, \text{A}, \text{B}\}$ . (In the remainder of this section we refer to the types in  $\mathcal{T}^{\text{ORAC}}$  as *roles*.) Define the type graph  $G^{\text{ORAC}}$  that is the complete graph on vertex set  $\mathcal{T}^{\text{ORAC}}$  (including self-loops on all vertices). Define the  $\mathcal{T}^{\text{ORAC}}$ -type sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  as follows. For a fixed index  $n \in \mathbb{N}$ , let  $V$  be the ambient space of  $\mathcal{S}$  and  $L^w$  for  $w \in \{\text{A}, \text{B}\}$  be the pair of CL functions of  $\mathcal{S}$  on index  $n$ .

Define two  $\mathcal{T}^{\text{ORAC}}$ -typed families of CL functions  $\{L_t^w : V \rightarrow V\}$ , for  $w \in \{\text{A}, \text{B}\}$  and  $t \in \mathcal{T}^{\text{ORAC}}$ , as follows:

$$L_t^w = \begin{cases} L^t & \text{if } t \in \{\text{A}, \text{B}\}, \\ \text{Id} & \text{if } t = \text{ORACLE}. \end{cases}$$

In other words, if a player gets the type  $t \in \{\text{A}, \text{B}\}$ , then they get the question that original player  $t$  would have received in the game played by  $\mathcal{V}_n$ . If they get type  $t = \text{ORACLE}$ , then they get the entire seed  $z$  that is used by the sampler  $\mathcal{S}$ , from which they can compute both  $L^{\text{A}}(z)$  and  $L^{\text{B}}(z)$ , the pair of questions sampled for the players in game  $\mathcal{V}_n$ .

By definition, the sampler distribution  $\mu_{\hat{\mathcal{S}}^{\text{ORAC}}, n}$  has the following properties.

1. Conditioned on both players receiving the ORACLE role, both players receive  $z$  for a uniformly random  $z \in V$ .
2. Conditioned on both players receiving the isolated player role, the player(s) with role A (respectively, B) receives  $L^{\text{A}}(z)$  (respectively,  $L^{\text{B}}(z)$ ) for a uniformly random  $z \in V$ .
3. Conditioned on player  $w \in \{\text{A}, \text{B}\}$  receiving the ORACLE role and player  $\bar{w}$  receiving the isolated player role, their question tuple is distributed according to  $((\text{ORACLE}, z), (v, L^v(z)))$  if  $w = \text{A}$  and

$((v, L^v(z)), (\text{ORACLE}, z))$  if  $w = \text{B}$ , where  $z \in V$  is uniformly random and  $v$  indicates the role of player  $\bar{w}$ .

**Decider.** The typed decider  $\hat{\mathcal{D}}^{\text{ORAC}}$  is specified in Figure 12.

Input to decider  $\hat{\mathcal{D}}^{\text{ORAC}}$ :  $(n, t_A, x_A, t_B, x_B, a_A, a_B)$ . For  $w \in \{A, B\}$ , if  $t_w = \text{ORACLE}$ , then parse  $a_w$  as a pair  $(a_{w,A}, a_{w,B})$ . Perform the following steps sequentially.

1. **(Game check).** For all  $w \in \{A, B\}$ , if  $t_w = \text{ORACLE}$ , then compute  $x_{w,v} = L^v(x_w)$  for  $v \in \{A, B\}$ . If  $\mathcal{D}$  rejects  $(n, x_{w,A}, x_{w,B}, a_{w,A}, a_{w,B})$ , then reject.
2. **(Consistency checks).**
  - (a) If  $t_A = t_B$  and  $a_A \neq a_B$ , then reject.
  - (b) If for some  $w \in \{A, B\}$ ,  $t_w = \text{ORACLE}$ ,  $t_{\bar{w}} \in \{A, B\}$ , and  $a_{w,t_{\bar{w}}} \neq a_{\bar{w}}$ , then reject.
3. Accept if none of the preceding steps rejects.

Figure 12: Specification of the typed decider  $\hat{\mathcal{D}}^{\text{ORAC}}$ .

### 9.3 Completeness and complexity of the oracularized verifier

We determine the complexity of the oracularized verifier and establish the completeness property.

**Theorem 9.1** (Completeness and complexity of the oracularized verifier). *Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier. Let  $\hat{\mathcal{V}}^{\text{ORAC}} = (\hat{\mathcal{S}}^{\text{ORAC}}, \hat{\mathcal{D}}^{\text{ORAC}})$  be the corresponding typed oracularized verifier. Then the following hold.*

- **(Completeness)** For all  $n \in \mathbb{N}$ , if  $\mathcal{V}_n$  has a PCC strategy of value 1, then  $\hat{\mathcal{V}}_n^{\text{ORAC}}$  has an SPCC strategy of value 1.
- **(Sampler complexity)** The sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  depends only on  $\mathcal{S}$  (and not on  $\mathcal{D}$ ). Moreover, the time complexity of  $\hat{\mathcal{S}}^{\text{ORAC}}$  satisfies

$$\text{TIME}_{\hat{\mathcal{S}}^{\text{ORAC}}}(n) = O(\text{TIME}_{\mathcal{S}}(n)),$$

Furthermore, if  $\mathcal{S}$  is an  $\ell$ -level sampler, then  $\hat{\mathcal{S}}^{\text{ORAC}}$  is a  $\ell$ -level typed sampler.

- **(Decider complexity)** The time complexity of  $\hat{\mathcal{D}}^{\text{ORAC}}$  satisfies

$$\text{TIME}_{\hat{\mathcal{D}}^{\text{ORAC}}}(n) = \text{poly}(\text{TIME}_{\mathcal{D}}(n), \text{TIME}_{\mathcal{S}}(n)).$$

- **(Efficient computability)** There is a Turing machine `ComputeOracleVerifier` which takes as input  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  and returns  $\hat{\mathcal{V}}^{\text{ORAC}} = (\hat{\mathcal{S}}^{\text{ORAC}}, \hat{\mathcal{D}}^{\text{ORAC}})$  in time  $\text{poly}(|\mathcal{V}|)$ .

**Remark 9.2.** Unlike with the Introspection transformation, we do not detype the oracularized verifier  $\hat{\mathcal{V}}^{\text{ORAC}}$ ; this is because the analysis of the Answer Reduction transformation in Section 10 directly reduces to the analysis of the typed oracularized verifier  $\hat{\mathcal{V}}^{\text{ORAC}}$  defined here.

*Proof.* We analyze the completeness and complexity properties of the typed verifier  $\hat{\mathcal{V}}^{\text{ORAC}}$ .

**Completeness.** For  $n \in \mathbb{N}$ , let  $\mathcal{S} = (|\psi\rangle, A, B)$  be a PCC strategy for  $\mathcal{V}_n$  with value 1. Consider the following symmetric strategy  $\mathcal{S}^{\text{ORAC}} = (|\psi\rangle, M)$  for  $\hat{\mathcal{V}}_n^{\text{ORAC}}$ . Depending on the role received, each player performs the following:

1. Suppose the player receives role  $v \in \{A, B\}$  and question  $x$ . Then the player performs the measurement that player  $v$  would on question  $x$  according to strategy  $\mathcal{S}$  to obtain outcome  $a$  (either  $\{A_a^x\}$  or  $\{B_a^x\}$ , depending on  $v$ ). The player replies with  $a$ .
2. Suppose the player receives role  $v = \text{ORACLE}$  and question  $x$ . The player first computes  $y_w = L^w(x)$  for  $w \in \{A, B\}$  where for  $w \in \{A, B\}$ ,  $L^w$  is the CL functions of  $\mathcal{S}$  corresponding to player  $w$ . Then, the player measures using the POVM  $\{M_{a_A, a_B}^{\text{ORACLE}, x}\}$  where

$$M_{a_A, a_B}^{\text{ORACLE}, x} = B_{a_B}^{y_B} A_{a_A}^{y_A}. \quad (105)$$

The projectors  $A_{a_A}^{y_A}$  and  $B_{a_B}^{y_B}$  commute because  $(y_A, y_B)$  is distributed according to  $\mu_{\mathcal{S}, n}$  (over the choice of  $x$ ) and  $\mathcal{S}$  is a commuting strategy for  $\mathcal{V}_n$ . Thus  $M_{a_A, a_B}^{\text{ORACLE}, x}$  is a projector. The player replies with  $(a_A, a_B)$ .

The strategy  $\mathcal{S}^{\text{ORAC}}$  is symmetric and projective by construction, and consistency follows from the consistency of  $\mathcal{S}$ . We now argue that the strategy is commuting and has value 1 in the game  $\hat{\mathcal{V}}_n^{\text{ORAC}}$ . We consider all possible pairs of roles.

1. (Oracle, isolated) Suppose without loss of generality that player  $w = A$  gets the ORACLE role and player  $\bar{w} = B$  gets the isolated player B role. Then player  $w$  gets question  $x$  and player  $\bar{w}$  gets question  $L^{\bar{w}}(x)$ , where  $x$  is uniformly sampled from  $V$ . The oracle player computes  $y_v = L^v(x)$  for all  $v \in \{A, B\}$ . Notice that  $(y_A, y_B)$  is distributed according to  $\mu_{\mathcal{S}, n}$ . The two players return  $((a_A, a_B), a'_B)$  with probability

$$\begin{aligned} \langle \psi | M_{a_A, a_B}^{\text{ORACLE}, x} \otimes B_{a'_B}^{y_B} | \psi \rangle &= \langle \psi | B_{a_B}^{y_B} A_{a_A}^{y_A} \otimes B_{a'_B}^{y_B} | \psi \rangle \\ &= \langle \psi | A_{a_A}^{y_A} \otimes B_{a_B}^{y_B} B_{a'_B}^{y_B} | \psi \rangle \\ &= \delta_{a_B, a'_B} \langle \psi | A_{a_A}^{y_A} \otimes B_{a_B}^{y_B} | \psi \rangle, \end{aligned}$$

where the first equality uses the definition of  $M_{a_A, a_B}^{\text{ORACLE}, x}$  from Eq. (105), the second equality uses the consistency of  $\mathcal{S}$ , and the third equality uses the projectivity of  $\mathcal{S}$ . Notice that when  $a_B = a'_B$ , this is exactly the probability of obtaining answers  $(a_A, a_B)$  when player A and player B get question pair  $(y_A, y_B)$  in the game  $\mathcal{V}_n$ . Since  $\mathcal{S}$  is value-1, the answers satisfy the decision procedure of  $\mathcal{V}_n$  with probability 1. Thus the oracle's answers pass the ‘‘Game check’’ of the oracularized decider with certainty, and furthermore the oracle's answers are consistent with the isolated player's answers and thus pass the ‘‘Consistency check’’ with certainty as well.

Commutativity of  $M_{a_A, a_B}^{\text{ORACLE}, x}$  and  $B_{a_B}^{y_B}$  follows from the commutativity of  $\mathcal{S}$  for the game  $\mathcal{V}_n$ .

2. (Both oracle) If both players get the ORACLE role, then both players receive the same question  $x \in V$ . Using a similar analysis as for the previous item, the players return the same answer pair (thus passing the ‘‘Consistency check’’) and pass the ‘‘Game check’’. Both players' measurements commute because they are identical.



3. (Both isolated) Suppose that both players receive the same isolated player role (e.g., they both receive the isolated player role A). They then perform the same measurements, which produce the same outcomes due to the consistency of the strategy  $\mathcal{S}$ , and thus they pass the ‘‘Consistency check’’. Otherwise, suppose that one player receives the A role and the other player receives the B role. Then the decider  $\hat{\mathcal{D}}^{\text{ORAC}}$  automatically accepts. Furthermore, their measurements commute because their questions are distributed according to  $\mu_{\mathcal{S},n}$ , and  $\mathcal{S}$  is a commuting strategy with respect to  $\mu_{\mathcal{S},n}$ .

**Complexity.** It is clear from the definition that  $\hat{\mathcal{S}}^{\text{ORAC}}$  depends only on  $\mathcal{S}$ . The time complexity of the sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  is dominated by those of the sampler  $\mathcal{S}$ . The complexity of  $\hat{\mathcal{D}}^{\text{ORAC}}$  is dominated by the complexity of both  $\mathcal{S}$  and  $\mathcal{D}$  and performing consistency checks. The sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  is a  $\max\{\ell, 1\}$ -level sampler because  $\mathcal{S}$  is an  $\ell$ -level sampler and the new CL functions for  $t = \text{ORACLE}$  are 1-level.

**Efficient computability.** The description of  $\hat{\mathcal{S}}^{\text{ORAC}}$  can be computed, in polynomial time, from the description of  $\mathcal{S}$  alone. The description of  $\hat{\mathcal{D}}^{\text{ORAC}}$  can be computed in polynomial time from the descriptions of  $\mathcal{S}$  and  $\mathcal{D}$ . Moreover, in each case the computation amounts to copying the description of  $\mathcal{S}$  and  $\mathcal{D}$  respectively, and adding constant-sized additional instructions. □

## 9.4 Soundness of the oracularized verifier

**Theorem 9.3** (Soundness of the oracularized verifier). *Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier and  $\hat{\mathcal{V}}^{\text{ORAC}} = (\hat{\mathcal{S}}^{\text{ORAC}}, \hat{\mathcal{D}}^{\text{ORAC}})$  be the corresponding typed oracularized verifier. Then there exists a function  $\delta(\varepsilon) = \text{poly}(\varepsilon)$  such that for all  $n \in \mathbb{N}$  the following hold.*

1. If  $\text{val}^*(\hat{\mathcal{V}}_n^{\text{ORAC}}) > 1 - \varepsilon$ , then  $\text{val}^*(\mathcal{V}_n) \geq 1 - \delta(\varepsilon)$ .
2. For all  $\varepsilon > 0$ , we have that

$$\mathcal{E}(\hat{\mathcal{V}}_n^{\text{ORAC}}, 1 - \varepsilon) \geq \mathcal{E}(\mathcal{V}_n, 1 - \delta(\varepsilon))$$

where  $\mathcal{E}(\cdot)$  is as in Definition 5.12.

*Proof.* Fix  $n \in \mathbb{N}$ . Let  $\mathcal{S}^{\text{ORAC}} = (|\psi\rangle, A, B)$  be a strategy for  $\hat{\mathcal{V}}_n^{\text{ORAC}}$  with value  $1 - \varepsilon$  for some  $0 < \varepsilon \leq 1$ . Using Lemma 5.7 we may without loss of generality assume that  $\mathcal{S}^{\text{ORAC}}$  is projective. Let  $(t, x) \in \mathcal{T}^{\text{ORAC}} \times V$  be a question to player  $w = A$ . In the event that  $t = \text{ORACLE}$  (which occurs with probability  $1/3$ ), let  $y_v = L^v(x)$  for each  $v \in \{A, B\}$ . From the consistency check performed by  $\hat{\mathcal{D}}^{\text{ORAC}}$  and item 1 of Fact 5.18, we have that for all  $v \in \{A, B\}$  and on average over  $x$  sampled by  $\hat{\mathcal{S}}^{\text{ORAC}}$ ,

$$A_{a_v}^{\text{ORACLE}, x} \otimes I_B \approx_\varepsilon I_A \otimes B_{a_v}^{v, y_v}. \quad (106)$$

Here, we used that with probability  $1/9$  player  $w = A$  gets the ORACLE role and player  $\bar{w} = B$  gets the isolated player  $v$  role; conditioned on this, player B gets question  $y_v$ .

Using Fact 5.19 on (106), with the C operators from Fact 5.19 chosen as  $\{A_{a_A, a_B}^{\text{ORACLE}, x}\}$  here we get

$$\begin{aligned} A_{a_A, a_B}^{\text{ORACLE}, x} \otimes I_B &= A_{a_A, a_B}^{\text{ORACLE}, x} \cdot (A_{a_B}^{\text{ORACLE}, x} \otimes I_B) \\ &\approx_\varepsilon A_{a_A, a_B}^{\text{ORACLE}, x} \cdot (I_A \otimes B_{a_B}^{B, y_B}) \\ &= A_{a_A, a_B}^{\text{ORACLE}, x} \otimes B_{a_B}^{B, y_B}, \end{aligned} \quad (107)$$

where for the first equality we used that the POVM elements  $\{A_{a_A, a_B}^{\text{ORACLE}, x}\}$  are projective. Using a similar calculation for the second and third approximations below we get

$$\begin{aligned}
A_{a_A, a_B}^{\text{ORACLE}, x} \otimes I_B &\approx_\varepsilon A_{a_A, a_B}^{\text{ORACLE}, x} \otimes B_{a_B}^{\text{B}, y_B} \\
&\approx_\varepsilon A_{a_A}^{\text{ORACLE}, x} \otimes B_{a_B}^{\text{B}, y_B} \\
&\approx_\varepsilon I_A \otimes B_{a_B}^{\text{B}, y_B} B_{a_A}^{\text{A}, y_A} \\
&\approx_\varepsilon I_A \otimes B_{a_A}^{\text{A}, y_A} B_{a_B}^{\text{B}, y_B},
\end{aligned} \tag{108}$$

where the last line uses that starting from  $A_{a_A, a_B}^{\text{ORACLE}, x} \otimes I_B$  we may perform all operations leading to the before-last line while reversing the order in which  $B^{\text{B}}$  and  $B^{\text{A}}$  are introduced.

Using Item 2a of the consistency check, we have that on average over a random  $y = L^{\text{A}}(x) \in L^{\text{A}}(V)$ ,

$$A_a^{\text{A}, y} \otimes I_B \approx_\varepsilon I_A \otimes B_a^{\text{A}, y}. \tag{109}$$

Define, for all  $y \in L^{\text{A}}(V)$ , measurement operators  $\{C_a^y\}_a$  where  $C_a^y = A_a^{\text{A}, y}$ . Similarly, define  $D_a^y = B_a^{\text{B}, y}$ . This defines a strategy  $\mathcal{S} = (|\psi\rangle, C, D)$  for the game  $\mathcal{V}_n$  that we now argue succeeds with high probability.

Let  $x \in V$  be uniformly random. Let  $y_A = L^{\text{A}}(x)$  and  $y_B = L^{\text{B}}(x)$ .

$$\begin{aligned}
A_{a_A, a_B}^{\text{ORACLE}, x} \otimes I_B &\approx_\varepsilon I_A \otimes B_{a_B}^{\text{B}, y_B} B_{a_A}^{\text{A}, y_A} \\
&\approx_\varepsilon A_{a_A}^{\text{A}, y_A} \otimes B_{a_B}^{\text{B}, y_B} \\
&= C_{a_A}^{y_A} \otimes D_{a_B}^{y_B}.
\end{aligned}$$

The first approximation follows from Equation (108). The second approximation follows from Equation (109) and Fact 5.19 (where we let  $C_b^y$  in Fact 5.19 represent  $I \otimes B_{a_B}^{\text{B}, y_B}$ ). The last equality follows from definition of  $C_{a_A}^{y_A}$  and  $D_{a_B}^{y_B}$ . The pair of questions  $(y_A, y_B) \in V \times V$  is distributed according to  $\mu_{\mathcal{S}, n}$ .

The game check part of  $\hat{\mathcal{D}}^{\text{ORAC}}$  succeeds with probability  $1 - O(\varepsilon)$ , which implies that the answer pair  $(a_A, a_B)$  that arises from the measurement  $A_{a_A, a_B}^{\text{ORACLE}, x} \otimes I_B$  is accepted by the decider  $\mathcal{D}$  on question pair  $(y_A, y_B)$  with probability  $1 - O(\varepsilon)$ . This in turn implies that the strategy  $\mathcal{S} = (|\psi\rangle, C, D)$  succeeds with probability  $1 - O(\sqrt{\varepsilon})$  in the game  $\mathcal{V}_n$ . As an additional consequence, the Schmidt rank of  $|\psi\rangle$  must be at least  $\mathcal{E}(\mathcal{V}_n, 1 - O(\sqrt{\varepsilon}))$ .  $\square$

## 10 Answer Reduction

In this section we show how to transform a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  into an “answer reduced” normal form verifier  $\mathcal{V}^{\text{AR}} = (\mathcal{S}^{\text{AR}}, \mathcal{D}^{\text{AR}})$  such that the values of the associated nonlocal games are directly related, yet the answer-reduced verifier’s decision runtime is only polylogarithmic in the answer length of the original verifier (the answer-reduced verifier’s sampling runtime remains polynomially related to the sampling runtime of  $\mathcal{S}$ ). The polylogarithmic dependence is achieved by composing a probabilistically checkable proof (PCP) with the oracularized verifier given in Section 9. This step generalizes the answer reduction technique of [NW19, Part V].

Given index  $n \in \mathbb{N}$ , the answer reduced verifier  $\mathcal{V}^{\text{AR}}$  simulates the oracularization  $\mathcal{V}^{\text{ORAC}}$  of  $\mathcal{V}$  on index  $n$ . To do so, it first samples questions  $x$  and  $y$  using the oracularized sampler  $\mathcal{S}^{\text{ORAC}}$  and distributes them to the players, who compute answers  $a$  and  $b$ . Let us suppose that the first player is assigned the ORACLE role, and parse their question and answer as pairs  $x = (x_A, x_B)$  and  $a = (a_A, a_B)$ , while the second player is an “isolated” player receiving the question  $x_A$  and responding with answer  $b$ . Instead of executing the decider  $\mathcal{D}^{\text{ORAC}}$  on the answers  $(a, b)$ , the verifier  $\mathcal{V}^{\text{AR}}$  asks the first player to compute a PCP  $\Pi$  of  $\mathcal{D}(n, x_A, x_B, a_A, a_B) = 1$ , and the second player to compute an encoding  $g_b$  of answer  $b$ .  $\mathcal{V}^{\text{AR}}$  then requests randomly chosen locations of the proof  $\Pi$  and the encoding  $g_b$ , and executes the PCP verifier on the players’ answers. By the soundness of the PCP,  $\mathcal{V}^{\text{AR}}$  accepts with high probability only if the player’s answers satisfy  $\mathcal{D}(n, x_A, x_B, a_A, a_B) = 1$  and  $b = a_A$ .

There are several challenges that arise when implementing answer reduction. One challenge, already encountered in [NW19], is that we need to ensure the PCP  $\Pi$  computed by the first player can be cross-tested against the encoding  $g_b$  computed by the second player (who doesn’t know the entire structure of the PCP  $\Pi$ ). This was handled in [NW19] by using a special type of PCP called a *probabilistically checkable proof of proximity (PCPP)*, which allows one to efficiently check that a *specific* string  $x$  is a satisfying assignment to a Boolean formula  $\varphi$ , as opposed to simply checking that  $\varphi$  is satisfiable. In a PCPP, an encoding of the specific string  $x$  is provided separately from the proof of satisfiability. The answer reduction scheme of [NW19] was able to use an “off-the-shelf” PCPP in a relatively black-box fashion to handle this.

In our answer reduction scheme, however, there is a further requirement: we need the question distribution of the answer reduced verifier to be conditionally linear. This is necessary to maintain the invariant that the verifier after each step of the compression procedure (introspection, answer reduction, parallel repetition) is a normal form verifier. Unfortunately, simulating the question distributions of off-the-shelf PCPPs with conditionally linear distributions can be quite cumbersome. Instead, we design a bespoke PCP verifier for the protocol whose question distribution is more easily seen to be conditionally linear.

This section is organized as follows. We start with some preliminaries on formulas and encodings in Section 10.1. In Section 10.2 we show how to use the Cook-Levin reduction to reduce the Bounded Halting problem for deciders to a succinct satisfiability problem called Succinct-3SAT. Following this, in Section 10.3, we reduce the Succinct-3SAT instance to an instance of a related problem called Succinct Decoupled 5SAT, which is easier to use in our answer reduction step. Then in Section 10.4 we introduce a PCP for Succinct Decoupled 5SAT. The verifier for the PCP expects a proof consisting of the evaluation tables of low-degree polynomials, including the low-degree encodings of the players’ answers  $a$  and  $b$ . In Section 10.5 we provide the definition of a normal-form verifier  $\mathcal{V}^{\text{AR}}$  that executes the composition of  $\mathcal{V}^{\text{ORAC}}$  with the PCP verifier from Section 10.4. In Section 10.6 we show completeness of the construction and analyze its complexity. In Section 10.7 we prove soundness.

## 10.1 Circuit preliminaries

Recall the definitions pertaining to Turing machines from Section 3.1. In this section it is useful to model computation using *Boolean circuits*. A Boolean circuit is a network of Boolean gates, connected by directed “wires.” Incoming wires encode the input to the circuit, and outgoing wires encode the output. For the most part, we will assume that the output of the circuit is a single bit, corresponding to a single output wire. The in-degree of each gate (the “fan-in”) is restricted to be at most 2, but there is no restriction on the out-degree (the “fan-out”). The *size* of a circuit is the total number of gates and wires that it contains. For a more detailed description, see Section 4.3 of [Pap94].

**Remark 10.1** (Plugging integers into circuits). *Let  $\mathcal{C}$  be a circuit with a single input of length  $n$ . Inputs to  $\mathcal{C}$  are strings  $x \in \{0, 1\}^n$ . In this section, we will also allow  $\mathcal{C}$  to receive inputs  $a \in \{0, 1, \dots, 2^n - 1\}$ . In doing so, we use the convention that a number  $a$  between 0 and  $2^n - 1$  is interpreted as its  $n$ -digit binary encoding  $\text{binary}_n(a)$  when provided as input to a set of  $n$  single-bit wires. In other words,  $\mathcal{C}(a) = \mathcal{C}(x)$ , where  $x = \text{binary}_n(a)$ .*

*More generally, if the circuit  $\mathcal{C}$  has  $k$  different inputs of length  $n_1, \dots, n_k$ , then we can evaluate it on inputs  $a_1 \in \{0, 1, \dots, 2^{n_1} - 1\}, \dots, a_k \in \{0, 1, \dots, 2^{n_k} - 1\}$  as follows:*

$$\mathcal{C}(a_1, \dots, a_k) = \mathcal{C}(x_1, \dots, x_k),$$

where  $x_1 = \text{binary}_{n_1}(a_1), \dots, x_k = \text{binary}_{n_k}(a_k)$ .

A 3SAT formula is a Boolean formula in conjunctive normal form in which at most three literals appear in each clause. More precisely,  $\varphi$  is a 3SAT formula on  $N$  variables  $x_1, x_2, \dots, x_N$  if it has the form  $\bigwedge_{j=1}^m C_j$  and each clause  $C_j$  is the disjunction of at most three literals, where a literal is either a variable  $x_i$  or its negation  $\neg x_i$ . We use  $x_i^o$  to denote the literal  $x_i$  if  $o = 1$  and  $\neg x_i$  if  $o = 0$ .

**Definition 10.2** (Succinct description of 3SAT formulas). Let  $N = 2^n$ , and let  $\varphi$  be a 3SAT formula on  $N$  variables named  $x_0, \dots, x_{N-1}$ . Let  $\mathcal{C}$  be a Boolean circuit with 3 inputs of length  $n$  and three single-bit inputs. Then  $\mathcal{C}$  is a *succinct description* of  $\varphi$  if for each  $i_1, i_2, i_3 \in \{0, 1, \dots, N - 1\}$  and  $o_1, o_2, o_3 \in \{0, 1\}$ ,

$$\mathcal{C}(i_1, i_2, i_3, o_1, o_2, o_3) = 1 \tag{110}$$

if and only if  $x_{i_1}^{o_1} \vee x_{i_2}^{o_2} \vee x_{i_3}^{o_3}$  is a clause in  $\varphi$ . In Equation (110), we use the notation from Remark 10.1.

**Definition 10.3** (Succinct-3SAT problem). The *Succinct-3SAT* problem is the language containing encodings of circuits  $\mathcal{C}$  in which  $\mathcal{C}$  is a succinct description of a satisfiable 3SAT formula  $\varphi$ .

The Tseitin transformation is a mapping from circuits to Boolean formulas. The following summarizes its properties; for an explicit construction, see [NW19, Section 3.8].

**Definition 10.4.** Let  $\mathcal{C}$  be a Boolean circuit with  $n$  inputs and size  $s$ . Then the corresponding *Tseitin formula*  $\mathcal{F}$  is a Boolean formula on  $n + s$  variables with the property that for all  $x \in \{0, 1\}^n$ ,  $\mathcal{C}(x) = 1$  if and only if there exists  $w \in \{0, 1\}^s$  such that  $\mathcal{F}(x, w) = 1$ . The formula  $\mathcal{F}$  has size  $O(s)$  and a description of it is computable from  $\mathcal{C}$  in time  $O(s)$ .

**Definition 10.5.** Let  $\mathcal{F}$  be a Boolean formula over  $m$  variables. The arithmetization  $\mathcal{F}_{\text{arith}}$  over  $\mathbb{F}_q$  is a function  $\mathcal{F}_{\text{arith}} : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  such that

$$\forall x \in \{0, 1\}^m, \quad \mathcal{F}_{\text{arith}}(x) = \mathcal{F}(x). \tag{111}$$

**Proposition 10.6.** *Let  $\mathcal{C}$  be a circuit on  $n$  inputs with size  $s$ . Then the arithmetization  $\mathcal{F}_{\text{arith}}$  of its Tseitin formula is a polynomial over  $\mathbb{F}_q$  on  $n + s$  variables with individual degree at most 2 in each variable. For admissible field sizes  $q$ , a description of the polynomial  $\mathcal{F}_{\text{arith}}$  can be computed in  $\text{poly}(s, \log q)$  time it can be evaluated at a specific point  $z \in \mathbb{F}_q^{n+s}$  in time  $\text{poly}(s, \log q)$ .*

*Proof.* The properties follow by inspecting the construction in Definition 3.27 and 3.28 of [NW19], together with Lemma 3.18. In particular, the degree bound follows by observing that every variable in  $\mathcal{F}$  occurs at most twice, and therefore the arithmetization  $\mathcal{F}_{\text{arith}}$  has individual degree 2.  $\square$

## 10.2 A Cook-Levin theorem for bounded deciders

**Definition 10.7** (Bounded Halting problem). The  $k$ -input *Bounded Halting* problem is the language  $\text{BH}_k$  containing the set of tuples  $(\alpha, T, z_1, \dots, z_k)$  where  $\alpha$  is the description of a  $k$ -input Turing machine,  $T \in \mathbb{N}$ ,  $z_1, \dots, z_k \in \{0, 1\}^*$ , and  $\mathcal{M}_\alpha$  accepts input  $(z_1, \dots, z_k)$  in at most  $T$  time steps.

We begin by defining natural encodings of a decider’s tape alphabet and set of states.

**Definition 10.8** (Decider encodings). Let  $\mathcal{D}$  be a decider with tape alphabet  $\Gamma = \{0, 1, \sqcup\}$  and set of states  $K$ . We will write  $\text{enc}_\Gamma : \Gamma \cup \{\square\} \rightarrow \{0, 1\}^2$  for the function which encodes the elements of  $\Gamma$ , and a special “ $\square$ ” symbol described below, as length-two binary strings in the following manner:

$$\begin{aligned} \text{enc}_\Gamma(0) &= 00, & \text{enc}_\Gamma(1) &= 01, \\ \text{enc}_\Gamma(\sqcup) &= 10, & \text{enc}_\Gamma(\square) &= 11. \end{aligned}$$

In addition, we write  $\text{enc}_K : K \rightarrow \{0, 1\}^\kappa$  for some arbitrary fixed  $\kappa$ -bit encoding of the elements of  $K$ , where  $\kappa = \lceil \log(|K|) \rceil$ .

Now, we give the main result of this section. It states that any decider  $\mathcal{D}$  can be converted into a circuit  $\mathcal{C}$  which succinctly represents a 3SAT formula  $\varphi_{3\text{SAT}}$  that carries out the time  $T$  computation of  $\mathcal{D}$ . In addition,  $\mathcal{C}$  is extremely small—size  $\text{poly} \log(T)$  rather than  $\text{poly}(T)$ .

**Proposition 10.9** (Succinct representation of deciders). *There is an algorithm with the following properties. Let  $\mathcal{D}$  be a decider; let  $n, T, Q$ , and  $\sigma$  be integers with  $Q \leq T$  and  $|\mathcal{D}| \leq \sigma$ , and let  $x$  and  $y$  be strings of length at most  $Q$ . Then on input  $(\mathcal{D}, n, T, Q, \sigma, x, y)$ , the algorithm outputs a circuit  $\mathcal{C}$  on  $3m + 3$  inputs which succinctly describes a 3SAT formula  $\varphi_{3\text{SAT}}$  on  $M = 2^m$  variables. Furthermore,  $\varphi_{3\text{SAT}}$  has the following property:*

- For all  $a, b \in \{0, 1\}^{2T}$ , there exists a  $c \in \{0, 1\}^{M-4T}$  such that  $w = (a, b, c)$  satisfies  $\varphi_{3\text{SAT}}$  if and only if there exist  $a_{\text{prefix}}, b_{\text{prefix}} \in \{0, 1\}^*$  of lengths  $\ell_a, \ell_b \leq T$ , respectively, such that

$$a = \text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{T-\ell_a}) \quad \text{and} \quad b = \text{enc}_\Gamma(b_{\text{prefix}}, \sqcup^{T-\ell_b})$$

and  $\mathcal{D}$  accepts  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  in time  $T$ .

Finally, the following statements hold:

1. The parameter  $m$  controlling the number of inputs to the circuit depends only on  $T$  and  $\sigma$ , and  $m(T, \sigma) = O(\log(T) + \log(\sigma))$ ,
2.  $\mathcal{C}$  has at most  $s(n, T, Q, \sigma) = \text{poly}(\log(n), \log(T), Q, \sigma)$  gates,

3. The algorithm runs in time  $\text{poly}(\log(n), \log(T), Q, \sigma)$ ,
4. Furthermore, explicit values for  $m(T, \sigma)$  and  $s(n, T, Q, \sigma)$  can be computed in time polynomial in  $n, \log(T), Q, \sigma$ .

Proposition 10.9 is essentially the standard fact that Succinct-3SAT is an NEXP-complete language, i.e. that every nondeterministic computation which takes time  $2^n$  can be represented as a Succinct-3SAT instance of size only  $\text{poly}(n)$ . However, it has several peculiarities which requires us to prove it from scratch rather than simply appealing to the NEXP-completeness of Succinct-3SAT. First, we require that the coordinates of  $a$  and  $b$  embed into  $w$  not randomly but as its lexicographically first coordinates (for reasons that are explained below in Section 10.3). Second, we need explicit bounds on how quantities such as the size of  $\mathcal{C}$  relate to quantities such as  $\sigma$ , an upper bound on the description length of  $\mathcal{D}$  in bits.

To prove Proposition 10.9, we follow the standard proof that Succinct-3SAT is NEXP-complete as presented in [Pap94]. This proof observes that the Cook-Levin reduction, which is used to show that 3SAT is NP-complete, produces a 3SAT instance whose clauses follow such a simple pattern that they can be described succinctly using an exponentially-smaller circuit. One key difference in our proof is that we will apply the Cook-Levin reduction directly to the 5-input Turing machine  $\mathcal{D}$ , which by Section 3.1 has 7 tapes; traditional proofs such as the one in [Pap94] would first convert  $\mathcal{D}$  to a single-tape Turing machine  $\mathcal{D}_{\text{single}}$ , and then apply the Cook-Levin reduction for single-tape Turing machines to  $\mathcal{D}_{\text{single}}$ . Though this adds notational overhead to our proof, it allows us to more easily track of which variables in  $\varphi_{3\text{SAT}}$  correspond to the strings  $a$  and  $b$  (see Proposition 10.9 to see what these refer to).

*Proof of Proposition 10.9.* The Cook-Levin reduction considers the *execution tableau* of  $\mathcal{D}$  when run for time  $T$ . The execution tableau contains, for each time  $t \in \{1, \dots, T\}$ , variables describing the state of  $\mathcal{D}$  and the contents of each of its tape cells at time  $t$ . More formally, it consists of the following three sets of variables.

1. For each time  $t \in \{1, \dots, T\}$ , tape  $i \in \{1, \dots, 7\}$ , and tape position  $j \in \{0, 1, \dots, T + 1\}$ , the tableau contains two Boolean-valued variables

$$c_{t,i,j} = (c_{t,i,j,1}, c_{t,i,j,2}) \in \{0, 1\}^2$$

which are supposed to correspond to the contents of the  $j$ -th tape cell on tape  $i$  at time  $t$  according to  $\text{enc}_\Gamma(\cdot)$ . The variables with  $j \in \{0, T + 1\}$  do not correspond to any cell on the tape; rather, the  $j = 0$  variables correspond to the left-boundary of the tape, and the  $j = T + 1$  variables correspond to the right-boundary of the first  $T$  cells on the tape. These are expected to always contain the special boundary symbol “ $\square$ ”, i.e.  $c_{t,i,j}$  should be equal to  $\text{enc}_\Gamma(\square)$  whenever  $j \in \{0, T + 1\}$ . As we will see below, it is convenient to define these so that for each  $t \in \{1, \dots, T\}$ ,  $i \in \{1, \dots, 7\}$ , and  $j \in \{1, \dots, T\}$ , the variable  $c_{t,i,j}$  also has a variable to its left  $c_{t,i,j-1}$  and to its right  $c_{t,i,j+1}$ .

2. For each time  $t \in \{1, \dots, T\}$ , tape  $i \in \{1, \dots, 7\}$ , and tape position  $j \in \{0, 1, \dots, T + 1\}$ , the tableau contains Boolean-valued variables  $h_{t,i,j} \in \{0, 1\}$  which are supposed to indicate whether the  $i$ -th tape head is in cell  $j$  at time  $t$ . For the boundary cells  $j \in \{0, T + 1\}$ , we expect that  $h_{t,i,j} = 0$  for all  $t \in \{1, \dots, T\}$  and  $i \in \{1, \dots, 7\}$ .
3. For each time  $t \in \{1, \dots, T\}$ , the tableau contains  $\kappa$  Boolean-valued variables

$$s_t = (s_{t,1}, \dots, s_{t,\kappa}) \in \{0, 1\}^\kappa$$

which are supposed to correspond to the state of  $\mathcal{D}$  at time  $t$  according to  $\text{enc}_\kappa(\cdot)$ .

Finally, we let  $\mathcal{V}$  denote the set of all of these variables. In other words,

$$\mathcal{V} = \{c_{t,i,j,k}\}_{t,i,j,k} \cup \{h_{t,i,j}\}_{t,i,j} \cup \{s_{t,k}\}_{t,k}.$$

In total, the number of variables in the execution tableau is given by

$$|\mathcal{V}| = O(T^2 + T \cdot \log(|K|)) = O(T^2 + T \cdot \log(|\mathcal{D}|)). \quad (112)$$

The first term in Equation (112) corresponds to the tape cell encodings  $c_{t,i,j}$  and  $h_{t,i,j}$ , and the second term corresponds to the Turing machine state encodings  $s_t$ . The second equality uses the fact that  $|K| \leq |\mathcal{D}|$ .

As stated above, we expect the variables in  $\mathcal{V}$  to correspond to some time- $T$  execution of the decider  $\mathcal{D}$ . However, in general these are just arbitrary  $\{0,1\}$ -valued variables. We now describe a set of constraints placed on these variables which, if satisfied, ensure they *do* indeed correspond to some time- $T$  execution of  $\mathcal{D}$ . These constraints will be split into two categories: (i) the constraints corresponding to the boundary, which ensure that the  $t = 1$  variables are initialized to a valid starting configuration and the  $j \in \{0, T + 1\}$  variables are set according to Items 1 and 2, and (ii) the constraints corresponding to the execution of  $\mathcal{D}$ , which ensure that the variables at each time  $(t + 1)$  follow from the variables at time  $t$  according to the computation of  $\mathcal{D}$ . We start with the boundary constraints, which are simple enough to be described with a 3SAT formula.

**Definition 10.10.** The *boundary formula*  $\varphi_{\text{Boundary}}$  is the 3SAT formula on the variables  $\mathcal{V}$  described as follows. Let  $o = \text{enc}_K(\text{start}) \in \{0,1\}^\kappa$ , where  $\text{start} \in K$  is the start state of  $\mathcal{D}$ . For the  $t = 1$  boundary,  $\varphi_{\text{Boundary}}$  contains the following set of clauses.

Indices	Clauses	
$i \in \{1, \dots, 7\}$	$h_{1,i,1}$	(113)
$i \in \{1, \dots, 7\}, j \neq 1$	$\neg h_{1,i,j}$	(114)
$k \in \{1, \dots, \kappa\}$	$s_{1,k}^{o_k}$	(115)
$i \in \{1, \dots, 7\}, j \in \{1, \dots, T\}$	$\neg c_{1,i,j,1} \vee \neg c_{1,i,j,2}$	(116)
$i \in \{1, \dots, 5\}, j < j' \in \{1, \dots, T\}$	$\neg c_{1,i,j,1} \vee c_{1,i,j',1}$	(117)
$i \in \{6, 7\}, j \in \{1, \dots, T\}$	$c_{1,i,j,1}$ and $\neg c_{1,i,j,2}$	(118)

This is meant to be read as follows: for each row, the ‘‘Indices’’ column specifies the range of the indices that the clauses in the ‘‘Clauses’’ column are quantified over. For example, row (113) specifies that for all  $i \in \{1, \dots, 7\}$ ,  $\varphi_{\text{Boundary}}$  contains the clause  $h_{1,i,1}$ . For the  $j \in \{0, T + 1\}$  boundary,  $\varphi_{\text{Boundary}}$  contains the following set of clauses.

Indices	Clauses	
$t \in \{1, \dots, T\}, i \in \{1, \dots, 7\}, j \in \{0, T + 1\}, k \in \{1, 2\}$	$c_{t,i,j,k}$	(119)

Rows (113) and (114) ensure that at time  $t = 1$ , each tape has exactly one tape head, and it is located on cell  $j = 1$ . Row (115) ensures that at time  $t = 1$ , the state is given by the start state  $\text{start}$ . (Recall the notation  $x_i^o$  to denote the literal  $x_i$  if  $o = 1$  and  $\neg x_i$  if  $o = 0$ .) For the remaining rows, we recall that under the encoding of the tape alphabet,  $\text{enc}_\Gamma(\sqcup) = 10$  and  $\text{enc}_\Gamma(\square) = 11$ . As a result, (i) row (119) ensures that for all times and tapes, the cells  $j \in \{0, T + 1\}$  contain the  $\square$  symbol, (ii) row (116) ensures that for time  $t = 1$ , no cell  $j \notin \{0, T + 1\}$  contains the  $\square$  symbol, and (iii) row (118) ensures that for time  $t = 1$

and tapes 6 and 7, all cells  $j \in \{1, \dots, T\}$  contain the  $\sqcup$  symbol. Finally, row (117) says that for tapes  $i \in \{1, \dots, 5\}$ , if cell  $j$  contains  $\sqcup$ , then every cell  $j' > j$  must contain  $\sqcup$  as well. This means that the five strings encoded by  $c_{1,1}, \dots, c_{1,5}$  each consist of a string of 0's and 1s followed by a string of  $\sqcup$ 's. In short, suppose we write  $n, x, y, a$ , and  $b$  for the prefixes of these strings with no  $\sqcup$ 's. If  $\varphi_{\text{Boundary}}$  is satisfied, then the execution tableau correctly encodes that the tapes of  $\mathcal{D}$  contain inputs  $n, x, y, a$ , and  $b$  at time  $t = 1$ .

Next, we describe the execution constraints. These are more complicated than the boundary constraints, and so we will begin by describing them in terms of a general Boolean circuit known as the *local check circuit*. For any time  $t \in \{1, \dots, T-1\}$  and tape positions  $j_1, \dots, j_7 \in \{1, \dots, T\}$ , the local check circuit can check that the execution tableau properly encodes these tape positions at time  $t+1$  by looking only at the encodings of these tape positions and their neighbors (i.e. the tape positions  $j_i \pm 1$  for each  $i \in \{1, \dots, 7\}$ ) at time  $t$ .

**Definition 10.11.** In this definition, we will define the *local check circuit*  $\mathcal{C}_{\text{Check}}$ . It has the following inputs.

- For each  $i \in \{1, \dots, 7\}$ , it has the eight inputs

$$\begin{array}{ccccccc} c_{\text{Check},0,i,-1} & c_{\text{Check},0,i,0} & c_{\text{Check},0,i,1} & h_{\text{Check},0,i,-1} & h_{\text{Check},0,i,0} & h_{\text{Check},0,i,1} & \\ & c_{\text{Check},1,i,0} & & & h_{\text{Check},1,i,0} & & \end{array} \quad (120)$$

where the  $c$ -inputs are in  $\{0, 1\}^2$  and the  $h$ -inputs are in  $\{0, 1\}$ .

- It has two inputs  $s_{\text{Check},0}, s_{\text{Check},1} \in \{0, 1\}^k$ .

In addition, for each time  $t \in \{1, \dots, T-1\}$  and tape positions  $j_1, \dots, j_7 \in \{1, \dots, T\}$ , we will define a circuit  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  by associating the inputs of  $\mathcal{C}_{\text{Check}}$  with certain variables in  $\mathcal{V}$ . We do this by associating the following eight inputs from  $\mathcal{V}$  with the corresponding variables in Equation (120):

$$\begin{array}{ccccccc} c_{t,i,j_i-1} & c_{t,i,j_i} & c_{t,i,j_i+1} & h_{t,i,j_i-1} & h_{t,i,j_i} & h_{t,i,j_i+1} & \\ & c_{t+1,i,j_i} & & & h_{t+1,i,j_i} & & \end{array}$$

as well as by associating  $s_t$  and  $s_{t+1}$  from  $\mathcal{V}$  with  $s_{\text{Check},0}$  and  $s_{\text{Check},1}$ , respectively.

We first define  $\mathcal{C}_{\text{Check}}$  as a function. In Proposition 10.12 below we give an implementation of this function as a circuit. To do so it is convenient to define a function  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  for each value of  $t, j_1, \dots, j_7$ . It will be clear that each of these is in fact the same function applied to different inputs, and hence this will define  $\mathcal{C}_{\text{Check}}$  as well. Now, the functions  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  perform the following computation.

- Suppose for each  $i \in \{1, \dots, 7\}$ , exactly one of the tape positions  $j_i - 1, j_i$ , and  $j_i + 1$  at time  $t$  contains a tape head. First,  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  computes the transition function of the Turing machine applied to the contents of these 7 tape positions, when the state of  $\mathcal{D}$  at time  $t$  is  $s_{\text{Check},0}$ . This transition function produces the state of  $\mathcal{D}$  and contents of these tape positions at time  $t+1$ , as well as directions to move the 7 tape heads in. Then  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  checks that the variables for the tape positions  $j_1, \dots, j_7$  and the state of  $\mathcal{D}$  at time  $t+1$  match what they should be according to the computed transition function. In addition, if a tape head is marked as moving into a tape cell containing a  $\sqcup$  symbol, that tape head remains in place instead.
- Suppose for each  $i \in \{1, \dots, 7\}$ , none of the tape positions  $j_i - 1, j_i$ , or  $j_i + 1$  contain a tape head at time  $t$ . Then  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  checks that the variables for the tape positions  $j_1, \dots, j_7$  at time  $t+1$  match the variables at time  $t$ .
- Otherwise,  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  accepts.



This defines  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$ , and therefore  $\mathcal{C}_{\text{Check}}$ .

Definition 10.11 shows the utility of introducing the boundary variables  $c_{t,i,0}$  and  $c_{t,i,T+1}$ . The function  $\mathcal{C}_{\text{Check}}$  checks the contents of a cell  $c_{t,i,j}$  at time  $t + 1$  by looking at the cell and its neighbors  $c_{t,i,j-1}, c_{t,i,j+1}$  at time  $t$ . However, those cells with  $j \in \{1, T\}$  only have either a left neighbor or a right neighbor, and so without the boundary variables we'd have to introduce two other local check circuits designed just for these boundary cases. The boundary variables then allow us to use the same local check circuit for all cells.

**Proposition 10.12.** *A circuit  $\mathcal{C}_{\text{Check}}$  of size at most  $\text{poly}(|\mathcal{D}|)$  which computes the function described above can be computed in time  $\text{poly}(|\mathcal{D}|)$ .*

*Proof.* The circuit has  $7 \cdot 12 + 2 \cdot \kappa = 84 + 2\kappa$  total Boolean inputs, giving a total of  $2^{84} \cdot 4^\kappa = O(|K|^2)$  possible input strings. For each possible fixed input string, the circuit  $\mathcal{C}_{\text{Check}}$  checks if the actual input is equal to the fixed input, which takes  $O(\kappa)$  gates, and then it accepts if the fixed input should be accepting, i.e. for each fixed input the correct output is hard-coded in the circuit. Doing so takes  $O(|K|^2 \cdot \kappa)$  gates. Computing  $\mathcal{C}_{\text{Check}}$  requires looping over all possible input strings and checking which ones are accepting or rejecting. This requires computing the transition function of  $\mathcal{D}$ , a task which takes time  $\text{poly}(|\mathcal{D}|)$ . The proposition follows by noting that  $|K| \leq |\mathcal{D}|$ .  $\square$

**Proposition 10.13.** *Suppose that the execution tableau satisfies  $\varphi_{\text{Boundary}}$  and the circuit  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  for each  $t \in \{1, \dots, T-1\}$  and  $j_1, \dots, j_7 \in \{1, \dots, T\}$ . Then the execution tableau correctly encodes the execution of  $\mathcal{D}$  on input  $(n, x, y, a, b)$  when run for time  $T$ .*

*Proof.* To show this, we will show for each time  $t \in \{1, \dots, T\}$  that the variables in the execution tableau corresponding to time  $t$  correctly encode the state of  $\mathcal{D}$  and the contents of the seven tapes at time  $t$ . The proof is by induction on  $t$ . The base case of  $t = 1$  follows from the tableau satisfying  $\varphi_{\text{Boundary}}$ .

Next we perform the induction step. Assuming the statement holds for time  $t \in \{1, \dots, T-1\}$ , we will show it holds for time  $t + 1$  as well. Let  $i^* \in \{1, \dots, 7\}$  be a tape, and consider a tape position  $j_{i^*} \in \{1, \dots, T\}$ . We will show that the variables correctly encode the contents of this tape position at time  $t + 1$ . Suppose one of the tape positions  $j_{i^*} - 1, j_{i^*},$  or  $j_{i^*} + 1$  at time  $t$  has a tape head. For each of the other tapes  $i \neq i^*$ , we select a tape position  $j_i$  such that either  $j_i - 1, j_i,$  or  $j_i + 1$  has a tape head at time  $t$ . (These positions are guaranteed to exist since each tape has exactly one tape head.)

By the induction hypothesis, for each tape  $i \in \{1, \dots, 7\}$  the variables corresponding to tape cells  $j_i - 1, j_i,$  and  $j_i + 1$  correctly encode the contents of these cells at time  $t$ . By assumption,  $\mathcal{C}_{\text{Check},t,j_1,\dots,j_7}$  evaluates to 1. In this case, it calculates the transition function of  $\mathcal{D}$  to compute the contents of the tape cells  $j_1, \dots, j_7$  at time  $t + 1$  and checks that the corresponding variables encode these contents. As a result, the tape position  $j_{i^*}$  on tape  $i^*$  is correctly encoded. In addition, it computes the state of  $\mathcal{D}$  at time  $t + 1$  and checks that the corresponding variables encode this state. This completes the induction step. The case when none of the tape positions  $j_{i^*} - 1, j_{i^*},$  and  $j_{i^*} + 1$  at time  $t$  contain a tape head follows similarly. Finally, the variables for all tape positions  $j \in \{0, T+1\}$  are correctly encoded due to  $\varphi_{\text{Boundary}}$  being satisfied.  $\square$

Proposition 10.13 gives a set of constraints that ensure the execution tableau properly encodes the execution of  $\mathcal{D}$ . Our next step will be to convert these constraints into a single 3SAT formula. This entails transforming each circuit  $\mathcal{C}_{\text{Check},t,i_1,\dots,i_7}$  into a 3SAT formula. We do so using the following reduction.

**Proposition 10.14 (Circuit-to-3SAT).** *There is an algorithm which, on input a size- $r$  circuit  $\mathcal{C}$  on variables  $x \in \{0, 1\}^n$ , runs in time  $\text{poly}(r)$  and outputs a 3SAT formula  $\varphi$  on variables  $x \in \{0, 1\}^n$  and  $y \in \{0, 1\}^r$  with  $O(r)$  clauses such that for all  $x$ ,  $\mathcal{C}(x) = 1$  if and only if there exists a  $y$  such that  $x$  and  $y$  satisfy  $\varphi$ .*

*Proof.* This is the textbook circuit-to-3SAT reduction. For each gate  $i \in \{1, \dots, r\}$ , the algorithm introduces a variable  $y_i \in \{0, 1\}$ , so that the total collection of variables is  $\mathcal{V}_{\text{total}} = \{x_i\}_{i \in \{1, \dots, n\}} \cup \{y_i\}_{i \in \{1, \dots, r\}}$ . Consider gate  $i \in \{1, \dots, r\}$ , and let  $z_1, z_2 \in \mathcal{V}_{\text{total}}$  be the pair of variables feeding into it. If gate  $i$  is an AND gate, then  $\varphi$  includes the constraints

$$(\neg z_1 \vee \neg z_2 \vee y_i), \quad (z_1 \vee z_2 \vee \neg y_i), \quad (z_1 \vee \neg z_2 \vee \neg y_i), \quad (\neg z_1 \vee z_2 \vee \neg y_i). \quad (121)$$

These constraints are satisfied if and only if  $y_i = z_1 \wedge z_2$ . The case of gate  $i$  being an OR gate follows similarly. Finally,  $\varphi$  includes the constraint  $(y_{i^*})$ , where  $i^* \in \{1, \dots, r\}$  is the output gate. It follows that  $x, y$  satisfy  $\varphi$  if and only if  $\mathcal{C}(x) = 1$  and for each  $i \in \{1, \dots, r\}$ ,  $y_i$  is the value computed by gate  $i$  in circuit  $\mathcal{C}$  on input  $x$ . In total,  $\varphi$  has  $4r + 1$  clauses and is computable in time  $\text{poly}(r)$ .  $\square$

We now apply the algorithm from Proposition 10.14 to  $\mathcal{C}_{\text{Check}}$ , which by Proposition 10.12 has size  $r \leq \text{poly}(|\mathcal{D}|)$ . This produces a 3SAT formula  $\varphi_{\text{Check}}$  on the variables in  $\mathcal{C}_{\text{Check}}$  plus auxiliary variables  $a_k$  for  $k \in \{1, \dots, r\}$  added by the reduction. Now, for each  $t \in \{1, \dots, T-1\}$  and  $j_1, \dots, j_7 \in \{1, \dots, T\}$ , we define a 3SAT formula  $\varphi_{\text{Check}, t, j_1, \dots, j_7}$  analogously to  $\mathcal{C}_{\text{Check}, t, j_1, \dots, j_7}$ . We begin by associating those variables in  $\varphi_{\text{Check}}$  which come from  $\mathcal{C}_{\text{Check}}$ 's inputs with the variables in  $\mathcal{V}$  as in Definition 10.11. Next, for each  $k \in \{1, \dots, r\}$ , we introduce a new variable  $a_{t, j_1, \dots, j_7, k} \in \{0, 1\}$  and associate it with the variable  $a_k$ . This defines  $\varphi_{\text{Check}, t, j_1, \dots, j_7}$ . In summary, the final 3SAT instance produced by the Cook-Levin reduction is

$$\varphi := \varphi_{\text{Boundary}} \wedge \left( \bigwedge_{t, j_1, \dots, j_7} \varphi_{\text{Check}, t, j_1, \dots, j_7} \right). \quad (122)$$

By Proposition 10.14, the execution tableau properly encodes the execution of  $\mathcal{D}$  if and only if there exists a setting to the auxiliary variables satisfying the 3SAT formula  $\varphi$ . In total,  $\varphi$  contains

$$|\mathcal{V}| + O(T^8) \cdot r \leq O(T^2 + T \cdot \log(|\mathcal{D}|) + T^8 \cdot \text{poly}(|\mathcal{D}|)) = \text{poly}(T, |\mathcal{D}|) \quad (123)$$

variables. The second term on the left-hand side of Equation (123) corresponds to the auxiliary variables in each of the  $O(T^8)$  copies of  $\varphi_{\text{Check}}$ . The inequality follows by Equation (112) and the fact that  $r \leq \text{poly}(|\mathcal{D}|)$ .

Our next task is to represent the 3SAT formula  $\varphi$  succinctly. To do this, we will provide a circuit  $\mathcal{C}$  which succinctly describes a 3SAT formula  $\varphi_{3\text{SAT}}$  which, while not literally equal to  $\varphi$ , will be isomorphic to it. This means that, for example,  $\varphi_{3\text{SAT}}$  may not even have the same number of variables as  $\varphi$ , but any variable in  $\varphi$  will correspond in a clear and direct manner to a variable in  $\varphi_{3\text{SAT}}$ , and any remaining variables in  $\varphi_{3\text{SAT}}$  do not appear in any clauses. This circuit is constructed as follows.

**Definition 10.15.** In this definition we construct the circuit  $\mathcal{C}$ . It has three inputs  $z_1, z_2, z_3$  of length  $m$ , which we specify below in Equation (124), and three inputs  $o_1, o_2, o_3 \in \{0, 1\}$ . For each  $v \in \{1, 2, 3\}$ , each  $z_v$  is supposed to specify a variable in  $\varphi$  according to a format we will now specify. If  $z_v$  is not properly formatted, then it does not correspond to a variable in  $\varphi$ ; if any of  $z_1, z_2, z_3$  is not properly formatted, then  $\mathcal{C}$  automatically outputs 0. Below, we will often write substrings of the  $z_v$ 's as though they are integers from some specified range, i.e.  $a \in \{b, \dots, c\}$ . This means that  $a$  is represented as a binary string of length  $\lceil \log(c+1) \rceil$ , which is to be interpreted as the binary encoding of an integer between  $b$  and  $c$ .

The input  $z_v$  is formatted as a string  $(\omega, \alpha, \beta_1, \beta_2, \beta_3, \beta_4)$ . The first substring  $\omega$  has length  $m - (|\alpha| + |\beta_1| + \dots + |\beta_4|)$  bits and is formatted to be the all-zeroes string. Its purpose is to pad the inputs to have the length  $m$  we specify below. Next,  $\alpha$  is formatted as an integer  $\alpha \in \{1, 2, 3, 4\}$ . The variable encoded by  $z_v$  is specified by  $\beta_\alpha$ , and the other three  $\beta$ 's should be the all-zeros string. We now specify the encoding of  $\beta_\alpha$ , conditioned on the value of  $\alpha$ .

1.  $\beta_1$  is formatted as  $(t, i, j, k)$ , where

$$t \in \{1, \dots, T\}, \quad i \in \{1, \dots, 7\}, \quad j \in \{0, 1, \dots, T+1\}, \quad k \in \{1, 2\}.$$

This corresponds to the variable  $c_{t,i,j,k}$ .

2.  $\beta_2$  is formatted as  $(t, i, j)$ , where

$$t \in \{1, \dots, T\}, \quad i \in \{1, \dots, 7\}, \quad j \in \{0, 1, \dots, T+1\}.$$

This corresponds to the variable  $h_{t,i,j}$ .

3.  $\beta_3$  is formatted as  $(t, k)$ , where

$$t \in \{1, \dots, T\}, \quad k \in \{1, \dots, \kappa\}.$$

This corresponds to the variable  $s_{t,k}$ .

4.  $\beta_4$  is formatted as  $(t, j_1, \dots, j_7, k)$ , where

$$t \in \{1, \dots, T-1\}, \quad j_1, \dots, j_7 \in \{1, 2, \dots, T\}, \quad k \in \{1, \dots, r\}.$$

This corresponds to the variable  $a_{t,j_1, \dots, j_7, k}$ .

In total, the length of these substrings is

$$|\alpha| + |\beta_1| + \dots + |\beta_4| = O(\log(T) + \log(\kappa) + \log(r)) = O(\log(T) + \log(|\mathcal{D}|)).$$

As a result, because  $|\mathcal{D}| \leq \sigma$ , the length of the “padding”  $\omega$  can be chosen so that each  $z_i$  has length

$$m = O(\log(T) + \log(\sigma)). \quad (124)$$

Checking that each  $z_\nu$  is properly formatted can be done by checking that certain substrings of  $z_1, z_2$ , and  $z_3$  encode integers which fall within specified ranges. This can be done using  $O(m)$  gates.

Having specified the inputs, we can now specify the execution of the circuit, and we may assume that  $z_1, z_2, z_3$  are properly formatted. Implementing the clauses from  $\varphi_{\text{Boundary}}$  is simple; we specify how to implement the clause from Equation (113).

- Suppose for input  $z_1$ ,  $\alpha = 2$ . Then  $\beta_2$  can be parsed as  $(t, i, j)$ . The circuit accepts if  $t = j = 1$  and  $o_1 = 1$ , regardless of  $z_2$  and  $z_3$ . This ensures that  $\varphi_{3\text{SAT}}$  includes  $x_{z_1} \vee x_{z_2}^{o_2} \vee x_{z_3}^{o_3}$  for any  $z_2, z_3, o_2, o_3$ , which is equivalent to including the arity-one clause  $x_{z_1}$ .

This can be implemented with  $O(m)$  gates. Similar arguments can be used to implement the clauses from Equations (114)-(119) using  $O(m)$  gates apiece.

Implementing the clauses from the formulas  $\varphi_{\text{Check}, t, j_1, \dots, j_7}$  is more challenging. From Equation (121), we can see that any constraint in this formula always involves a variable of the form  $a_{t, j_1, \dots, j_7, k}$  for some  $k \in \{1, \dots, r\}$ .

1. First check if one of its inputs  $z_1, z_2, z_3$  corresponds to such a variable. This can be done with  $O(1)$  gates simply by checking if for any of the  $z_i$ 's,  $a = 4$ . If so, this specifies the values of  $t, j_1, \dots, j_7$ .

2. Each variable in  $\varphi_{\text{Check},t,j_1,\dots,j_7}$  is associated with a variable in  $\varphi_{\text{Check}}$ . The circuit checks if all the  $z_i$ 's are contained in  $\varphi_{\text{Check},t,j_1,\dots,j_7}$  and then it computes which variable in  $\varphi_{\text{Check}}$  they are associated with. We include below the example of checking whether  $z_1$  is associated with the variable  $c_{\text{Check},1,1,0,0}$ .
  - The circuit  $\mathcal{C}$  first computes  $t + 1$ , which takes  $O(m)$  gates. It then looks at  $z_1$  and checks if  $\alpha = 1$ . If so, then  $\beta_1 = (t', i', j', k')$ , and so it tests the equalities  $t' = t + 1$ ,  $i' = 1$ ,  $j' = j_1$ , and  $k' = 0$ , each of which takes  $O(m)$  gates to test. If so, then  $z_1$  is associated with  $c_{\text{Check},1,1,0,0}$ .

There are  $\text{poly}(|\mathcal{D}|)$  variables in  $\varphi_{\text{Check}}$  and, for each variable  $z_i$ , it takes  $O(m)$  gates to determine whether  $z_i$  is associated with this variable. As a result, because  $|\mathcal{D}| \leq \sigma$ , this takes  $\text{poly}(\sigma, \log(T))$  gates to compute.

3. Whether  $z_1, z_2$ , and  $z_3$  share a clause in  $\varphi_{\text{Check},t,j_1,\dots,j_7}$  depends only on which variables in  $\varphi_{\text{Check}}$  they are associated with. As a result, after having computed these variables, the algorithm can hard-code whether the circuit  $\mathcal{C}$  should accept.

This completes the description of  $\mathcal{C}$ . In total, it contains  $\text{poly}(\sigma, \log(T))$  gates. Computing  $\mathcal{C}$  first requires computing  $\varphi_{\text{Check}}$ , which takes time  $\text{poly}(|\mathcal{D}|) \leq \text{poly}(\sigma)$  by Propositions 10.12 and 10.14. After that, the steps outlined above for construction  $\mathcal{C}$  take time  $\text{poly}(\sigma, \log(T))$ .

The circuit  $\mathcal{C}$  succinctly describes  $\varphi$  in the loose sense described above. Recall that  $\varphi$  accepts if and only if it encodes the execution of  $\mathcal{D}$  up to time  $T$ . We now modify  $\mathcal{C}$  to (i) hard code  $n, x$ , and  $y$  onto  $\mathcal{D}$ 's input tapes, and (ii) ensure that  $\mathcal{D}$  accepts. We demonstrate how to do so by hard coding  $n$  as an example.

- The input  $n$  is described by some string  $v$  of length  $\ell = O(\log(n))$ . We would like to hard-code the string  $(v, \sqcup^{T-\ell})$  into the first tape of  $\mathcal{D}$ . To do this, we first check if  $z_1$  corresponds to the variable  $c_{t,i,j,k}$  for some values of  $t, i, j, k$ . Then, we check if  $t = 1$ , the time where the inputs appear on the tapes, and  $i = 1$ , corresponding to the first tape. If so, we branch on whether  $j \leq \ell$ . If it is, then if  $k = 1$  the circuit accepts if  $o_1 = 0$ , and if  $k = 2$  the circuit accepts if  $o_1 = v_j$ . Otherwise, if  $j > \ell$ , then if  $k = 1$  the circuit accepts if  $o_1 = 1$ , and if  $k = 2$  the circuit accepts if  $o_1 = 0$ . This can be done with  $\text{poly}(\log(n), \log(T))$  gates.

This modifies  $\varphi$  so that it only accepts if  $n$  is written on its first tape at input. We can similarly hard-code  $x$  and  $y$  onto the second and third input tapes and hard-code the accepting state as the final state of  $\mathcal{D}$ . In total, this takes  $\text{poly}(\log(n), |x|, |y|, \log(T), \sigma)$ , which is  $\text{poly}(\log(n), Q, \log(T), \sigma)$  because  $x$  and  $y$  have length at most  $Q$ .

It remains to ensure that the variables corresponding to the 4th and 5th at time  $t = 1$  are the lexicographically-first named variables in  $\varphi$ . However, this is simple and can be done using  $\text{poly}(\log(n), Q, \log(T), \sigma)$  gates. This concludes the construction.  $\square$

### 10.3 A succinct 5SAT description for deciders

Proposition 10.9 allows us to convert any decider  $\mathcal{D}$  and inputs  $n, x, y$  into a 3SAT formula  $\varphi_{3\text{SAT}}$ , succinctly described by a circuit  $\mathcal{C}$ , which represents it. However, there are two undesirable properties of this construction, which we describe below.

1. First, evaluating any clause  $(w_{i_1}^{o_{i_1}} \vee w_{i_2}^{o_{i_2}} \vee w_{i_3}^{o_{i_3}})$  of  $\varphi_{3\text{SAT}}$  requires evaluating the *same* assignment  $w$  at three separate points. While this is fine when the assignment  $w$  is provided in full to the verifier, it can be a problem when the verifier is only able to query the points in  $w$  by interacting with a

prover. In this case, the verifier might send the prover the values  $i_1, i_2, i_3$ , who responds with three bits  $b_1, b_2, b_3 \in \{0, 1\}$ , purported to be the values  $w_{i_1}, w_{i_2}, w_{i_3}$  for some assignment  $w \in \{0, 1\}^M$ . As it will turn out, in the answer reduced protocol below, the verifier will actually be able to force the prover to reply using *three* different assignments. In other words, the prover will have three assignments  $w_1, w_2, w_3 \in \{0, 1\}^M$  such that, for any  $i_1, i_2, i_3$  provided to it by the verifier, it will respond with  $b_1 = w_{1,i_1}, b_2 = w_{2,i_2}, b_3 = w_{3,i_3}$ . However, even given this there is no guarantee that the three assignments are the *same* assignments (i.e. that  $w_1 = w_2 = w_3$ ). In the work of [NW19], this was accomplished by an additional subroutine called the *intersecting lines test*, which would enforce consistency between  $w_1, w_2$ , and  $w_3$ . In this work, on the hand, we would like to relax the assumption that  $w_1, w_2$ , and  $w_3$  must be the same. This will allow us to not use the intersecting lines test, simplifying the answer reduction protocol. (In fact, the answer reduced verifier will not be querying the assignments directly, but rather low-degree encodings of these assignments.)

2. Second, we are guaranteed that  $w = (a, b, c)$  satisfies  $\varphi_{3\text{SAT}}$  if and only if  $\mathcal{D}$  accepts  $(n, x, y, a, b)$ . However, it is inconvenient that  $a$  and  $b$  are contained as substrings of  $w$ . To see why, recall from Section 9 that the oracularized verifier sometimes gives one prover a pair of questions  $(x, y)$  and another prover just one of the questions—say,  $x$ . The answer reduced verifier will sample its questions similarly; as for its answers, it might expect the first prover to respond with a string  $w = (a, b, c)$  that satisfies  $\varphi_{3\text{SAT}}$  and the second prover to respond with a string  $a'$  such that  $a = a'$ . Verifying that  $a = a'$  requires the verifier to sample a uniformly random point from  $w$ , restricted to the coordinates in  $a$ . As it turns out, generating a uniform point from a substring is extremely cumbersome, though not impossible, to do when the verifier's questions are expected to be sampled from conditional linear functions. To remove this complication, though, we would instead prefer if the provers' answers were formatted in a way that gave the verifier direct access to the strings  $a$  and  $b$ .

In the remainder of this section, we will show how to modify the Succinct-3SAT circuits produced by Proposition 10.9 in order to ameliorate these two difficulties. Doing so entails modifying the  $\varphi_{3\text{SAT}}$  formula from Proposition 10.9 to produce a 5SAT formula  $\varphi_{5\text{SAT}}$ . The clauses of  $\varphi_{5\text{SAT}}$  will be of the form

$$a_{i_1}^{o_1} \vee b_{i_2}^{o_2} \vee w_{1,i_3}^{o_3} \vee w_{2,i_4}^{o_4} \vee w_{3,i_5}^{o_5},$$

where  $a, b, w_1, w_2$ , and  $w_3$  are five separate assignments which are not assumed to be equal. The guarantee is that  $(a, b, w_1, w_2, w_3)$  satisfies  $\varphi_{5\text{SAT}}$  if and only if  $\mathcal{D}$  accepts  $(n, x, y, a, b)$ . This addresses the two concerns from above: each clause is totally *decoupled*, meaning it samples 5 variables from 5 different assignments, and so no consistency check must be performed between the assignments. In addition, the first two strings exactly correspond to  $a$  and  $b$ , addressing the second item.

We now formally define decoupled 5SAT instances and how they succinctly represent bounded deciders. Following that, we show how the succinct 3SAT instance  $\varphi_{3\text{SAT}}$  produced by Proposition 10.9 can be modified to produce a succinct 5SAT instance  $\varphi_{5\text{SAT}}$  which represents  $\mathcal{D}$ .

**Definition 10.16** (Decoupled 5SAT and its succinct descriptions). A *block* of variables  $x_i$  is a tuple  $x_i = (x_{i,0}, \dots, x_{i,N_i-1})$ . A formula  $\varphi$  on 5 blocks  $x_1, x_2, \dots, x_5$  of variables is called a *decoupled 5SAT formula* if every clause is of the form

$$x_{1,i_1}^{o_1} \vee x_{2,i_2}^{o_2} \vee x_{3,i_3}^{o_3} \vee x_{4,i_4}^{o_4} \vee x_{5,i_5}^{o_5}, \quad (125)$$

for  $i_j \in \{0, 1, \dots, N_j - 1\}$  and  $o_1, \dots, o_5 \in \{0, 1\}$ . (Recall from Definition 10.2 that the notation  $x^o$  means  $x$  if  $o = 1$  and  $\neg x$  if  $o = 0$ .)

For each  $i \in \{1, 2, \dots, 5\}$ , suppose each  $N_i$  is a power of two, and write it as  $N_i = 2^{n_i}$ . Let  $\mathcal{C}$  be a circuit with five inputs of length  $n_1, n_2, \dots, n_5$  and five single-bit inputs. Then  $\mathcal{C}$  succinctly describes decoupled  $\varphi$  if, for all  $i_j \in \{0, 1, \dots, N_j - 1\}$  and  $o_1, o_2, \dots, o_5 \in \{0, 1\}$ ,

$$\mathcal{C}(i_1, i_2, \dots, i_5, o_1, o_2, \dots, o_5) = 1 \quad (126)$$

if and only if the clause in (125) is included in  $\varphi$ . As in Definition 10.2, we slightly abuse notation and use the convention that a number  $a$  between 0 and  $2^{n_i} - 1$  is interpreted as its binary encoding  $\text{binary}_{n_i}(a)$  when provided as input to a set of  $n_i$  single-bit wires.

**Definition 10.17** (Succinct descriptions for bounded deciders). Let  $\mathcal{D}$  be a decider. Fix an index  $n \in \mathbb{N}$  and a time  $T \in \mathbb{N}$ . Let  $L = 2^\ell$  be a power of two that is at least as large as  $2T$ . Let  $x$  and  $y$  be strings,  $r \in \mathbb{N}$  and  $R = 2^r$ .

Consider a circuit  $\mathcal{C}$  with two inputs of length  $\ell$ , three inputs of length  $r$ , and 5 single-bit inputs. Let  $\varphi_{\mathcal{C}}$  be the decoupled 5SAT instance with two blocks of variables of size  $L$  and three blocks of size  $R$  which  $\mathcal{C}$  succinctly describes. Then we say that  $\mathcal{C}$  succinctly describes  $\mathcal{D}$  (on inputs  $n$ ,  $x$ , and  $y$  and time  $T$ ) if, for all  $a, b \in \{0, 1\}^L$ , there exists  $w_1, w_2, w_3 \in \{0, 1\}^R$  such that  $a, b, w_1, w_2, w_3$  satisfy  $\varphi_{\mathcal{C}}$  if and only if there exist  $a_{\text{prefix}}, b_{\text{prefix}} \in \{0, 1\}^*$  of lengths  $\ell_a, \ell_b \leq T$ , respectively, such that

$$a = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{L/2-\ell_a}) \quad \text{and} \quad b = \text{enc}_{\Gamma}(b_{\text{prefix}}, \sqcup^{L/2-\ell_b})$$

and  $\mathcal{D}$  accepts  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  in time  $T$ .

In this definition of succinct descriptions, the answers  $a$  and  $b$  are isolated, in that the first input of  $\mathcal{C}$  of length  $\ell$  indexes into  $a$  and the second input of length  $\ell$  indexes into  $b$ . The next proposition shows how to construct such descriptions.

**Proposition 10.18** (Explicit succinct descriptions). *There is a Turing machine SuccinctDecider with the following properties. Let  $\mathcal{D}$  be a decider, let  $n, T, Q$ , and  $\sigma$  be integers with  $Q \leq T$  and  $|\mathcal{D}| \leq \sigma$ , and let  $x$  and  $y$  be strings of length at most  $Q$ . Then on input  $(\mathcal{D}, n, T, Q, \sigma, x, y)$ , SuccinctDecider outputs a circuit  $\mathcal{C}$  with two inputs of length  $\ell_0(T)$ , three of length  $r_0(T, \sigma)$ , and five single-bit inputs which succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ . Moreover, the following hold.*

1.  $\ell_0(T) = \lceil \log(2T) \rceil$ .
2.  $r_0(T, \sigma) = O(\log(T) + \log(\sigma))$ ,
3.  $\mathcal{C}$  has at most  $s_0(n, T, Q, \sigma) = \text{poly}(\log(T), \log(n), Q, \sigma)$  gates,
4. SuccinctDecider runs in time  $\text{poly}(\log(T), \log(n), Q, \sigma)$ , and the parameters  $\ell_0, r_0, s_0$  can be computed from  $n, T, Q, \sigma$  in time  $\text{poly}(\log(T), \log(n), \log(Q), \sigma)$ .

*Proof.* The Turing machine SuccinctDecider begins by running the algorithm in Proposition 10.9 on input  $(\mathcal{D}, n, T, Q, \sigma, x, y)$  to produce a circuit  $\mathcal{C}_{3\text{SAT}}$  on  $3r_0 + 3$  inputs, where  $r_0 = O(\log(T) + \log(\sigma))$  is the parameter  $m$  from the proposition. Set  $\ell_0 = \lceil \log(2T) \rceil$ ,  $L = 2^{\ell_0}$  and  $R = 2^{r_0}$ . Given this, SuccinctDecider returns the circuit  $\mathcal{C}$  with inputs  $i_1, i_2 \in \{0, 1, \dots, L - 1\}$ ,  $i_3, i_4, i_5 \in \{0, 1, \dots, R - 1\}$ ,  $o_1, o_2, \dots, o_5 \in \{0, 1\}$ , and

$$\mathcal{C}(i_1, i_2, \dots, i_5, o_1, o_2, \dots, o_5) = 1,$$

if one of the following conditions hold.

$$\begin{aligned}
& \mathcal{C}_{3\text{SAT}}(i_3, i_4, i_5, o_3, o_4, o_5) = 1, \\
& (i_1 < 2T) \wedge (i_1 = i_3) \wedge (o_1 \neq o_3), \\
& (i_2 < 2T) \wedge (i_2 = i_3 - 2T) \wedge (o_2 \neq o_3), \\
& (i_1 \geq 2T) \wedge (i_1 \text{ is odd}) \wedge (o_1 = 1), \\
& (i_1 \geq 2T) \wedge (i_1 \text{ is even}) \wedge (o_1 = 0), \\
& (i_2 \geq 2T) \wedge (i_2 \text{ is odd}) \wedge (o_2 = 1), \\
& (i_2 \geq 2T) \wedge (i_2 \text{ is even}) \wedge (o_2 = 0), \\
& (i_3 = i_4) \wedge (o_3 \neq o_4), \\
& (i_4 = i_5) \wedge (o_4 \neq o_5).
\end{aligned}$$

It is not hard to verify that testing “ $(i_1 < 2T)$ ” can be done with  $O(\ell_0)$  AND and OR gates, and testing  $(i_2 = i_3 - 2T)$  can be done with  $O(m)$  AND and OR gates. Using similar estimates for the remaining sub-circuits, we compute

$$\begin{aligned}
\text{size}(\mathcal{C}) &= \text{size}(\mathcal{C}_{3\text{SAT}}) + O(r_0 + \ell_0) = \text{size}(\mathcal{C}_{3\text{SAT}}) + O(r_0) \\
&\leq \text{poly}(\log(T), \log(n), Q, \sigma) + O(\log(T) + \log(\sigma)).
\end{aligned}$$

In addition, due to the simplicity of these modifications, we conclude that the runtime of SuccinctDecider is dominated by the runtime of the algorithm from Proposition 10.9, which is  $\text{poly}(\log(T), \log(n), Q, \sigma)$ .

Now we show that  $\mathcal{C}$  succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ . To begin, we describe the decoupled 5SAT formula  $\varphi_{\mathcal{C}}$ . Let us first consider the constraints in  $\varphi_{\mathcal{C}}$  which are implied by the final constraint, i.e. those of the form

$$a_{i_1}^{o_1} \vee b_{i_2}^{o_2} \vee (w_1)_{i_3}^{o_3} \vee (w_2)_{i_4}^{o_4} \vee (w_3)_{i_5}^{o_5}$$

whenever  $i_4 = i_5$  and  $o_4 \neq o_5$ . For any fixed  $i_1, i_2, i_3$ , the negations  $o_1, o_2, o_3$  can take any values, and as a result, the first three bits in the constraint vary over all assignments in  $\{0, 1\}^3$ . This means that these constraints are satisfied if and only if  $(w_2)_{i_4}^{o_4} \vee (w_3)_{i_5}^{o_5}$  is satisfied whenever  $i_4 = i_5$  and  $o_4 \neq o_5$ . This, in turn, is equivalent to the constraint  $w_2 = w_3$ . Carrying out similar arguments for the entire circuit, we can express the formula  $\varphi_{\mathcal{C}}$  as follows.

$$\begin{aligned}
\varphi_{\mathcal{C}}(a, b, w_1, w_2, w_3) &= \varphi_{3\text{SAT}}(w_1, w_2, w_3) \wedge (w_{1,1} = a_1) \wedge (w_{1,2} = b_1) \\
&\wedge (a_2 = (10)^{L/2-T}) \wedge (b_2 = (10)^{L/2-T}) \wedge (w_1 = w_2) \wedge (w_2 = w_3).
\end{aligned}$$

Here, we write  $\varphi_{3\text{SAT}}(w_1, w_2, w_3)$  for the formula in which, for each constraint in  $\varphi_{3\text{SAT}}$ , the first variable is taken from  $w_1$ , the second from  $w_2$ , and the third from  $w_3$ . In addition, we write  $a = (a_1, a_2)$ , where  $a_1$  is the first  $2T$  bits in  $a$  and  $a_2$  is the remaining  $L - 2T$  bits, and similarly for  $b = (b_1, b_2)$ . We also write  $w_1 = (w_{1,1}, w_{1,2}, w_{1,3})$ , where  $w_{1,1}$  contains the first  $2T$  bits in  $w_1$ ,  $w_{1,2}$  contains the second  $2T$  bits, and  $w_{1,3}$  contains the remaining  $R - 4T$  bits. As a result,  $\varphi_{\mathcal{C}}$  is satisfied only if  $w_1 = w_2 = w_3 = (a_1, b_1, c)$  for some string  $c \in \{0, 1\}^{R-4T}$ . In this case, calling  $w = (a_1, b_1, c)$ ,  $\varphi_{\mathcal{C}}$  is satisfied only if  $\varphi_{3\text{SAT}}(w)$  is. By Proposition 10.9, this implies that there exists a string  $a_{\text{prefix}}$  of length  $\ell_a \leq T$  such that  $a_1 = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{T-\ell_a})$ . This, in turn, implies that

$$a = (a_1, a_2) = (\text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{T-\ell_a}), (10)^{L-2T}) = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{L/2-\ell_a}),$$

using the fact that  $\text{enc}_\Gamma(\sqcup) = 10$ , and similarly for  $b$ . Finally, Proposition 10.9 implies that  $\mathcal{D}$  accepts  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  in time  $T$ . This completes the proof.  $\square$

As stated above, moving from 3SAT to 5SAT allows us to devote the first two inputs to  $a$  and  $b$ . In addition, we have added extra constraints into  $\varphi_{\mathcal{C}}$  which enforce that  $w_1 = w_2 = w_3$ , which means that we can relax this assumption on these assignments.

We now show a simple transformation that takes in a succinct circuit  $\mathcal{C}$  and outputs another succinct circuit  $\mathcal{C}'$  whose 5 inputs are “padded” to contain more input bits.

**Proposition 10.19** (Padding). *Let  $\mathcal{C}$  be a circuit of size  $s$  with two inputs of length  $\ell$ , three inputs of length  $r$ , and 5 single-bit inputs. Suppose  $\mathcal{C}$  succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ . Then there is an algorithm which takes as input  $(\mathcal{C}, \ell, r, \ell', r')$ , with  $\ell' \geq \ell$  and  $r' \geq r$ , and in time  $\text{poly}(s, \ell', r')$  outputs a circuit  $\mathcal{C}'$  with the following properties. First,  $\mathcal{C}'$  has two inputs of length  $\ell'$ , three inputs of length  $r'$ , and 5 single-bit inputs, and its size is  $s + \text{poly}(\ell', r')$ . Second, it succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ .*

*Proof.* Write  $L = 2^\ell, R = 2^r$  and  $L' = 2^{\ell'}, R' = 2^{r'}$ . The algorithm constructs the circuit  $\mathcal{C}'$  which on inputs  $i_1, i_2 \in \{0, \dots, L' - 1\}$ ,  $i_3, i_4, i_5 \in \{0, \dots, R' - 1\}$ , and  $o_1, \dots, o_5 \in \{0, 1\}$ , outputs 1 if and only if one of the following conditions hold:

$$\begin{aligned} & (i_1, i_2 < L) \wedge (i_3, i_4, i_5 < R) \wedge \mathcal{C}(i_1, i_2, i_3, i_4, i_5, o_1, o_2, o_3, o_4, o_5) = 1, \\ & (i_1 \geq L) \wedge (i_1 \text{ is odd}) \wedge (o_1 = 1), \\ & (i_1 \geq L) \wedge (i_1 \text{ is even}) \wedge (o_1 = 0), \\ & (i_2 \geq L) \wedge (i_2 \text{ is odd}) \wedge (o_2 = 1), \\ & (i_2 \geq L) \wedge (i_2 \text{ is even}) \wedge (o_2 = 0). \end{aligned}$$

Now we show that  $\mathcal{C}'$  succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ . To begin, we describe the decoupled 5SAT formula  $\varphi_{\mathcal{C}'}$ . The second and third constraints imply that for each  $i_1 \geq L$ ,  $\varphi_{\mathcal{C}'}$  contains the constraint  $(a_{i_1})$  if  $i_1$  is odd and  $(\neg a_{i_1})$  if  $i_1$  is even. Likewise, the fourth and fifth constraints imply that for each  $i_2 \geq L$ ,  $\varphi_{\mathcal{C}'}$  contains the constraint  $(b_{i_2})$  if  $i_2$  is odd and  $(\neg a_{i_2})$  if  $i_2$  is even. Thus, we can express the formula  $\varphi_{\mathcal{C}'}$  as follows.

$$\varphi_{\mathcal{C}'}(a, b, w_1, w_2, w_3) = \varphi_{\mathcal{C}}(a_1, b_1, w_{1,1}, w_{2,1}, w_{3,1}) \wedge (a_2 = (10)^{(L'-L)/2}) \wedge (b_2 = (10)^{(L'-L)/2}). \quad (127)$$

Here, we write  $a = (a_1, a_2)$  and  $b = (b_1, b_2)$ , where  $a_1, b_1$  have length  $L$  and  $a_2, b_2$  have length  $L' - L$ , and for each  $i \in \{1, 2, 3\}$ , we write  $w_i = (w_{i,1}, w_{i,2})$ , where  $w_{i,1}$  has length  $R$  and  $w_{i,2}$  has length  $R' - R$ .

Now, suppose there exist  $w_1, w_2, w_3$  such that  $a, b, w_1, w_2, w_3$  satisfy  $\varphi_{\mathcal{C}'}$ . Then because  $\mathcal{C}$  succinctly represents  $\mathcal{D}$ , there exists  $a_{\text{prefix}}, b_{\text{prefix}}$  of lengths  $\ell_a, \ell_b \leq T$  such that  $\mathcal{D}$  accepts  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$ . In addition,

$$a_1 = \text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{L/2-\ell_a}),$$

and likewise for  $b_1$ . Equation (127) then implies that

$$\begin{aligned} a &= (a_1, a_2) = (\text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{L/2-\ell_a}), (10)^{(L'-L)/2}) \\ &= (\text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{L/2-\ell_a}), \text{enc}_\Gamma(\sqcup)^{(L'-L)/2}) \\ &= \text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{L'/2-\ell_a}), \end{aligned}$$

where the third step used the fact that  $\text{enc}_\Gamma(\sqcup) = 10$ . As a similar statement holds for  $b$ , this establishes that  $\mathcal{C}'$  succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ .  $\square$



By combining Proposition 10.18 and Proposition 10.19 (choosing  $\ell' = r' = r_0$  where  $r_0$  is from Proposition 10.18), we get the following:

**Proposition 10.20** (Explicit padded succinct descriptions). *There is a Turing machine PaddedSuccinctDecider with the following properties. Let  $\mathcal{D}$  be a decider; let  $n, T, Q$ , and  $\sigma$  be integers with  $Q \leq T$  and  $|\mathcal{D}| \leq \sigma$ , and let  $x$  and  $y$  be strings of length at most  $Q$ . Then on input  $(\mathcal{D}, n, T, Q, \sigma, x, y)$ , PaddedSuccinctDecider outputs a circuit  $\mathcal{C}$  with five inputs of length  $m(T, \sigma)$  and five single-bit inputs which succinctly describes  $\mathcal{D}$  on inputs  $n, x$ , and  $y$  and time  $T$ . Moreover, the following hold.*

1.  $m(T, \sigma) = O(\log(T) + \log(\sigma))$  and  $2^m \geq 2T$ .
2.  $\mathcal{C}$  has at most  $s(n, T, Q, \sigma)$  gates, where  $s$  is such that  $s(n, T, Q, \sigma) = \text{poly}(\log(T), \log(n), Q, \sigma)$  and  $5m(T, \sigma) + 5 + s(n, T, Q, \sigma)$  is a power of 2.
3. PaddedSuccinctDecider runs in time  $\text{poly}(\log(T), \log(n), Q, \sigma)$ , and the parameters  $m(T, \sigma), s(n, T, Q, \sigma)$  can be computed from  $n, T, Q, \sigma$  in time  $\text{poly}(\log(T), \log(n), \log(Q), \sigma)$ .

## 10.4 A PCP for normal form deciders

We give a probabilistically checkable proof (PCP) for the Bounded Halting problem specialized to the case of normal form deciders. Our PCP will use standard techniques from the algebraic, low-degree-code-based PCP literature. In particular, we slightly modify the PCP for Succinct-3SAT described in [NW19, Section 11] (which itself is based on the proof of the PCP theorem in [Har04]) to apply it to the decoupled Succinct-5SAT instances described in Section 10.3. We follow their treatment closely. As the PCPs constructed in this section are only an intermediate object towards the normal form verifier introduced in the next section we do not include standard definitions on PCPs, and refer to these references (in particular [Har04]) for background. We begin with some preliminaries.

### 10.4.1 Preliminaries

A key part of the PCP will be to design a function  $f : \mathbb{F}^n \rightarrow \mathbb{F}$  which is zero on the subcube  $H_{\text{subcube}} = \{0, 1\}^n$ . Our next proposition shows that given such a function  $f$ , there is a way of writing it so that the fact that it is zero on  $H_{\text{subcube}}$  is self-evidently true. Doing so involves showing that  $f$  can be written in a simple basis of polynomials which are constructed to be zero on the subcube. This fact is standard in the literature (see, for example, [BSS08, Lemma 4.11]), and we include its proof for completeness.

**Proposition 10.21** (Polynomial basis of zero functions). *Let  $\mathbb{F}$  be a field. Define  $\text{zero} : \mathbb{F} \rightarrow \mathbb{F}$  as the univariate polynomial  $x \mapsto x(1 - x)$ . Define  $H_{\text{subcube}} = \{0, 1\}^n$ . Suppose  $f : \mathbb{F}^n \rightarrow \mathbb{F}$  is an individual degree  $d$  polynomial such that  $f(x) = 0$  for all  $x \in H_{\text{subcube}}$ . Then there exist polynomials  $c_1, \dots, c_n : \mathbb{F}^n \rightarrow \mathbb{F}$  such that for all  $x \in \mathbb{F}^n$ ,*

$$f(x) = \sum_{i=1}^n c_i(x) \cdot \text{zero}(x_i).$$

In addition, for each  $i \in \{1, \dots, n\}$ ,  $c_i$  has individual degree- $d$ .

*Proof.* To prove this, we first prove the following statement for each  $k \in \{0, 1, \dots, n\}$ : there exists an individual degree- $d$  polynomial  $r_k : \mathbb{F}^n \rightarrow \mathbb{F}$  and polynomials  $c_1, \dots, c_k : \mathbb{F}^n \rightarrow \mathbb{F}$  such that

$$f(x) = \sum_{i=1}^k c_i(x) \cdot \text{zero}(x_i) + r_k(x). \tag{128}$$

In addition, for each  $i \in \{1, \dots, k\}$ ,  $c_i$  has individual degree  $d$  and the degree of  $x_i$  in  $r_k$  is at most 1.

The proof is by induction on  $k$ , the base case of  $k = 0$  being trivial. Now, we perform the induction step. Assuming that Equation (128) holds for  $k$ , we will show that it holds for  $k + 1$  as well. Let  $r_k$  be the polynomial guaranteed by the inductive hypothesis. We now divide  $r_k$  by  $\text{zero}(x_{k+1})$  using polynomial division. This guarantees a polynomial  $c_{k+1}$  and an individual degree- $d$  polynomial  $r_{k+1}(x)$  such that

$$r_k(x) = c_{k+1}(x) \cdot \text{zero}(x_{k+1}) + r_{k+1}(x).$$

In addition,  $c_{k+1}$  still has individual degree  $d$  (and in fact the degree of  $x_{k+1}$  in  $c_{k+1}$  is at most  $d - 2$ ). Furthermore, for each  $i \in \{1, \dots, k + 1\}$ ,  $r_{k+1}$  has degree at most 1 in  $x_i$ . Plugging this into Equation (128), we see that

$$f(x) = \sum_{i=1}^{k+1} c_i(x) \cdot \text{zero}(x_i) + r_{k+1}(x). \quad (129)$$

This completes the induction.

Applying the  $k = n$  case of Equation (128), we see that

$$f(x) = \sum_{i=1}^n c_i(x) \cdot \text{zero}(x_i) + r(x), \quad (130)$$

where the degree of each variables in  $r$  is at most 1. Since  $f(x)$  vanishes on the subcube  $H_{\text{subcube}}$ , it must also be that  $r(x)$  vanishes on  $H_{\text{subcube}}$  as well. We claim that  $r$  must therefore be the zero polynomial.

We prove this by showing the following statement: for every polynomial  $s(x_1, \dots, x_n)$  with individual degree 1 (i.e. a multilinear polynomial) that vanishes on all points  $x \in \{0, 1\}^n$ ,  $s$  must be the zero polynomial. We show this by induction on the number of variables.

Consider the base case  $n = 1$ . Then  $s$  is a univariate polynomial with degree at most 1. However it is zero on two points, so therefore it must be the zero polynomial. Now for the inductive step: assuming the proposition holds for some  $n \geq 1$ , we show that it holds for  $n + 1$  as well. For any multilinear polynomials, we can write

$$s(x_1, \dots, x_{n+1}) = x_{n+1}s_1(x_1, \dots, x_n) + s_2(x_1, \dots, x_n)$$

where both  $s_1$  and  $s_2$  are  $n$ -variate multilinear polynomials. Fix  $x_{n+1} = 0$ . Then  $s(x_1, \dots, x_n, 0) = s_2(x_1, \dots, x_n)$ . Since  $s$  vanishes on  $\{0, 1\}^{n+1}$ , this implies that  $s_2$  vanishes on  $\{0, 1\}^n$  as well. By the inductive hypothesis,  $s_2$  is the identically zero polynomial. Now fix  $x_{n+1} = 1$ . Then  $s(x_1, \dots, x_n, 1) = s_1(x_1, \dots, x_n)$ . Again, since  $s$  vanishes on  $\{0, 1\}^{n+1}$ , this implies that  $s_1$  vanishes on  $\{0, 1\}^n$  as well. Again by the inductive hypothesis,  $s_1$  is the identically zero polynomial. This shows that  $s$  is the identically zero polynomial, completing the induction.

Thus,  $r$  is the zero polynomial. Applying this fact to Equation (130), we arrive at the statement in the proposition.  $\square$

## 10.4.2 The PCP

**The problem.** The input to the PCP verifier is a tuple  $(\mathcal{D}, n, T, Q, \sigma, \gamma, x, y)$ . Here,  $\mathcal{D}$  is a decider,  $n, T, Q, \sigma$  and  $\gamma$  are integers with  $Q \leq T$  and  $|\mathcal{D}| \leq \sigma$ , and  $x$  and  $y$  are a pair of strings of length at most  $Q$  each. The goal of the verifier is to check whether there exists two strings  $a_{\text{prefix}}$  and  $b_{\text{prefix}}$  of length at most  $T$  such that  $\mathcal{D}$  halts on input  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  in time at most  $T$ . To do that the verifier makes random queries to a specially encoded *PCP proof*  $\Pi$ , and decides whether to accept or reject based on the parts of  $\Pi$  that it reads. The proof  $\Pi$  is encoded using a low-degree code whose parameters are controlled by  $\gamma$ : the

larger  $\gamma$ , the smaller the soundness error in testing the code. We first set the parameters used in the PCP construction.

**Definition 10.22** (Parameters for the PCP). For all integers  $n, T, Q, \sigma, \gamma \in \mathbb{N}$  such that  $Q \leq T$  and  $|\mathcal{D}| \leq \sigma$  define the tuple  $\text{pcpparams}(n, T, Q, \sigma, \gamma) = (q, m, d, m', s)$  as follows. Let  $m = m(T, \sigma)$ , and  $s = s(n, t, Q, \sigma)$  be as in Proposition 10.20. Let  $a' > 1$  and  $0 < b' < 1$  be the universal constants from Theorem 7.8. Define the following integers.

1. Let  $m' = 5m + 5 + s$  (recall that by Proposition 10.20,  $s$  is chosen so that  $m'$  is a power of 2).
2. Let  $q = 2^k$  where  $k$  is the smallest odd integer satisfying the following:
  - (a)  $k \geq ((\gamma b' + 3a')/b') \cdot \log s$ .
  - (b)  $(2 + 5k)m'/2^k < 1/2$ .
  - (c)  $km'/2^k \leq s^{-b'\gamma}$ .
  - (d)  $2^k$  is divisible by  $m'$ .
3. Let  $d = k$ .

Given  $n, T, Q, \sigma, \gamma$  represented in binary, the parameter tuple  $\text{pcpparams}(n, T, Q, \sigma, \gamma)$  can be computed in time  $\text{poly}(\log(n), \log(T), \log(Q), \log(\gamma), \log(\sigma))$ .

Next, we define the format of a valid PCP proof, which for our construction consists of evaluation tables of low-degree polynomials.

**Definition 10.23.** Given  $n, T, Q, \sigma, \gamma \in \mathbb{N}$  and  $(q, m, d, m', s) = \text{pcpparams}(n, T, Q, \sigma, \gamma)$ , a *low-degree PCP proof* is a tuple  $\Pi$  of evaluation tables of polynomials  $g_1, \dots, g_5 : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  and  $c_0, \dots, c_{m'} : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  with all polynomials having individual degree at most  $d$ . We divide the  $m'$  input variables of  $c_0, \dots, c_{m'}$  into blocks as follows:

$$\mathbb{F}_q^{m'} \ni z = \left( \underbrace{x_1}_{\mathbb{F}_q^m}, \dots, \underbrace{x_5}_{\mathbb{F}_q^m}, \underbrace{o}_{\mathbb{F}_q^5}, \underbrace{w}_{\mathbb{F}_q^s} \right).$$

**Definition 10.24.** Given a low-degree PCP proof  $\Pi$  and a point  $z = (x_1, \dots, x_5, o, w) \in \mathbb{F}_q^{m'}$ , where  $x_1, \dots, x_5 \in \mathbb{F}_q^m$ ,  $o \in \mathbb{F}_q^5$ , and  $w \in \mathbb{F}_q^s$ , the *evaluation* of  $\Pi$  at  $z$  is given by

$$\text{eval}_z(\Pi) = (\alpha_1, \dots, \alpha_5, \beta_0, \dots, \beta_{m'}) \in \mathbb{F}_q^{6+m'},$$

where  $\alpha_i = g_i(x_i)$  and  $\beta_j = c_j(z)$ .

**Theorem 10.25.** *There exists a Turing machine  $\mathcal{M}_{\text{AR}}$  with the following properties.*

1. **(Input format)** *The input to  $\mathcal{M}_{\text{AR}}$  consists of two parts: a “decider specification” and a “PCP view.”*

- (a) **(Decider specification)** *Let  $\mathcal{D}$  be a decider,  $n, T, Q, \sigma$  and  $\gamma$  be integers with  $Q \leq T$ , and  $\sigma = |\mathcal{D}|$ , and let  $x$  and  $y$  be strings of length at most  $Q$ . Let  $(q, m, d, m', s) = \text{pcpparams}(n, T, Q, \sigma, \gamma)$  be as in Definition 10.22. Then the corresponding decider specification is the tuple*

$$(\mathcal{D}, n, T, Q, \sigma, \gamma, x, y).$$

(b) (**PCP view**) Let  $z \in \mathbb{F}_q^{m'}$  and let  $\Xi \in \mathbb{F}_q^{6+m'}$ . Then the PCP view is the pair  $(z, \Xi)$ .

The Turing machine  $\mathcal{M}_{\text{AR}}$  returns either 1 (accept) or 0 (reject).

2. (**Complexity**):  $\mathcal{M}_{\text{AR}}$  runs in time at most  $\text{poly}(\log(T), \log(n), Q, \sigma, \gamma)$ .

For the remaining items, fix a decider specification  $(\mathcal{D}, n, T, Q, \sigma, \gamma, x, y)$ ; we think of  $\mathcal{M}_{\text{AR}}$  as a function of the PCP view input only.

3. (**Completeness**): Suppose  $a_{\text{prefix}}, b_{\text{prefix}} \in \{0, 1\}^*$  are two strings of length  $\ell_a, \ell_b \leq T$ , respectively, such that  $\mathcal{D}$  halts and accepts in time  $T$  on input  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$ . Let  $M = 2^m$  and write

$$a = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{M/2-\ell_a}) \quad \text{and} \quad b = \text{enc}_{\Gamma}(b_{\text{prefix}}, \sqcup^{M/2-\ell_b}),$$

where recall that by Proposition 10.20,  $m$  is chosen so that  $2T \leq M$ , and  $\sqcup$  is a special symbol that is encoded using two bits; thus  $a$  and  $b$  are each an  $M$ -bit string. Then there exists a low-degree PCP proof (Definition 10.23)  $\Pi = (g_1, \dots, g_5, c_0, \dots, c_{m'})$  with  $g_1 = g_a$  and  $g_2 = g_b$ , the low-degree encodings of  $a$  and  $b$ , respectively (see Section 3.4), which causes  $\mathcal{M}_{\text{AR}}$  to accept with probability 1 over a uniformly random  $z \in \mathbb{F}_q^{m'}$ :

$$\Pr_{z \in \mathbb{F}_q^{m'}} (\mathcal{M}_{\text{AR}}(z, \text{eval}_z(\Pi)) = 1) = 1.$$

4. (**Soundness**): Let  $\Pi = (g_1, \dots, g_5, c_0, \dots, c_{m'})$  be a low-degree PCP proof such that  $\mathcal{M}_{\text{AR}}$  at a uniformly random  $z$  accepts with probability larger than  $p_{\text{sound}} = \frac{1}{2}$ :

$$\Pr_{z \in \mathbb{F}_q^{m'}} (\mathcal{M}_{\text{AR}}(z, \text{eval}_z(\Pi)) = 1) > p_{\text{sound}}.$$

Then there exist strings  $a, b \in \{0, 1\}^M$  with the following properties.

(a) There exist strings  $a_{\text{prefix}}, b_{\text{prefix}} \in \{0, 1\}^*$  of length  $\ell_a, \ell_b$ , respectively, such that

$$a = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{M/2-\ell_a}) \quad \text{and} \quad b = \text{enc}_{\Gamma}(b_{\text{prefix}}, \sqcup^{M/2-\ell_b}).$$

(b)  $\mathcal{D}$  halts in time  $T$  and accepts on input  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$ .

(c)  $a = \text{Dec}(g_1)$  and  $b = \text{Dec}(g_2)$ , where  $\text{Dec}(\cdot)$  is the Boolean decoding map of the low-degree encoding defined in Section 3.4.

It is important to note that the soundness in Theorem 10.25 is *only* against low-degree proofs, which is why we are able to obtain a soundness error independent of  $\gamma$ . The eventual answer-reduced verifier (Theorem 10.27) will combine this PCP with the low-degree test, and the latter will introduce a dependence on  $\gamma$  in the soundness error.

*Proof of Theorem 10.25.* We present the description of the Turing machine  $\mathcal{M}_{\text{AR}}$  in Figure 13.

Before establishing the Complexity, Completeness, and Soundness properties of  $\mathcal{M}_{\text{AR}}$ , we point out several important items. Recall that  $M = 2^m$ .

The Turing machine  $\mathcal{M}_{\text{AR}}$  takes the following as input:

- Decider specification  $(\mathcal{D}, n, T, Q, \sigma, \gamma, x, y)$
- PCP view  $(z, \Xi)$

The Turing machine performs the following steps sequentially:

1. Compute  $\mathcal{C} = \text{PaddedSuccinctDecider}(\mathcal{D}, n, T, Q, \sigma, x, y)$ . The circuit  $\mathcal{C}$  has five  $m$ -bit inputs and five single-bit inputs, and contains at most  $s$  AND and OR gates.
2. Compute the Tseitin formula  $\mathcal{F}$  (Definition 10.4) corresponding to  $\mathcal{C}$ , which is a boolean formula on  $m' = 5m + 5 + s$  variables. Let  $\mathcal{F}_{\text{arith}} : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  denote the arithmetization of  $\mathcal{F}$  (Definition 10.5).
3. Parse  $z = (x, o, w) \in \mathbb{F}_q^{m'}$  where  $x_1, \dots, x_5 \in \mathbb{F}_q^m$ ,  $o \in \mathbb{F}_q^5$ , and  $w \in \mathbb{F}_q^s$ . (Here  $x$  is a variable which should not be confused with an input  $x$  to  $\mathcal{D}$ .) Parse  $\Xi = (\alpha_1, \dots, \alpha_5, \beta_0, \dots, \beta_{m'}) \in \mathbb{F}_q^{6+m'}$ .
4. **(Formula test)** Reject if  $\beta_0 \neq \mathcal{F}_{\text{arith}}(x, o, w) \cdot (\alpha_1 - o_1) \cdots (\alpha_5 - o_5)$ . Otherwise, continue.
5. **(Zero on subcube test)** Reject if  $\beta_0 \neq \sum_{i=1}^{m'} \beta_i \cdot \text{zero}(z_i)$ . Otherwise, accept.

Figure 13: The decision procedure  $\mathcal{M}_{\text{AR}}$ .

1. We recall what it means for the circuit  $\mathcal{C}$  computed in Step 1 of Figure 13 to succinctly describe the 5SAT formula  $\varphi_{\mathcal{C}}$ . Let the inputs of the formula  $\varphi_{\mathcal{C}}$  be strings  $a, b, u_3, u_4, u_5 \in \{0, 1\}^M$ . Then for all  $x_1, \dots, x_5 \in \{0, 1\}^m$  and  $o \in \{0, 1\}^5$ , we have that  $\mathcal{C}(x_1, \dots, x_5, o) = 1$  if and only if  $a_{x_1}^{o_1} \vee b_{x_2}^{o_2} \vee u_{3, x_3}^{o_3} \vee u_{4, x_4}^{o_4} \vee u_{5, x_5}^{o_5}$  is a clause in  $\varphi_{\mathcal{C}}$ , where the coordinates of  $a, b, u_3, u_4, u_5$  are indexed by strings of length  $m$ .

Furthermore, the formula  $\varphi_{\mathcal{C}}$  is related to the decider  $\mathcal{D}$  in the following way. For all  $a, b \in \{0, 1\}^M$ , there exist  $u_3, u_4, u_5 \in \{0, 1\}^M$  such that  $a, b, u_3, u_4, u_5$  satisfy  $\varphi_{\mathcal{C}}$  if and only if there exist  $a_{\text{prefix}}, b_{\text{prefix}} \in \{0, 1\}^*$  of lengths  $\ell_a, \ell_b \leq T$ , respectively, such that

$$a = \text{enc}_{\Gamma}(a_{\text{prefix}}, \sqcup^{M/2-\ell_a}) \quad \text{and} \quad b = \text{enc}_{\Gamma}(b_{\text{prefix}}, \sqcup^{M/2-\ell_b})$$

and  $\mathcal{D}$  accepts  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  in time  $T$ .

2. We recall from Definition 10.4 that the boolean formula  $\mathcal{F}$  is related to  $\mathcal{C}$  in the following way. For all  $x_1, \dots, x_5 \in \{0, 1\}^m$  and  $o \in \{0, 1\}^5$ ,  $\mathcal{C}(x_1, \dots, x_5, o) = 1$  if and only if there exists a  $w \in \{0, 1\}^s$  such that  $\mathcal{F}(x_1, \dots, x_5, o, w) = 1$ . The size of the formula  $\mathcal{F}$  is linear in the size of the circuit  $\mathcal{C}$ , which is  $s$ .
3. We recall from Definition 10.5 that the arithmetization  $\mathcal{F}_{\text{arith}} := \text{arith}_q(\mathcal{F})$  is a function  $\mathcal{F}_{\text{arith}} : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  such that

$$\forall (x, o, w) \in \{0, 1\}^{m'} \quad \mathcal{F}_{\text{arith}}(x, o, w) = \mathcal{F}(x, o, w). \quad (131)$$

By Proposition 10.6,  $\mathcal{F}_{\text{arith}}$  is an individual degree 2 polynomial.

**Complexity.** We bound the complexity of the Turing machine  $\mathcal{M}_{\text{AR}}$ . From Proposition 10.20, Step 1 takes time  $\text{poly}(\log(T), \log(n), Q, \sigma)$ . Computing the Tseitin formula in Step 2 takes time that is polynomial in the size of the circuit  $\mathcal{C}$ , which is  $\text{poly}(s) = \text{poly}(\log(T), \log(n), Q, \sigma)$ . The Formula Test, Step 4, requires computing the arithmetization  $\mathcal{F}_{\text{arith}}$  at a point  $z \in \mathbb{F}_q^{m'}$ , which takes time  $\text{poly}(s, \log q)$ . The Zero on Subcube Test, Step 5, takes time  $\text{poly}(m', \log q)$  to compute a sum of  $m'$  products of  $\mathbb{F}_q$  elements (evaluating  $\text{zero}(\cdot)$  takes time  $\text{poly}(\log q)$ ). Thus the complexity of  $\mathcal{M}_{\text{AR}}$  is bounded by

$$\text{poly}(\log(T), \log(n), Q, \sigma, \gamma)$$

using our choice of pcpparams from Definition 10.22. (The dependence on  $\gamma$  comes from the setting of  $q$ .)

**Completeness.** We now establish the Completeness property of  $\mathcal{M}_{\text{AR}}$  by concocting a PCP proof  $\Pi$  that is accepted with probability 1.

Let  $a, b \in \{0, 1\}^M$  be as specified in Item 3 of Theorem 10.25. Then by definition of the formula  $\varphi_{\mathcal{C}}$  and the assumption that  $\mathcal{D}$  halts and accepts in time  $T$  on input  $(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$ , there exist strings  $u_3, u_4, u_5 \in \{0, 1\}^M$  such that  $(a, b, u_3, u_4, u_5)$  satisfies  $\varphi_{\mathcal{C}}$ .

Let  $g_1, g_2$  denote the low-degree encodings of  $a$  and  $b$ , respectively (see Section 3.4 for definition of low-degree encoding), and let  $g_3, g_4, g_5$  denote the low-degree encodings of  $u_3, u_4$ , and  $u_5$ , respectively. In particular,  $g_1, \dots, g_5$  are  $m$ -variate multilinear polynomials, and for all  $x_1, \dots, x_5 \in \{0, 1\}^m$ ,

$$g_1(x_1) = a_{x_1}, \quad g_2(x_2) = b_{x_2}, \quad g_3(x_3) = u_{3,x_3}, \quad g_4(x_4) = u_{4,x_4}, \quad g_5(x_5) = u_{5,x_5},$$

where we index the coordinates of  $a, b, u_3, u_4, u_5$  by  $m$ -bit strings in the natural way.

Next, define the polynomial  $c_0 : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  as

$$c_0(x, o, w) = \mathcal{F}_{\text{arith}}(x, o, w) \cdot (g_1(x_1) - o_1) \cdots (g_5(x_5) - o_5).$$

Note that  $c_0$  has individual degree 3, since as recalled in item 3 above  $\mathcal{F}_{\text{arith}}(x, o, w)$  has individual degree 2 and each of the  $g_i$ 's are multilinear).

We now show that  $c_0(x, o, w) = 0$  for all  $(x, o, w) \in \{0, 1\}^{m'}$ . By construction, the arithmetization  $\mathcal{F}_{\text{arith}}(x, o, w)$  is either 0 or 1. If it is 0, then we are done. If it is 1, then this means that  $\mathcal{C}(x, o) = 1$ , which by definition means that  $a_{x_1}^{o_1} \vee b_{x_2}^{o_2} \vee u_{3,x_3}^{o_3} \vee u_{4,x_4}^{o_4} \vee u_{5,x_5}^{o_5}$  is a clause in the formula  $\varphi_{\mathcal{C}}$ . But since  $(a, b, u_3, u_4, u_5)$  satisfies  $\varphi_{\mathcal{C}}$ , this means that at least one of  $a_{x_1} = o_1$ ,  $b_{x_2} = o_2$ ,  $u_{3,x_3} = o_3$ ,  $u_{4,x_4} = o_4$ , or  $u_{5,x_5} = o_5$ . But this means that the product  $(g_1(x_1) - o_1) \cdots (g_5(x_5) - o_5) = 0$ , by definition of the  $g_i$ 's. Thus  $c_0(x, o, w) = 0$ .

Thus, by Proposition 10.21, there exist polynomials  $c_1, \dots, c_{m'} : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  of individual degree at most 3 that certify that  $c_0$  vanishes on  $\{0, 1\}^{m'}$ . In other words, for all  $z \in \mathbb{F}_q^{m'}$  (not just over the boolean cube), we have

$$c_0(z) = \sum_{i=1}^{m'} c_i(z) \cdot \text{zero}(z_i) \tag{132}$$

where  $\text{zero}(x) = x(1 - x)$ .

Let the PCP proof  $\Pi$  be the collection of polynomials  $(g_1, \dots, g_5, c_0, c_1, \dots, c_{m'})$ , where the  $g_i$ 's act on disjoint variables and the  $c_i$ 's act on all of them. Let  $z \in \mathbb{F}_q^{m'}$  and let  $\Xi = (\alpha_1, \dots, \alpha_5, \beta_0, \beta_1, \dots, \beta_{m'}) =$

$\text{eval}_z(\Pi)$ . The PCP view  $(z, \Xi)$  passes the Formula Test always, because  $\beta_0 = c_0(z)$  and  $\alpha_i = g_i(z)$  for  $i \in \{1, 2, \dots, 5\}$ . The PCP view also passes the Zero on Subcube Test, because of Equation (132).

Thus the PCP view is accepted by the Turing machine  $\mathcal{M}_{\text{AR}}$  with probability 1. This establishes the Completeness property.

**Soundness.** Fix a low-degree PCP proof  $\Pi = (g_1, \dots, g_5, c_0, \dots, c_{m'})$  such that the PCP view  $(z, \Xi)$  for  $\Xi = \text{eval}_z(\Pi)$  causes  $\mathcal{M}_{\text{AR}}$  to accept with probability greater than  $p_{\text{sound}}$ . This, in particular, implies that the Formula Test is satisfied with high probability. With probability greater than  $p_{\text{sound}}$  over the choice of  $z \sim \mathbb{F}_q^{m'}$ , we have

$$c_0(z) = \mathcal{F}_{\text{arith}}(z) \cdot (g_1(x_1) - o_1) \cdots (g_5(x_5) - o_5)$$

Note that  $c_0(z)$ , by definition of low-degree PCP proof, has individual degree at most  $d$ . The right-hand side has individual degree at most  $2 + 5d$ , because  $\mathcal{F}_{\text{arith}}$  has individual degree 2 and the  $g_i$ 's have individual degree at most  $d$ . Thus both sides have total degree at most  $(2 + 5d)m'$ . Suppose  $c_0(z)$  was not identical to  $\mathcal{F}_{\text{arith}}(z) \cdot (g_1(x_1) - o_1) \cdots (g_5(x_5) - o_5)$ . Then the Schwartz-Zippel lemma implies that the probability they agree on a randomly chosen  $z$  is at most  $(2 + 5d)m'/q$ . By our choice of  $q$ ,  $m'$ , and  $d$  in Definition 10.22, this is less than  $p_{\text{sound}}$ , which is a contradiction. Thus  $c_0(z)$  is equal to  $\mathcal{F}_{\text{arith}}(z) \cdot (g_1(x_1) - o_1) \cdots (g_5(x_5) - o_5)$  for all  $z \in \mathbb{F}_q^{m'}$ .

Next, we consider the Zero on Subcube Test. With probability greater than  $p_{\text{sound}}$  over the choice of  $z \sim \mathbb{F}_q^{m'}$ , we have

$$c_0(z) = \sum_{i=1}^{m'} c_i(z) \cdot \text{zero}(z_i).$$

Again, the polynomials on both sides of the equation have individual degree at most  $d + 2$ , and therefore total degree at most  $(2 + d)m'$ . By the Schwartz-Zippel lemma, the probability that both sides would agree on the evaluation of a random  $z$ , if they weren't equal polynomials, would be at most  $(2 + d)m'/q$ , which is also less than  $p_{\text{sound}}$ . Thus  $c_0(z) = \sum_{i=1}^{m'} c_i(z) \cdot \text{zero}(z_i)$  for all  $z \in \mathbb{F}_q^{m'}$ . This in particular implies that  $c_0(z)$  vanishes on the subcube  $\{0, 1\}^{m'}$ .

We can now decode assignments  $a, b, u_3, u_4, u_5 \in \{0, 1\}^M$  that satisfy the formula  $\varphi_C$ . Let  $a = \text{Dec}(g_1), b = \text{Dec}(g_2), u_3 = \text{Dec}(g_3), u_4 = \text{Dec}(g_4), u_5 = \text{Dec}(g_5)$  where  $\text{Dec}(\cdot)$  is the decoding map defined in Section 3.4. The fact that  $c_0(z)$  vanishes on the subcube  $\{0, 1\}^{m'}$  implies that all clauses of  $\varphi_C$  are satisfied by  $a, b, u_3, u_4, u_5$ . By construction of  $\varphi_C$ , this implies that there exists  $a_{\text{prefix}}, b_{\text{prefix}}$  such that  $\mathcal{D}(n, x, y, a_{\text{prefix}}, b_{\text{prefix}})$  accepts in time  $T$ . This completes the proof of the Soundness property, and the proof of the Theorem.  $\square$

## 10.5 A normal form verifier for the PCP

In this section we show how to convert the PCP from Section 10.4 into a normal form verifier. This results in an ‘‘answer reduction’’ scheme: a way to map a verifier  $\mathcal{V}$  into a new (typed) verifier  $\hat{\mathcal{V}}^{\text{AR}}$  with a smaller answer size.

Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier and  $(\lambda, \mu, \gamma)$  a tuple of integers. In the rest of this section we define the (typed) answer-reduced verifier  $\hat{\mathcal{V}}^{\text{AR}} = (\hat{\mathcal{S}}^{\text{AR}}, \hat{\mathcal{D}}^{\text{AR}})$  associated with  $\mathcal{V}$  and parameters  $(\lambda, \mu, \gamma)$ . Completeness, complexity and soundness of the construction are shown in the following sections.

### 10.5.1 Parameters and notation

We establish some notation and parameters that are used throughout Section 10.5. First, define the functions

$$T(n) = (2^{\lambda n})^\mu \quad \text{and} \quad Q(n) = (\lambda n)^\mu . \quad (133)$$

In the main theorem of this section, Theorem 10.27, we assume that the input verifier  $\mathcal{V}$  satisfies

$$\text{TIME}_{\mathcal{D}}(n) \leq T(n) \quad \text{and} \quad \text{TIME}_{\mathcal{S}}(n) \leq Q(n) . \quad (134)$$

In what follows we write  $T$  and  $Q$  as free parameters (though they are implicitly functions of the index  $n$ ).

Next, for all integers  $n \in \mathbb{N}$  define the PCP parameters  $(q, m, d, m', s) = \text{pcpparams}(n, T, Q, \sigma, \gamma)$  where  $\sigma = |\mathcal{D}|$ . Note that the parameters  $(q, m, d, m', s)$  are all implicitly functions of  $n$ .

Recall from Definition 4.16 the notation  $\mu_{\mathcal{S}}$  denoting the distribution over pairs of questions  $(x_A, x_B)$  generated by  $\mathcal{S}$ . We use  $\mu_{\mathcal{S},A}$  to indicate the marginal distribution of  $\mu_{\mathcal{S}}$  on the first question  $x_A$ , and  $\text{SUPP}(\mu_{\mathcal{S}})$  to indicate the set of question pairs that have nonzero probability under  $\mu_{\mathcal{S}}$ .

**Remark 10.26.** *Throughout Section 10.5, for convenience we often identify the label A with 1 and B with 2. For example,  $g_A$  is another label for the polynomial  $g_1$ .*

### 10.5.2 The answer-reduced verifier

Let  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  be a normal form verifier and  $(\lambda, \mu, \gamma)$  integers. All other required parameters and notation are introduced in Section 10.5.1.

**Sampler.** In this section we define the (typed) sampler  $\hat{\mathcal{S}}^{\text{AR}}$  for the (typed) answer reduced verifier  $\hat{\mathcal{V}}^{\text{AR}}$ .

We first give an intuitive description of the sampler, and then proceed to give a formal definition. The sampler  $\hat{\mathcal{S}}^{\text{AR}}$  is a *product* of the oracularized sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  corresponding to  $\mathcal{S}$ , and a typed PCP sampler  $\hat{\mathcal{S}}^{\text{PCP}}$ . The corresponding question distribution  $\mu_{\hat{\mathcal{S}}^{\text{AR}}}$  is then a product distribution  $\mu_{\hat{\mathcal{S}}^{\text{ORAC}}} \times \mu_{\hat{\mathcal{S}}^{\text{PCP}}}$ .

The PCP distribution  $\mu_{\hat{\mathcal{S}}^{\text{PCP}}}$  corresponds to six copies of the classical low-degree test question distribution (see Section 7.1.1). Recall that the classical low-degree test is a nonlocal game that checks whether the players' answers are consistent with low-degree polynomials. The answer-reduced verifier will use the classical low-degree test to check that the players respond with evaluations of low-degree polynomials  $g_1, \dots, g_5, c_0, \dots, c_{m'}$ , and then process the evaluations using the PCP verifier  $\mathcal{M}_{\text{AR}}$  from Theorem 10.25. Five of the six copies are meant to certify the “low-degreeness” of  $g_1, \dots, g_5$ , and the sixth copy is used to certify the “low-degreeness” of the entire bundle of polynomials  $g_1, \dots, g_5, c_0, \dots, c_{m'}$ . The reason that the  $g_i$ 's are checked individually is to ensure that they only depend on certain blocks of input variables, because ultimately the soundness of the answer-reduced verifier reduces to the soundness of the PCP verifier  $\mathcal{M}_{\text{AR}}$ , and Theorem 10.25 assumes that the polynomials  $g_1, \dots, g_5$  of the PCP proof only depend on certain subsets of variables.

We now formally define the typed PCP sampler  $\hat{\mathcal{S}}^{\text{PCP}}$ . The type set is

$$\mathcal{T}^{\text{PCP}} = \{\text{POINT}_1, \dots, \text{POINT}_6\} \cup \{\text{ALINE}_1, \dots, \text{ALINE}_6\} \cup \{\text{DLINE}_1, \dots, \text{DLINE}_6\} ,$$

and the type graph  $G^{\text{PCP}} = (\mathcal{T}^{\text{PCP}}, E^{\text{PCP}})$  uses the complete edge set  $E^{\text{PCP}} = \mathcal{T}^{\text{PCP}} \times \mathcal{T}^{\text{PCP}}$ . The ambient vector space for the sampler is

$$V^{\text{PCP}} = \left( \bigoplus_{i=1}^5 V_{i,X} \oplus V_{i,I} \oplus V_{i,V} \right) \oplus V_{\text{AUX},X} \oplus V_{\text{AUX},I} \oplus V_{\text{AUX},V} , \quad (135)$$



where the spaces  $V_{i,X}, V_{i,V}$  are each isomorphic to  $\mathbb{F}_q^m$ , the space  $V_{i,I}$  is isomorphic to  $\mathbb{F}_q$ , the spaces  $V_{\text{AUX},X}, V_{\text{AUX},V}$  are each isomorphic to  $\mathbb{F}_q^{5+s}$ , and the space  $V_{\text{AUX},I}$  is isomorphic to  $\mathbb{F}_q$ . In addition, define the following direct sums:

$$\begin{aligned} V_{6,X} &= \left( \bigoplus_{i=1}^5 V_{i,X} \right) \oplus V_{\text{AUX},X}, \\ V_{6,I} &= \left( \bigoplus_{i=1}^5 V_{i,I} \right) \oplus V_{\text{AUX},I}, \\ V_{6,V} &= \left( \bigoplus_{i=1}^5 V_{i,V} \right) \oplus V_{\text{AUX},V}. \end{aligned}$$

For all  $i \in \{1, 2, \dots, 6\}$ , we call the space  $V_{i,X}$  the  $i$ -th *point register*, the space  $V_{i,I}$  the  $i$ -th *coordinate register*, and the space  $V_{i,V}$  the  $i$ -th *direction register* (compare this with the subspaces defined in Section 7.1.1).

For every type  $t \in \mathcal{T}^{\text{PCP}}$ , we define the following CL functions on  $V^{\text{PCP}}$ :

- For the types  $t = \text{POINT}_i$  for  $i \in \{1, \dots, 6\}$ , the corresponding 1-level CL function  $L_{\text{POINT}_i}$  is identical to the CL function  $L_{\text{POINT}}$  defined in Equation (48), but acts on the subspace  $V_{i,X} \oplus V_{i,I} \oplus V_{i,V}$  (and zeroes out all the other registers).
- For the types  $t = \text{ALINE}_i$  for  $i \in \{1, \dots, 6\}$ , the corresponding 2-level CL function  $L_{\text{ALINE}_i}$  is identical to the CL function  $L_{\text{ALINE}}$  defined in Equation (49), but acts on the subspaces  $V_{i,X} \oplus V_{i,I} \oplus V_{i,V}$ .
- For the types  $t = \text{DLINE}_i$  for  $i \in \{1, \dots, 6\}$ , the corresponding 3-level CL function  $L_{\text{DLINE}_i}$  is identical to the CL function  $L_{\text{DLINE}}$  defined in Equation (51), but acts on the subspaces  $V_{i,X} \oplus V_{i,I} \oplus V_{i,V}$ .

Observe that for  $i \in \{1, \dots, 5\}$ , the CL distributions  $\mu_{L_{\text{ALINE}_i}, L_{\text{POINT}_i}}$  and  $\mu_{L_{\text{DLINE}_i}, L_{\text{POINT}_i}}$  correspond to the question distributions of the classical low-degree test parameterized by  $q$  and  $m$  (see Section 7.1.1 for the definition of the low-degree test distributions, which only depend on the first two arguments of the 4-tuple  $\text{ldparams}$ ), and the CL distributions  $\mu_{L_{\text{ALINE}_6}, L_{\text{POINT}_6}}$  and  $\mu_{L_{\text{DLINE}_6}, L_{\text{POINT}_6}}$  correspond to the classical low-degree test parameterized by  $q$  and  $m'$ .

We finally give the formal definition of the sampler  $\hat{\mathcal{S}}^{\text{AR}}$ . The type set of  $\hat{\mathcal{S}}^{\text{AR}}$  is  $\mathcal{T}^{\text{AR}} = \mathcal{T}^{\text{ORAC}} \times \mathcal{T}^{\text{PCP}}$  and the type graph is  $G^{\text{AR}} = G^{\text{ORAC}} \times G^{\text{PCP}}$  with edge set

$$E^{\text{AR}} = \{ \{(u, v), (u', v')\} : \{u, u'\} \in E^{\text{ORAC}} \text{ and } \{v, v'\} \in E^{\text{PCP}} \}.$$

Let  $V^{\text{ORAC}}$  denote the ambient space of the oracularized sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$ . The ambient space of  $\hat{\mathcal{S}}^{\text{AR}}$  is then the direct sum  $V^{\text{ORAC}} \oplus V^{\text{PCP}}$ . For all type pairs  $(t_{\text{ORAC}}, t_{\text{PCP}}) \in \mathcal{T}^{\text{AR}}$ , the corresponding CL function  $L_{t_{\text{ORAC}}, t_{\text{PCP}}}$  is simply the direct sum of the CL functions  $L_{t_{\text{ORAC}}}$  (coming from  $\hat{\mathcal{S}}^{\text{ORAC}}$ ) and  $L_{t_{\text{PCP}}}$  (coming from  $\hat{\mathcal{S}}^{\text{PCP}}$ ). Thus one can see that the distribution corresponding to  $\hat{\mathcal{S}}^{\text{AR}}$  is the product distribution  $\mu_{\hat{\mathcal{S}}^{\text{ORAC}}} \times \mu_{\hat{\mathcal{S}}^{\text{PCP}}}$ .

Note that  $\hat{\mathcal{S}}^{\text{AR}}$  is an  $\max\{\ell, 3\}$ -level sampler, where  $\ell$  is the number of levels of the sampler  $\mathcal{S}$ .

**Decider.** The decider  $\hat{\mathcal{D}}^{\text{AR}}$  is described in Fig. 14.

Type	Question Format	Answer Format
POINT <sub><i>i</i></sub> for $i = 1, \dots, 5$	$y_i \in \mathbb{F}_q^m$	$\alpha_i \in \mathbb{F}_q$
ALINE <sub><i>i</i></sub> for $i = 1, \dots, 5$	$v_i \in \mathbb{F}_q^m \times \mathbb{F}_q$	$h_i : \mathbb{F}_q \rightarrow \mathbb{F}_q$
DLINE <sub><i>i</i></sub> for $i = 1, \dots, 5$	$v_i \in \mathbb{F}_q^m \times \mathbb{F}_q \times \mathbb{F}_q^m$	$h_i : \mathbb{F}_q \rightarrow \mathbb{F}_q$
POINT <sub>6</sub>	$z = (y, o, w) \in \mathbb{F}_q^{m'}$	$(\alpha'_1, \dots, \alpha'_5, \beta_0, \dots, \beta_{m'}) \in \mathbb{F}_q^{m'+6}$
ALINE <sub>6</sub>	$v \in \mathbb{F}_q^{m'} \times \mathbb{F}_q$	$(h'_1, \dots, h'_5, f_0, \dots, f_{m'}) : \mathbb{F}_q \rightarrow \mathbb{F}_q^{m'+6}$
DLINE <sub>6</sub>	$v \in \mathbb{F}_q^{m'} \times \mathbb{F}_q \times \mathbb{F}_q^{m'}$	$(h'_1, \dots, h'_5, f_0, \dots, f_{m'}) : \mathbb{F}_q \rightarrow \mathbb{F}_q^{m'+6}$

Table 1: Question and answer formats for types in  $\mathcal{T}^{\text{PCP}}$ .

On input  $(n, t_A, x_A, t_B, x_B, a_A, a_B)$ , the decider  $\hat{\mathcal{D}}^{\text{AR}}$  parses  $t_A, t_B$  as  $(t_{Q,A}, t_{\Pi,A})$ , and  $(t_{Q,B}, t_{\Pi,B})$  respectively in  $\mathcal{T}^{\text{ORAC}} \times \mathcal{T}^{\text{PCP}}$ , parses  $x_A$  and  $x_B$  as  $(x_{Q,A}, x_{\Pi,A})$  and  $(x_{Q,B}, x_{\Pi,B})$  respectively. The answer format depends only on  $t_{\Pi,A}$  and  $t_{\Pi,B}$  respectively and is as indicated in Table 1. The decider performs the following steps sequentially, for all  $w \in \{A, B\}$ :

1. **(Global consistency check)**: If  $t_A = t_B$ , reject if  $a_A \neq a_B$ .
2. **(Input consistency check)**: If  $t_{Q,w} = \text{ORACLE}$  and  $t_{Q,\bar{w}} = v \in \{A, B\}$ , and if  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_6, \text{POINT}_v)$ , reject if  $\alpha_v \neq \alpha'_v$  (where  $A \leftrightarrow 1$  and  $B \leftrightarrow 2$ , as per Remark 10.26).
3. **(Input low degree test)** If  $t_{Q,w} = t_{Q,\bar{w}} = v \in \{A, B\}$ , and if  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_v, \text{ALINE}_v)$  (resp. DLINE<sub>*v*</sub> instead of ALINE<sub>*v*</sub>), execute  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  on input  $(\text{POINT}, x_{\Pi,w}, \text{ALINE}, x_{\Pi,\bar{w}}, a_w, a_{\bar{w}})$  (resp. DLINE instead of ALINE), where  $\text{ldparams} = (q, m, d, 1)$ . Reject if  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  rejects.
4. **(Proof encoding checks)**: If  $t_{Q,w} = t_{Q,\bar{w}} = \text{ORACLE}$ ,
  - (a) (Consistency test) If  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_i, \text{POINT}_6)$  for some  $i \in \{3, \dots, 5\}$ , reject if  $\alpha_i \neq \alpha'_i$ .
  - (b) (Individual low degree test) If  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_i, \text{ALINE}_i)$  (resp. DLINE<sub>*i*</sub>) for some  $i \in \{3, \dots, 5\}$ , execute  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  on input  $(\text{POINT}, x_{\Pi,w}, \text{ALINE}, x_{\Pi,\bar{w}}, a_w, a_{\bar{w}})$  (resp. DLINE). Reject if  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  rejects.
  - (c) (Simultaneous low degree test) If  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_6, \text{ALINE}_6)$  (resp. DLINE<sub>6</sub>), execute  $\mathcal{D}_{\text{ldparams}'}^{\text{LD}}$  on input  $(\text{POINT}, x_{\Pi,w}, \text{ALINE}, x_{\Pi,\bar{w}}, a_w, a_{\bar{w}})$  (resp. DLINE), where  $\text{ldparams}' = (q, m', d, m' + 6)$ . Reject if  $\mathcal{D}_{\text{ldparams}'}^{\text{LD}}$  rejects.
5. **(Game check)**: If  $t_{Q,w} = \text{ORACLE}$ , then for  $v \in \{A, B\}$ , compute  $x_{w,v} = L^v(x_{Q,w})$ . If  $t_{\Pi,w} = \text{POINT}_6$ , reject if  $\mathcal{M}_{\text{AR}}((\mathcal{D}, n, T, Q, \gamma, x_{w,A}, x_{w,B}), (z, a_w))$  rejects. Otherwise, accept.

Figure 14: The decision procedure  $\hat{\mathcal{D}}^{\text{AR}}$ . Parameters  $T, Q, q, m, m', d, \gamma$  are defined in Section 10.5.1.

### 10.5.3 Main theorem for answer reduction

**Theorem 10.27.** *There exists a polynomial time Turing machine  $\text{ComputeAnsRedVerifier}$  that, on input  $(\mathcal{V}, \lambda, \mu, \gamma)$  where  $\mathcal{V}$  is an  $\ell$ -level normal form verifier and  $\lambda, \mu, \gamma \in \mathbb{N}$ , outputs the description of the answer reduced verifier  $\mathcal{V}^{\text{AR}} = (\mathcal{S}^{\text{AR}}, \mathcal{D}^{\text{AR}})$ . Let  $T(n)$  and  $Q(n)$  be the functions specified in Eq. (133),*

and suppose that  $\mathcal{V}$  satisfies the complexity conditions specified in Eq. (134). Then  $\mathcal{V}^{\text{AR}}$  is  $\max\{\ell + 2, 5\}$ -level, has time complexity bounds

$$\begin{aligned}\text{TIME}_{\mathcal{S}^{\text{AR}}}(n) &= O(\text{TIME}_{\hat{\mathcal{S}}^{\text{ORAC}}}(n) + \text{poly}(\log(T(n)), |\mathcal{D}|, \gamma)) = \text{poly}((\lambda n)^\mu, |\mathcal{D}|, \gamma), \\ \text{TIME}_{\mathcal{D}^{\text{AR}}}(n) &= \text{poly}(\log(T(n)), Q(n), |\mathcal{D}|, \gamma) = \text{poly}((\lambda n)^\mu, |\mathcal{D}|, \gamma),\end{aligned}$$

and furthermore the sampler  $\mathcal{S}^{\text{AR}}$  only depends on  $\mathcal{S}$ , the parameter tuple  $(\lambda, \mu, \gamma)$ , and the description length  $|\mathcal{D}|$  (but nothing else about  $\mathcal{D}$ ). Moreover, there exists a function

$$\delta(\varepsilon, n) = a\gamma^a((\lambda \cdot |\mathcal{D}| \cdot n)^{a\mu} \cdot \varepsilon^b + (\lambda \cdot |\mathcal{D}| \cdot n)^{-\mu b \gamma})$$

for universal constants  $a > 0$ ,  $0 < b < 1$ , such that the following hold for all  $n \geq 2$ :

1. (**Completeness**) If  $\mathcal{V}_n$  has a projective, consistent, and commuting (PCC) strategy of value 1, then  $\mathcal{V}_n^{\text{AR}}$  has an SPCC strategy with value 1.
2. (**Soundness**) If  $\text{val}^*(\mathcal{V}_n^{\text{AR}}) > 1 - \varepsilon$  for some  $\varepsilon > 0$  then  $\text{val}^*(\mathcal{V}_n) \geq 1 - \delta(\varepsilon, n)$ .
3. (**Entanglement**) Let  $\mathcal{E}(\cdot)$  be as defined in Definition 5.12. Then for all  $\varepsilon \geq 0$ ,

$$\mathcal{E}(\mathcal{V}_n^{\text{AR}}, 1 - \varepsilon) \geq \mathcal{E}(\mathcal{V}_n, 1 - \delta(\varepsilon, n)).$$

Observe that  $\text{TIME}_{\mathcal{D}^{\text{AR}}}(n)$  is polynomial in the *logarithm* of  $T(n)$ , the runtime of the decider of  $\mathcal{V}$ , achieving the desired reduction in runtime and answer complexity.

*Proof.* The Turing machine `ComputeAnsRedVerifier` can be described as follows:

1. From the description of  $\mathcal{V}$ , compute the description of the (typed) oracularized verifier  $\hat{\mathcal{V}}^{\text{ORAC}} = (\hat{\mathcal{S}}^{\text{ORAC}}, \hat{\mathcal{D}}^{\text{ORAC}})$  using the Turing machine `ComputeOracleVerifier` from Theorem 9.1.
2. From the description of the typed sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  and the parameter tuple  $(\lambda, \mu, \gamma)$ , compute the descriptions of the typed sampler  $\hat{\mathcal{S}}^{\text{AR}}$  as described in Section 10.5.2.
3. From the descriptions of the decider  $\mathcal{D}$  and the parameter tuple  $(\lambda, \mu, \gamma)$ , compute the descriptions of the typed decider  $\hat{\mathcal{D}}^{\text{AR}}$  as described in Figure 14.
4. Compute the detyped verifier  $\mathcal{V}^{\text{AR}} = (\mathcal{S}^{\text{AR}}, \mathcal{D}^{\text{AR}}) = \text{Detype}(\hat{\mathcal{V}}^{\text{AR}})$ .

The Turing machine `ComputeAnsRedVerifier` takes time that is  $\text{poly}(|\mathcal{V}|, \log \lambda, \log \mu, \log \gamma)$ , because each step, including the detyping procedure, runs in time that is polynomial in the length of the input  $(\mathcal{V}, \lambda, \mu, \gamma)$ .

**Properties of the sampler  $\mathcal{S}^{\text{AR}}$ .** It can be verified via inspection that the typed sampler  $\hat{\mathcal{S}}^{\text{AR}}$  depends on  $\hat{\mathcal{S}}^{\text{ORAC}}$  (which itself only depends on  $\mathcal{S}$ ; see Theorem 9.1), and  $\hat{\mathcal{S}}^{\text{PCP}}$  (which only depends on the parameter tuple  $(\lambda, \mu, \gamma)$  and the description length  $|\mathcal{D}|$ ). Using Theorem 9.1,  $\hat{\mathcal{S}}^{\text{ORAC}}$  is a  $\ell$ -level typed sampler and, therefore,  $\hat{\mathcal{S}}^{\text{AR}}$  is a  $\max\{\ell, 3\}$ -level sampler as  $\hat{\mathcal{S}}^{\text{PCP}}$  is a typed 3-level sampler. Using Lemma 6.18 for the detyping it follows that  $\mathcal{S}^{\text{AR}}$  is a  $\max\{\ell + 2, 5\}$ -level typed sampler.

**Complexity.** In addition to the running time of  $\hat{\mathcal{S}}^{\text{ORAC}}$ , the time complexity of  $\hat{\mathcal{S}}^{\text{AR}}$  also includes the running time of  $\hat{\mathcal{S}}^{\text{PCP}}$ , which is dominated by the complexity of computing the CL functions  $L_{\text{POINT}_i}$ ,  $L_{\text{ALINE}_i}$ , and  $L_{\text{DLINE}_i}$ . From Lemma 7.9, this takes time

$$\text{poly}(m', \log q) = \text{poly}(\gamma, s) = \text{poly}(\log T, \log n, Q, |\mathcal{D}|, \gamma) = \text{poly}(\log T, |\mathcal{D}|, \gamma),$$

where the last equality uses the formulas (133). Therefore the time complexity of  $\hat{\mathcal{S}}^{\text{AR}}$  is

$$O(\text{TIME}_{\hat{\mathcal{S}}^{\text{ORAC}}}(n) + \text{poly}(\log T(n), |\mathcal{D}|, \gamma)).$$

The complexity of  $\mathcal{S}^{\text{AR}} = \text{Detype}(\hat{\mathcal{S}}^{\text{AR}})$  then follows by Lemma 6.18 and Equation (134).

The decider  $\hat{\mathcal{D}}^{\text{AR}}$  executes subroutines  $\mathcal{D}^{\text{LD}}$  and  $\mathcal{M}_{\text{AR}}$ . The runtime of  $\mathcal{D}^{\text{LD}}$  is  $\text{poly}(m, d, m', \log q)$ . The runtime of  $\mathcal{M}_{\text{AR}}$  is given in Theorem 10.25; for  $T$  and  $Q$  as in (133) it is  $\text{poly}(\log T, \log n, \log q, Q, |\mathcal{D}|)$ .

In addition to these subroutines,  $\hat{\mathcal{D}}^{\text{AR}}$  computes the CL functions  $L^{\text{A}}$  and  $L^{\text{B}}$  of the oracularized sampler  $\hat{\mathcal{S}}^{\text{ORAC}}$  in Step 5 of Figure 14. By Theorem 9.1, the complexity of  $\hat{\mathcal{S}}^{\text{ORAC}}$  is polynomial in the time complexity of  $\mathcal{S}$ , which by Equation (134) is at most  $Q(n)$ .

Thus the overall time complexity of  $\hat{\mathcal{D}}^{\text{AR}}$ , using both the PCP parameter settings of Definition 10.22 and the assumptions of Equation (134), is  $\text{poly}((\lambda n)^\mu, |\mathcal{D}|, \gamma)$ . The complexity of  $\text{Detype}(\hat{\mathcal{D}}^{\text{AR}})$  follows by Lemma 6.18.

**Completeness, Soundness, and Entanglement.** The Completeness property is established in Section 10.6 and the Soundness and Entanglement properties are proven in Section 10.7.  $\square$

## 10.6 Completeness of the answer-reduced verifier

We establish the Completeness property of the answer reduced verifier as stated in Theorem 10.27.

*Proof of the Completeness part of Theorem 10.27.* We establish the Completeness property of  $\hat{\mathcal{V}}^{\text{AR}}$ , which by the Detying Lemma (Lemma 6.18) establishes the Completeness property of  $\mathcal{V}^{\text{AR}}$ .

Let  $n \geq 1$  be an index for  $\hat{\mathcal{V}}^{\text{AR}}$ . Let  $\mathcal{S}$  be a PCC strategy for  $\mathcal{V}_n$  with value 1. By Theorem 9.1 it follows that there exists a symmetric PCC strategy  $\mathcal{S}^{\text{ORAC}}$  using a state  $|\psi\rangle$  with value 1 for  $\hat{\mathcal{V}}_n^{\text{ORAC}}$ . We define a strategy  $\mathcal{S}^{\text{AR}}$  for the typed verifier  $\hat{\mathcal{V}}_n^{\text{AR}}$  as follows. The shared state is  $|\psi\rangle$ . Given the index  $n$  and  $(\lambda, \mu, \gamma)$ , each player can compute  $(q, m, d, m', s) = \text{pcpparams}(n, T, Q, \sigma, \gamma)$  (see Definition 10.22), where  $T = T(n)$  and  $Q = Q(n)$  are as in (133), and  $\sigma = |\mathcal{D}|$ . Let  $M = 2^m$ .

1. On receipt of a question  $((t_Q, t_\Pi), (x_Q, x_\Pi))$  a player first measures their share of  $|\psi\rangle$  using the projective measurement for  $\mathcal{S}^{\text{ORAC}}$  for the typed question  $(t_Q, x_Q)$  to obtain an outcome  $a_Q$ . The player then computes an answer, depending on  $t_Q, a_Q$  and  $t_\Pi, x_\Pi$ , as follows:
  - (a) Suppose  $t_Q = v \in \{\text{A}, \text{B}\}$  and  $t_\Pi \in \{\text{POINT}_v, \text{ALINE}_v\}$  (resp.  $\text{DLINE}_v$  instead of  $\text{ALINE}_v$ ). Let  $a'_Q = a_Q$  if  $a_Q$  has length at most  $T$ , and let  $a'_Q$  be the truncation of  $a_Q$  to its first  $T$  symbols otherwise. Let  $\ell_a \leq T$  be the length of  $a'_Q$ , and set

$$a''_Q = \text{enc}_\Gamma(a'_Q, \sqcup^{M/2-\ell_a}).$$

Next, the player computes the low-degree encoding  $g_{a''_Q}$  of  $a''_Q$  using the low-degree encoding described in Section 3.4, in the same way as described in Section 10.4.2. The player then returns the restriction of  $g_{a''_Q}$  to the axis-parallel line (resp. the diagonal line) specified by  $x_\Pi$ .

- (b) If  $t_Q = \text{ORACLE}$ , for  $v \in \{A, B\}$  the player computes questions  $x_v = L^v(x_Q)$ , as in Step 1 of  $\hat{\mathcal{D}}^{\text{ORAC}}$ . The player parses  $a_Q$  as a pair  $(a_A, a_B)$ . Let  $a'_A = a_A$  if  $a_A$  has length at most  $T$ , and let  $a'_A$  be the truncation of  $a_A$  to its first  $T$  symbols otherwise. Let  $\ell_A \leq T$  be the length of  $a'_A$ , and set

$$a''_A = \text{enc}_\Gamma(a'_A, \sqcup^{M/2-\ell_A}).$$

Define  $a''_B$  similarly. The player computes a PCP proof  $\Pi = (g_1, \dots, g_5, c_0, \dots, c_{m'})$  as described in the completeness case of Theorem 10.25 for the tuple  $(\mathcal{D}, n, T, x_A, x_B)$ , where the polynomials  $g_1, g_2$  are low-degree encodings of  $a''_A$  and  $a''_B$ , respectively.

- i. If  $t_\Pi \in \{\text{POINT}_i, \text{ALINE}_i\}$  (resp.  $\text{DLINE}_i$  instead of  $\text{ALINE}_i$ ) for  $i \in \{1, \dots, 5\}$ , the player returns the restriction of  $g_i$  to the axis-parallel line (resp. diagonal line) of  $\mathbb{F}_q^m$  specified by  $x_\Pi$ .
  - ii. If  $t_\Pi \in \{\text{POINT}_6, \text{ALINE}_6\}$  (resp.  $\text{DLINE}_6$  instead of  $\text{ALINE}_6$ ), the player returns the restriction of all the polynomials  $g_1, \dots, g_5, c_0, \dots, c_{m'}$  to the axis-parallel line of  $\mathbb{F}_q^{m'}$  (resp. the diagonal line) specified by  $x_\Pi$ .
- (c) In all other cases the player returns 0.

The strategy  $\mathcal{S}^{\text{AR}}$  is projective and consistent because  $\mathcal{S}^{\text{ORAC}}$  is. To show that it has value 1, we first observe that by definition it satisfies all consistency checks. Moreover, the strategy passes all low-degree tests with certainty because it always returns restrictions of consistent polynomials. Finally, it also passes the game check with probability 1. This follows from the completeness statement of the PCP made in Theorem 10.25 and the fact that, if  $\mathcal{D}$  accepts the input  $(n, x_A, x_B, a_A, a_B)$  in time at most  $T$  then it also accepts  $(n, x_A, x_B, a'_A, a'_B)$  in time at most  $T$ , where  $a'_A$  and  $a'_B$  are obtained from  $a_A$  and  $a_B$  by truncating them to strings of length  $T$  if their lengths exceed  $T$ .

To show that  $\mathcal{S}^{\text{AR}}$  is commuting, note that using the product structure of  $\hat{\mathcal{S}}^{\text{AR}}$  every typed question pair with positive probability consists of a pair of questions  $((t_{Q,A}, x_{Q,A}), (t_{Q,B}, x_{Q,B}))$  with positive probability for  $\hat{\mathcal{S}}^{\text{ORAC}}$ , together with an arbitrary pair  $((t_{\Pi,A}, x_{\Pi,A}), (t_{\Pi,B}, x_{\Pi,B}))$ . Using that  $\mathcal{S}^{\text{ORAC}}$  is commuting and that the additional operations associated with  $((t_{\Pi,A}, x_{\Pi,A}), (t_{\Pi,B}, x_{\Pi,B}))$  amount to classical post-processing it follows that  $\mathcal{S}^{\text{AR}}$  is commuting.

This establishes the existence of a symmetric PCC strategy for  $\hat{\mathcal{V}}_n^{\text{AR}}$  with value 1. By Lemma 6.18 it follows that there exists a symmetric PCC strategy for  $\mathcal{V}_n^{\text{AR}} = \text{Detype}(\hat{\mathcal{V}}_n^{\text{AR}})_n$  with value 1.  $\square$

## 10.7 Soundness of the answer-reduced verifier

*Proof of the soundness part of Theorem 10.27.* We first show the soundness for the typed verifier  $\hat{\mathcal{V}}^{\text{AR}}$ . Soundness for the detyped verifier  $\mathcal{V}^{\text{AR}}$  follows from Lemma 6.18, with a constant-factor loss using that the type set  $\mathcal{T}^{\text{AR}}$  for  $\hat{\mathcal{V}}^{\text{AR}}$  has constant size.

We proceed in two steps. Fix an index  $n \geq 1$  and suppose that  $\text{val}^*(\hat{\mathcal{V}}_n^{\text{AR}}) > 1 - \varepsilon$  for some  $\varepsilon > 0$ . Observe that  $\hat{\mathcal{S}}^{\text{ORAC}}$  and  $\hat{\mathcal{S}}^{\text{PCP}}$  both sample distributions that are invariant under permutation of the two players; therefore, the same holds for  $\hat{\mathcal{S}}^{\text{AR}}$ . Moreover, the decider  $\hat{\mathcal{D}}^{\text{AR}}$  treats both players symmetrically. Therefore, the game played by  $\hat{\mathcal{V}}_n^{\text{AR}}$  is a symmetric game. Applying Lemma 5.7 it follows that there exists a symmetric projective strategy  $\mathcal{S} = (|\psi\rangle, M)$  for  $\hat{\mathcal{V}}_n^{\text{AR}}$  with value greater than  $1 - \varepsilon$ .

We use the following shorthand notation. A pair of questions to the players is  $((t_A, x'_A), (t_B, x'_B))$  where for  $w \in \{A, B\}$ ,  $t_w = (t_{Q,w}, t_{\Pi,w})$  and  $x'_w = (x'_{Q,w}, x'_{\Pi,w})$ . When  $w$  is clear from context we omit it from the subscript. Fixing a  $w$ , whenever  $t_Q = \text{ORACLE}$  we introduce  $x_A = L^A(x_Q)$  and  $x_B = L^B(x'_Q)$  and

often write directly the player's question as  $x_Q = (x_A, x_B)$ . Whenever  $t_Q = v \in \{A, B\}$  we slightly abuse notation and write the question as  $x_Q = (x_v, v)$ , explicitly including the type to clarify which player it points to.

We denote the measurements used by both players in strategy  $\mathcal{S}$  by  $\{M(x_Q)_a^{x_\Pi}\}$ , where for the sake of clarity we have notationally separated the two parts  $x_Q$  and  $x_\Pi$  of the question and omitted explicit mention of the associated types  $t_Q$  and  $t_\Pi$  (we include the type and write  $M(x_Q)_a^{t_\Pi, x_\Pi}$  when it is needed for clarity). First we show that the strategy  $\mathcal{S}$  is close to a strategy  $\mathcal{S}'$  that performs “low-degree” measurements: upon receipt of a typed question  $(t, x) = ((t_Q, t_\Pi), (x_Q, x_\Pi))$  a player first performs a measurement depending on  $x_Q$  to obtain a tuple of low-degree polynomials, and then returns evaluations of those polynomials on the subspaces (either axis-parallel or diagonal lines) specified by  $x_\Pi$ . This step of the argument uses the quantum soundness of the low-degree test performed in Steps 3 and 4 of Figure 14. Next, we “decode” this strategy to produce a strategy  $\mathcal{S}''$  for  $\hat{\mathcal{Y}}^{\text{ORAC}}$  with a high value. This step makes use of the classical soundness of the underlying PCP shown in Section 10.4. The conclusion of the theorem then follows from the soundness of  $\hat{\mathcal{Y}}^{\text{ORAC}}$  (Theorem 9.3). We proceed with the details.

We start by showing a sequence of claims that establish approximations implied by the assumption that  $\mathcal{S}^{\text{AR}}$  succeeds with probability greater than  $1 - \varepsilon$  in the decision procedure implemented by the decider in Figure 14.

**Claim 10.28** (Global consistency check, Step 1). *On average over questions  $(t_A, x_A) = ((t_Q, t_\Pi), (x_Q, x_\Pi))$  sampled from the marginal distribution of  $\mu_{\mathcal{S}^{\text{AR}}}$  on the first player it holds that*

$$M(x_Q)_a^{x_\Pi} \otimes I \simeq_\varepsilon I \otimes M(x_Q)_a^{x_\Pi}. \quad (136)$$

*Proof.* First we observe that the condition  $t_A = t_B$  for the global consistency check, Step 1 in Figure 14, holds with constant probability over the choice of a pair of questions  $(t_A, x_A), (t_B, x_B)$  sampled according to  $\mu_{\mathcal{S}^{\text{AR}}}$ . Thus  $\mathcal{S}$  must succeed in this test with probability  $1 - O(\varepsilon)$ , conditioned on the test being executed: this is because each of  $\hat{\mathcal{S}}^{\text{ORAC}}$  and  $\hat{\mathcal{S}}^{\text{PCP}}$  have a constant probability of returning a pair of questions of the same type.

Moreover, observe that conditioned on  $t_A = t_B$  a pair of questions  $((t_A, x_A), (t_A, x_B)) \sim \mu_{\mathcal{S}^{\text{AR}}}$  is such that  $x_A = x_B = L_{t_A}(z)$ , where  $z$  is the sampler seed and  $L_{t_A}$  the CL function of type  $t_A$  associated with  $\hat{\mathcal{S}}^{\text{AR}}$ . The claim then follows directly from the test and the definition of approximate consistency (Definition 5.15).  $\square$

**Claim 10.29** (Input consistency check, Step 2). *For all  $v \in \{A, B\}$ , on average over question pairs  $(x_A, x_B) \sim \mu_{\mathcal{S}}$  and  $z = (y_1, \dots, y_5, 0, w) \in \mathbb{F}_q^{m'}$  sampled uniformly at random,*

$$M(x_A, x_B)_{\alpha_v}^{\text{POINT}_6, z} \otimes I \simeq_\varepsilon I \otimes M(x_v, v)_{\alpha_v}^{\text{POINT}_v, y_v}, \quad (137)$$

where as in Remark 10.26 we made the identification  $1 \leftrightarrow A$  and  $2 \leftrightarrow B$ . Moreover, an analogous relation holds for operators acting on opposite sides of the tensor product.

*Proof.* For  $w = A$  and fixed  $v \in \{A, B\}$  there is a constant probability that  $t_{Q,w} = \text{ORACLE}$ ,  $t_{Q,\bar{w}} = v$ , and  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_6, \text{POINT}_v)$ . Therefore, the input consistency check in Step 2 is executed with constant probability, and  $\mathcal{S}$  must pass it with probability  $1 - O(\varepsilon)$ , conditioned on the test being executed.

Moreover, conditioned on  $t_{Q,w} = \text{ORACLE}$ ,  $t_{Q,\bar{w}} = v$ , and  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_6, \text{POINT}_v)$ , the distribution of  $(x_{Q,w}, x_{Q,\bar{w}})$  is  $((x_A, x_B), x_v)$  for  $(x_A, x_B) \sim \mu_{\mathcal{S}}$  and the distribution of  $(x_{\Pi,w}, x_{\Pi,\bar{w}})$  is  $(z, y_v)$  for a uniformly random  $z \in \mathbb{F}_q^{m'}$ . Eq. (137) then follows directly from the specification of the test and the definition of approximate consistency. The “moreover” part follows from the case  $w = B$ .  $\square$

**Claim 10.30** (Input low degree test, Step 3). *For each  $v \in \{A, B\}$  and for each  $x$  in the support of the marginal of  $\mu_S$  on player  $v$  there exists a measurement  $\{G_g^{x,v}\} \in \text{PolyMeas}(m, d, q)$  such that the following hold for some  $\delta_1 = O(\delta_{\text{LD}}(O(\varepsilon), q, m, d, 1))$ , where  $\delta_{\text{LD}}$  is defined in Theorem 7.8. For all  $v \in \{A, B\}$ , on average over  $x$  chosen from the marginal of  $\mu_S$  on player  $v$  and  $y_v \in \mathbb{F}_q^m$  sampled uniformly at random,*

$$M(x, v)_\alpha^{\text{POINT}_{v, y_v}} \otimes I \simeq_{\delta_1} I \otimes G_{[\text{eval}_{y_v}(\cdot)=\alpha]}^{x,v}, \quad (138)$$

$$I \otimes M(x, v)_\alpha^{\text{POINT}_{v, y_v}} \simeq_{\delta_1} G_{[\text{eval}_{y_v}(\cdot)=\alpha]}^{x,v} \otimes I, \quad (139)$$

$$G_g^{x,v} \otimes I \simeq_{\delta_1} I \otimes G_g^{x,v}, \quad (140)$$

where we used the notation  $\text{eval}_{y_v}(g) = g(y_v)$  for the evaluation map.

*Proof.* Fix  $v \in \{A, B\}$ . For any  $w \in \{A, B\}$  there is a constant probability that  $t_{Q,w} = t_{Q,\bar{w}} = v$  and  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_v, \text{ALINE}_v)$  (resp.  $\text{DLINE}_v$ ). Therefore, the input low degree test in Step 3 of Figure 14 is executed with constant probability, and  $\mathcal{S}$  must pass it with probability  $1 - O(\varepsilon)$ , conditioned on the test being executed.

Observe that by definition the distribution of  $(x_{\Pi,A}, x_{\Pi,B})$  conditioned on  $t_{Q,w} = t_{Q,\bar{w}} = v$ , uniformly random  $x_Q = (x_v, v)$ , and  $(t_{\Pi,w}, t_{\Pi,\bar{w}}) = (\text{POINT}_v, \text{ALINE}_v)$  (resp.  $\text{DLINE}_v$ ), where  $w \in \{A, B\}$  is uniformly random, is exactly the distribution of questions in the game  $\mathfrak{G}^{\text{LD}}$  described in Section 7.1.1, parametrized by  $\text{ldparams} = (q, m, d, 1)$ .

For every  $v \in \{A, B\}$  and question  $x = L^v(z)$  in the support of the marginal distribution of  $\mu_S$  on player  $v$  let  $\varepsilon_{x,v}$  be the probability that  $\mathcal{S}$  is rejected in Step 3, conditioned on the test being executed and on average over  $w \in \{A, B\}$ . Then  $\mathbb{E}[\varepsilon_{x,v}] = O(\varepsilon)$ , where the expectation is taken over a uniformly random  $v \in \{A, B\}$  and  $x = L^v(z)$  for uniformly random  $z$ .

By definition it follows that the strategy  $\hat{S}^{\text{AR}}$  conditioned on the first part of the players' questions being  $t_{Q,w} = t_{Q,\bar{w}} = v$  and  $x_{Q,A} = x_{Q,B} = x$  is a projective strategy that succeeds with probability  $1 - \varepsilon_{x,v}$  in the low-degree test  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$  executed in Item 3.

We may thus apply Theorem 7.8 to obtain  $\{G_g^{x,v}\} \in \text{PolyMeas}(m, d, q)$  such that (138), (139) and (140) each hold with approximation error  $O(\delta_{\text{LD}}(\varepsilon_{x,v}, q, m, d, 1))$ . Using that for fixed  $q, m, d$  the function  $\varepsilon \mapsto \delta_{\text{LD}}(\varepsilon, q, m, d, 1)$  is concave, the claim follows from Jensen's inequality.  $\square$

**Claim 10.31** (Proof encoding checks, Step 4). *For each  $x_Q = (x_A, x_B)$  in the support of  $\mu_S$  there exist measurements  $\{G_g^{(x_A, x_B), i}\} \in \text{PolyMeas}(m, d, q)$  for each  $i \in \{3, 4, 5\}$  and*

$$\{J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{(x_A, x_B)}\} \in \text{PolyMeas}(m', d, q, m' + 6)$$

such that the following hold for some

$$\delta_2 = O(\delta_{\text{LD}}(O(\varepsilon), q, m, d, 1) + \delta_{\text{LD}}(O(\varepsilon), q, m', d, m' + 6)).$$

First, for all  $i \in \{3, 4, 5\}$ , on average over  $(x_A, x_B) \sim \mu_S$  and  $z = (y_1, \dots, y_5, 0, w)$  of type  $\text{POINT}_6$  sampled uniformly at random,

$$I \otimes M(x_A, x_B)_{\alpha_i}^{\text{POINT}_{i, y_i}} \simeq_\varepsilon M(x_A, x_B)_{\alpha_i}^{\text{POINT}_{6, z}} \otimes I. \quad (141)$$

Second, for all  $i \in \{3, 4, 5\}$  and on average over  $(x_A, x_B) \sim \mu_S$  and  $y_i \in \mathbb{F}_q^m$  sampled uniformly at random,

$$M(x_A, x_B)_{\alpha}^{\text{POINT}_i, y_i} \otimes I \simeq_{\delta_2} I \otimes G_{[\text{eval}_{y_i}(\cdot)=\alpha]}^{(x_A, x_B), i}, \quad (142)$$

$$G_g^{(x_A, x_B), i} \otimes I \simeq_{\delta_2} I \otimes G_g^{(x_A, x_B), i}. \quad (143)$$

Third, for all  $i \in \{1, \dots, 5\}$  and  $j \in \{0, \dots, m'\}$ , on average over  $(x_A, x_B) \sim \mu_S$  and  $z \in \mathbb{F}_q^{m'}$  sampled uniformly at random,

$$M(x_A, x_B)_{\alpha_1, \dots, \alpha_5, \beta_0, \dots, \beta_{m'}}^{\text{POINT}_6, z} \otimes I \simeq_{\delta_2} I \otimes J_{[\text{eval}_z(\cdot)=(\alpha_1, \dots, \alpha_5, \beta_0, \dots, \beta_{m'})]}^{(x_A, x_B)}, \quad (144)$$

$$J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{(x_A, x_B)} \otimes I \simeq_{\delta_2} I \otimes J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{(x_A, x_B)}. \quad (145)$$

Moreover, analogous equations to (141), (142) and (144) hold with operators acting on opposite sides of the tensor product. Here, recall the definition of  $\text{PolyMeas}(\cdot, \cdot, \cdot)$  from Definition 7.7.

*Proof.* The proof of the first item is similar to the proof of Claim 10.29, and we omit it.

The proof of the second and third items is similar to the proof of Claim 10.30, and we include more details. Fix an  $i \in \{3, 4, 5\}$ . For any  $w \in \{A, B\}$  there is a constant probability that  $t_{Q, w} = t_{Q, \bar{w}} = \text{ORACLE}$  and  $(t_{\Pi, w}, t_{\Pi, \bar{w}}) = (\text{POINT}_i, \text{ALINE}_i)$  (or  $\text{DLINE}_i$  instead of  $\text{ALINE}_i$ ), in which case the individual low-degree test in Step 4b is executed. Therefore,  $\mathcal{S}$  must succeed in that part of the test with probability  $1 - O(\varepsilon)$  conditioned on the test being executed.

Furthermore, for fixed  $i \in \{3, 4, 5\}$  and uniformly random  $w \in \{A, B\}$ , conditioned on the test being executed for that  $i$  and  $w$  the distribution of  $(x_{\Pi, A}, x_{\Pi, B})$  is exactly the distribution of questions in the game  $\mathfrak{G}^{\text{LD}}$  described in Section 7.1.1, parametrized by  $\text{ldparams} = (q, m, d, 1)$ .

For every  $i \in \{3, 4, 5\}$  and  $x = (x_A, x_B)$  in the support of  $\mu_S$  let  $\varepsilon_{x, i}$  be the probability that  $\mathcal{S}$  is rejected in Step 4b, conditioned on the test being executed for that  $i$  and on average over  $w \in \{A, B\}$ . Then for each  $i$ ,  $\mathbb{E}[\varepsilon_{x, i}] = O(\varepsilon)$ , where the expectation is taken over a uniformly random  $x \sim \mu_S$ .

By definition of the individual low-degree test it follows from Theorem 7.8 that for every  $x = (x_A, x_B)$  in the support of  $\mu_S$  and  $i \in \{3, 4, 5\}$  there is a measurement  $\{G_g^{(x_A, x_B), i}\} \in \text{PolyMeas}(m, d, q)$  such that on average over  $y_i \in \mathbb{F}_q^{m'}$  of type  $\text{POINT}_i$  sampled uniformly at random, Eq. (142) and (143) both hold with approximation error  $O(\delta_{\text{LD}}(\varepsilon_{x, i}, q, m, d, 1))$ . Eq. (142) and (143) follow using the concavity of  $\delta_{\text{LD}}$  as a function of  $\varepsilon$ .

Finally we consider the simultaneous low-degree test, Step 4c. Here as well, using that there is a constant probability that  $t_{Q, w} = t_{Q, \bar{w}} = \text{ORACLE}$  and  $(t_{\Pi, w}, t_{\Pi, \bar{w}}) = (\text{POINT}_6, \text{ALINE}_6)$  (resp.  $\text{DLINE}_6$ ) it follows that  $\mathcal{S}^{\text{AR}}$  must succeed in that part of the test with probability  $1 - O(\varepsilon)$ . Using a similar argument as before it follows from Theorem 7.8 (this time for parameters  $(q, m', d, m' + 6)$ ) that for every  $(x_A, x_B)$  there is a family of measurements  $\{J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{(x_A, x_B)}\} \in \text{PolyMeas}(m', d, q, m' + 6)$  such that on average over  $z \in \mathbb{F}_q^{m'}$  sampled uniformly at random, Eq. (144) and (145) both hold with approximation error  $O(\delta_{\text{LD}}(O(\varepsilon), q, m', d, m' + 6))$ .  $\square$

The families of measurements  $\{G_g^{x_Q, i}\}$  and  $\{J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{x_Q}\}$ , for  $x_Q$  in the support of  $\mu_S$  and  $i \in \{1, \dots, 5\}$ , whose existence follows from Claim 10.30 and Claim 10.31 have outcomes that are low (individual) degree polynomials: for the first family, individual degree  $d$  polynomials  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$ , and for the second, tuples of individual degree  $d$  polynomials  $f_i, c_j : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$ . Recall that  $m' = 5m + 5 + s$  and



that an element  $z \in \mathbb{F}_q^{m'}$  is written as a triple  $(y, o, w)$  with  $x = (y_1, \dots, y_5) \in \mathbb{F}_q^{5m}$ ,  $o \in \mathbb{F}_q^5$  and  $w \in \mathbb{F}_q^s$ . The following claim, whose proof is based on Lemma 5.26, shows that we can reduce to a situation where the polynomials  $f_1, \dots, f_5$  returned by  $J$  are such that for each  $i \in \{1, \dots, 5\}$ ,  $f_i$  only depends on the  $y_i$ , and not on the entire variable  $z$ .

**Claim 10.32.** *For all  $(x_A, x_B)$  in the support of  $\mu_S$  and individual degree  $d$  polynomials  $g_1, \dots, g_5 : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  and  $c_0, \dots, c_{m'} : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$  define*

$$\Lambda_{g_1, \dots, g_5, c_0, \dots, c_{m'}}^{x_A, x_B} = G_{g_1}^{x_A, 1} G_{g_2}^{x_B, 2} G_{g_3}^{(x_A, x_B), 3} G_{g_4}^{(x_A, x_B), 4} G_{g_5}^{(x_A, x_B), 5} J_{c_0, \dots, c_{m'}}^{(x_A, x_B)} G_{g_5}^{(x_A, x_B), 5} G_{g_4}^{(x_A, x_B), 4} G_{g_3}^{(x_A, x_B), 3} G_{g_2}^{x_B, 2} G_{g_1}^{x_A, 1}, \quad (146)$$

where the outcomes  $(f_1, \dots, f_5)$  of the  $J$  operator in the middle have been marginalized over. Then there is a

$$\delta_3 = O\left(\left(\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6)^{1/2} + \frac{m'd}{q}\right)^{1/2}\right)$$

such that

$$\delta_3 \geq \max\{\varepsilon, \delta_1, \delta_2\}$$

and on average over  $(x_A, x_B) \sim \mu_S$  and  $z \in \mathbb{F}_q^{m'}$  sampled uniformly at random,

$$\Lambda_{[\text{eval}_z(\cdot)=(\alpha, \beta)]}^{x_A, x_B} \otimes I \simeq_{\delta_3} I \otimes J_{[\text{eval}_z(\cdot)=(\alpha, \beta)]}^{x_A, x_B}. \quad (147)$$

Moreover, a similar equation holds with the operators acting on opposite sides of the tensor product.

*Proof.* We apply Lemma 5.26 with the following setting of parameters. The number of sets of functions  $k$  is set to 6. The question set  $\mathcal{X}$  is set to the support of  $\mu_S$ , and the distribution  $\mu$  on it is the distribution  $\mu_S$ . The sets  $\mathcal{G}_i$  for  $i \in \{1, \dots, 5\}$  consist of individual degree  $d$  polynomials over  $\mathbb{F}_q^{m'}$  that depend only on the  $i$ -th block of  $m$  variables. The set  $\mathcal{G}_6$  consists of  $(m' + 1)$ -tuples of individual degree  $d$  polynomials over  $\mathbb{F}_q^{m'}$ .

We first verify the assumption on the sets of functions. Since all polynomials have individual degree at most  $d$  (and therefore total degree at most  $m'd$ ), by Lemma 3.20 the parameter  $\varepsilon$  in Lemma 5.26 can be set to  $m'd/q$ .

The family of measurements  $\{A_{g_1, \dots, g_6}^x\}$  in Lemma 5.26 is the family of measurements  $\{J_{f_1, \dots, f_5, c_0, \dots, c_{m'}}^{(x_A, x_B)}\}$  here, where we set  $g_i = f_i$  for  $i \in \{1, \dots, 5\}$  and  $g_6 = (c_0, \dots, c_{m'})$ . The measurements  $\{G_g^{i,x}\}$  in Lemma 5.26 are  $\{G_g^{x_A, i}\}$  for  $i \in \{1, 2\}$ ,  $\{G_g^{(x_A, x_B), i}\}$  for  $i \in \{3, 4, 5\}$ , and  $\{J_g^{(x_A, x_B)}\}$  for  $i = 6$ . To ensure that all polynomials are defined over the same range, we treat  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  that is an outcome of some  $\{G_g^{i,x}\}$  as a polynomial  $g' : \mathbb{F}_q^{m'} \rightarrow \mathbb{F}_q$ , where the role of the  $m$  variables of  $g$  is taken by the  $i$ -th block of  $m$  variables of  $g'$ .

We verify that assumption (40) in the lemma holds with an error  $\delta = O((\max\{\varepsilon, \delta_1, \delta_2\})^{1/2})$ . For  $i \in \{1, 2\}$  the assumption follows by combining (137) and (139) with (144) and Fact 5.24. For  $i \in \{3, 4, 5\}$  we use (141) instead of (137) and (142) instead of (139). Finally, for  $i = 6$  we use (145) and Fact 5.24. In these derivations, we use Fact 5.23, the triangle inequality for “ $\simeq$ ”.

The conclusion follows from Lemma 5.26, with an error  $\delta_3 = O((\delta + m'd/q)^{1/2})$ . Observing that  $\varepsilon, \delta_1, \delta_2 = O(\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6))$ , as can be verified from the definition of  $\delta_{\text{LD}}$  given in Theorem 7.8, and therefore  $\delta = O(\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6)^{1/2})$ , the claimed bound

$$\delta_3 = O\left(\left(\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6)^{1/2} + \frac{m'd}{q}\right)^{1/2}\right)$$

follows. □

At this point, we have constructed measurements  $G$  and  $\Lambda$  that return low-degree polynomials in a similar way as is expected from the honest strategy in  $\hat{\mathcal{V}}_n^{\text{AR}}$ , as described in the proof of Theorem 10.27. These measurements can be used to specify a new strategy  $\mathcal{S}'$  for the game  $\hat{\mathcal{V}}_n^{\text{AR}}$  as follows. The shared state remains the state  $|\psi\rangle$  used in  $\mathcal{S}$ . For  $w \in \{A, B\}$ , upon reception of a question  $(t_w, x_w)$  player  $w$  performs the following. If  $t_w = (t_{Q,w}, t_{\Gamma,w})$  is such that  $t_{Q,w} = \text{ORACLE}$ , the player measures their share of  $|\psi\rangle$  using the measurement  $\Lambda^{x_{Q,w}}$  defined in Claim 10.32 to obtain a tuple  $(g_1, \dots, g_5, c_0, \dots, c_{m'})$  of polynomials. The player then answers exactly as in the strategy described in the ‘‘completeness’’ part of the proof of Theorem 10.27. Similarly, if  $t_{Q,w} = v \in \{A, B\}$  the player first measures their share of  $|\psi\rangle$  using the measurement  $G^{x_{Q,w},v}$  from Lemma 10.30 to obtain a polynomial  $g$  as outcome; the player then answers according to the same honest strategy.

**Lemma 10.33.** *The strategy  $\mathcal{S}'$  succeeds with probability  $1 - O(\delta_3^{1/2})$  in the game  $\hat{\mathcal{V}}_n^{\text{AR}}$ .*

*Proof.* First we establish useful consistency relations. By combining Equation (147) and Equation (144) and applying Fact 5.24 we obtain that for all  $i \in \{1, \dots, 5\}$ , on average over  $(x_A, x_B) \sim \mu_S$  and  $z \in \mathbb{F}_q^{m'}$  sampled uniformly at random,

$$M(x_A, x_B)_{\alpha_i}^{\text{POINT}_{6,z}} \otimes I \simeq_{\delta_3} I \otimes \Lambda_{[\text{eval}_z(\cdot)_i = \alpha_i]}^{(x_A, x_B)}, \quad (148)$$

and a similar equation holds with the operators acting on opposite sides of the tensor product. Next, combining (148) together with Equation (137) and Equation (138) it follows that for each  $v \in \{A, B\}$ , on average over  $(x_A, x_B) \sim \mu_S$  and  $z = (y_1, \dots, y_5, 0, w) \in \mathbb{F}_q^{m'}$  sampled uniformly at random,

$$G_{[\text{eval}_{y_v}(\cdot) = \alpha_v]}^{x_v, v} \otimes I \simeq_{\delta_3} I \otimes \Lambda_{[\text{eval}_z(\cdot)_v = \alpha_v]}^{(x_A, x_B)}. \quad (149)$$

From the Schwartz-Zippel lemma (Lemma 3.20) it follows that the probability that any two distinct individual degree  $d$  polynomials  $g_v$  (an outcome of  $G^{x_v, v}$ ) and  $g'_v$  (an outcome of  $\Lambda^{(x_A, x_B)}$ ) agree at a uniformly random point  $y_v \in \mathbb{F}_q^m$  is at most  $m'd/q$ . It thus follows from (149) that for all  $v \in \{A, B\}$ , on average over  $(x_A, x_B) \sim \mu_S$  and  $y_v \in \mathbb{F}_q^M$  sampled uniformly at random,

$$G_{g_v}^{x_v, v} \otimes I \simeq_{\delta_3 + m'd/q} I \otimes \Lambda_{g_v}^{(x_A, x_B)}. \quad (150)$$

We now show that  $\mathcal{S}'$  is accepted by  $\hat{\mathcal{V}}_n^{\text{AR}}$  with high probability. We bound the probability of succeeding in each subtest.

First note that the strategy is accepted in Step 1 of Figure 14. For the  $G$  measurements, consistency follows from (140). For the  $\Lambda$  measurements, note first that by (147) and (145) it follows that on average over  $(x_A, x_B) \sim \mu_S$ ,

$$\Lambda_{[\text{eval}_z(\cdot) = (\alpha, \beta)]}^{x_A, x_B} \otimes I \simeq_{\delta_3} I \otimes \Lambda_{[\text{eval}_z(\cdot) = (\alpha, \beta)]}^{x_A, x_B}. \quad (151)$$

Using that all outcomes of  $\Lambda^{x_A, x_B}$  are individual degree  $d$  polynomials and the Schwartz-Zippel lemma (Lemma 3.20) it follows that whenever a measurement of  $\Lambda^{x_A, x_B} \otimes \Lambda^{x_A, x_B}$  returns distinct outcomes, the outcomes take a different value at a uniformly random  $z$  with probability at least  $1 - m'd/q$ . It then follows from (151) and the fact that  $\delta_3 \geq m'd/q$  by definition that the strategy  $\mathcal{S}'$  is accepted in Step 1 with probability  $1 - O(\delta_3)$ .

Next, the strategy is also accepted in the consistency check performed in item 2 due to (150), and the consistency check in item 4(a) for the same reasons as for item 1. Finally, for the low-degree tests performed in item 3 and items 4(b) and 4(c), the strategy succeeds due to consistency and the fact that, as long as both players obtain the same polynomial outcomes, they pass the low-degree tests with probability 1.

It remains to analyze the strategy's success probability in Step 5, the game check. Recall that this check is nontrivial only when  $t_{Q,w} = \text{ORACLE}$  and  $t_{\Pi,w} = \text{POINT}_6$ . By assumption the original strategy  $\mathcal{S}$  succeeds with probability  $1 - O(\varepsilon)$  in the game check. Thus, it will suffice to show that the answer distribution from  $\mathcal{S}'$  is close to the answer distribution from  $\mathcal{S}$  whenever  $t_{Q,w} = \text{ORACLE}$  and  $t_{\Pi,w} = \text{POINT}_6$  for either or both players  $w \in \{A, B\}$ . For these question types, the measurement operator used by  $\mathcal{S}$  is  $M(x_A, x_B)_{(\alpha, \beta)}^{\text{POINT}_{6,z}}$ , and the operator used by  $\mathcal{S}'$  is  $\Lambda_{[\text{eval}_z(\cdot) = (\alpha, \beta)]}^{(x_A, x_B)}$ . By converting Equation (148) and Equation (151) to  $\approx$  bounds using Fact 5.18, and then applying the triangle inequality (Fact 5.22) we obtain that on average over  $(x_A, x_B) \sim \mu_S$ ,

$$\Lambda_{[\text{eval}_z(\cdot) = (\alpha, \beta)]}^{x_A, x_B} \otimes I \approx_{\delta_3} M(x_A, x_B)_{(\alpha, \beta)}^{\text{POINT}_{6,z}} \otimes I_B.$$

Thus, restricting to the game check,  $\mathcal{S}$  and  $\mathcal{S}'$  are  $\delta_3$ -close in the sense of Definition 5.17. They both use the same state, and moreover,  $\mathcal{S}$  is projective. Therefore, by Lemma 5.28, this implies that  $\mathcal{S}'$  must succeed in the game check with probability  $1 - \varepsilon - O(\delta_3^{1/2}) = 1 - O(\delta_3^{1/2})$ . Thus, we have shown that  $\mathcal{S}'$  succeeds in all subtests of the game with probability  $1 - O(\delta_3^{1/2})$ , and thus the lemma follows.  $\square$

We now complete the proof by a reduction to the game  $\mathcal{V}_n^{\text{ORAC}}$ : from the strategy  $\mathcal{S}'$  we construct a symmetric strategy  $\mathcal{S}'' = (|\psi\rangle, A)$  for  $\hat{\mathcal{V}}^{\text{ORAC}}$  by “decoding” the low-degree measurements  $G$  and  $\Lambda$ . The state  $|\psi\rangle$  in  $\mathcal{S}''$  is identical to the state used in  $\mathcal{S}'$  (which is identical to the state used in  $\mathcal{S}$ ). To begin, we define a decoding map  $\Delta(\cdot)$ , which takes in a polynomial  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  and outputs a string in  $\{0, 1\}^*$ . This map is computed as follows:

- First, compute  $a = \text{Dec}(g) \in \{0, 1\}^M$ , where  $\text{Dec}$  is the (boolean) decoding of the low-degree code defined in Section 3.4.
- If there exists an  $a_{\text{prefix}} \in \{0, 1\}^*$  of length  $\ell_a \leq T$  such that  $a = \text{enc}_\Gamma(a_{\text{prefix}}, \sqcup^{M/2 - \ell_a})$ , then  $\Delta(g) = a_{\text{prefix}}$ . Otherwise,  $\Delta(g)$  is allowed to be arbitrary.

We can now define the “decoded” measurements  $\{A^{x_{v,v}}\}$  and  $\{A^{x_A, x_B}\}$  as follows:

$$A_a^{x_{v,v}} = G_{[\Delta(\cdot) = a]}^{x_{v,v}}, \quad A_{a_A, a_B}^{x_A, x_B} = \Lambda_{[\Delta(\cdot)_{A,B} = (a_A, a_B)]}^{x_A, x_B}. \quad (152)$$

**Lemma 10.34.** *The strategy  $\mathcal{S}''$  succeeds with probability  $1 - O(\delta_3^{1/2})$  in the game  $\hat{\mathcal{V}}_n^{\text{ORAC}}$ .*

*Proof.* We consider the different subtests executed by  $\hat{\mathcal{D}}^{\text{ORAC}}$  (see Figure 12). We start with Step 2, the consistency checks. Success in the first check, Step 2a, follows from the success of  $\mathcal{S}'$  in the global consistency check, Step 1 of  $\hat{\mathcal{D}}^{\text{AR}}$ , the definition (152), and the fact that conditioned on  $t_{Q,A} = t_{Q,B} = \text{ORACLE}$ , the distribution of  $(x_{Q,A}, x_{Q,B})$  in  $\hat{\mathcal{V}}_n^{\text{AR}}$  is the same as the distribution of  $(x_A, x_B)$  in  $\mathcal{V}_n^{\text{ORAC}}$ , conditioned on  $t_A = t_B = \text{ORACLE}$ . Similarly, success in the second check, Step 2b, follows from success of  $\mathcal{S}'$  in the input consistency check, Step 2 of  $\hat{\mathcal{D}}^{\text{AR}}$ .

Next we consider the game check of  $\hat{\mathcal{D}}^{\text{ORAC}}$ , Step 1. To analyze the success probability of  $\mathcal{S}''$  we use that  $\mathcal{S}'$  succeeds in the game check of  $\hat{\mathcal{D}}^{\text{AR}}$ , Step 5, and the soundness of the PCP, as shown in Theorem 10.25. Let  $p_{\text{sound}}$  be as in Theorem 10.25.

Let  $w \in \{A, B\}$ ,  $(x_A, x_B)$  be in the support of  $\mu_S$ , and  $\Pi = (g_1, \dots, g_5, c_0, \dots, c_{m'})$  an outcome of  $\Lambda^{x_A, x_B}$  such that conditioned on that outcome being obtained by player  $w$  in the game check of  $\hat{\mathcal{D}}^{\text{AR}}$ ,  $\mathcal{M}_{\text{AR}}$  accepts the pair of inputs  $(\mathcal{D}, n, T, Q, \gamma, x_A, x_B)$  and  $(z, a_w)$  with probability at least  $p_{\text{sound}}$  over the choice of a uniformly random  $z \in \mathbb{F}_q^{m'}$  and  $a_w = \text{eval}_z(\Pi)$ .

For any such  $\Pi$ , the soundness statement of Theorem 10.25 states that there exist  $a_A, a_B \in \{0, 1\}^*$  such that  $\mathcal{D}(n, x_A, x_B, a_A, a_B) = 1$  and furthermore  $a_v = \Delta(g_v)$  for  $v \in \{A, B\}$ . It follows that for any proof  $\Pi$  returned by  $\{\Lambda_{\Pi}^{x_A, x_B}\}$  which is accepted with probability greater than  $p_{\text{sound}}$  in the game check of  $\hat{\mathcal{D}}^{\text{AR}}$  it holds that  $\mathcal{D}(n, x_A, x_B, \Delta(g_A), \Delta(g_B)) = 1$ . Using this observation we evaluate the probability  $q_g''$  that the strategy  $\mathcal{S}''$  succeeds in the game check of  $\hat{\mathcal{V}}^{\text{ORAC}}$ . Let  $q_g'$  be the probability that  $\mathcal{S}'$  succeeds in the game check of  $\hat{\mathcal{D}}^{\text{AR}}$ .

$$\begin{aligned}
q_g'' &= \mathbb{E}_{(x_A, x_B) \sim \mu_S} \sum_{a_A, a_B: \mathcal{D}(n, x_A, x_B, a_A, a_B) = 1} \langle \psi | A_{a_A, a_B}^{(x_A, x_B)} \otimes I | \psi \rangle \\
&= \mathbb{E}_{(x_A, x_B) \sim \mu_S} \sum_{\Pi: \mathcal{D}(n, x_A, x_B, \Delta(g_1), \Delta(g_2)) = 1} \langle \psi | \Lambda_{\Pi}^{(x_A, x_B)} \otimes I | \psi \rangle \\
&\geq \mathbb{E}_{(x_A, x_B) \sim \mu_S} \sum_{\Pi: \mathcal{D}(n, x_A, x_B, \Delta(g_1), \Delta(g_2)) = 1} \langle \psi | \Lambda_{\Pi}^{(x_A, x_B)} \otimes I | \psi \rangle \cdot \Pr_{z \sim \mathbb{F}_q^{m'}} [\mathcal{M}_{\text{AR}}(z, \text{eval}_z(\Pi)) = 1] \\
&= q_g' - \mathbb{E}_{(x_A, x_B) \sim \mu_S} \sum_{\Pi: \mathcal{D}(n, x_A, x_B, \Delta(g_1), \Delta(g_2)) = 0} \langle \psi | \Lambda_{\Pi}^{(x_A, x_B)} \otimes I | \psi \rangle \cdot \Pr_{z \sim \mathbb{F}_q^{m'}} [\mathcal{M}_{\text{AR}}(z, \text{eval}_z(\Pi)) = 1] \\
&\geq q_g' - (1 - q_g'') \cdot p_{\text{sound}}.
\end{aligned}$$

Rearranging terms,

$$q_g'' \geq \frac{q_g' - p_{\text{sound}}}{1 - p_{\text{sound}}} = 1 - \frac{1 - q_g'}{1 - p_{\text{sound}}}. \quad (153)$$

Altogether, using Lemma 10.33 we have shown that  $\mathcal{S}''$  is accepted in each subtest performed by  $\hat{\mathcal{D}}^{\text{ORAC}}$  with probability at least  $1 - O(\delta_3^{1/2})$ . Since every subtest occurs with constant probability, the lemma follows.  $\square$

To conclude the proof of the soundness, we appeal to the soundness statement for  $\hat{\mathcal{V}}^{\text{ORAC}}$ , given in Theorem 9.3. Now we establish the form of the error function  $\delta$  as stated in the theorem.

Let  $a', b'$  be the universal constants from Theorem 7.8 such that

$$\delta_{\text{LD}}(\varepsilon, q, m, d, t) = a' (dmt)^{a'} (\varepsilon^{b'} + q^{-b'} + 2^{-b' md}).$$

By our choice of PCP parameters in Definition 10.22, we have that  $m \leq s$ , so therefore we can upper bound the product  $dm'(m' + 6)$  by  $c\gamma s^3$  where  $c > 1$  is a universal constant. Furthermore, we see from Definition 10.22 that  $2^{dm} \geq q \geq s^{(\gamma b' + 3a')/b'}$ , so we have

$$\begin{aligned}
\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6) &\leq a' (c\gamma)^{a'} (s^{3a'} \cdot \varepsilon^{b'} + 2 \cdot s^{3a'} \cdot q^{-b'}) \\
&\leq 2a' (c\gamma)^{a'} (s^{3a'} \cdot \varepsilon^{b'} + s^{3a'} \cdot s^{-(\gamma b' + 3a')}) \\
&\leq a'' \gamma^{a''} (s^{a''} \cdot \varepsilon^{b''} + s^{-\gamma b''})
\end{aligned} \quad (154)$$

where we set  $a'' = \max\{2a'c^{a'}, 3a'\}$  and  $b'' = b'$ . From Proposition 10.20 and Eq. (134) we have that  $s$  is at most  $\text{poly}((\lambda n)^\mu, \sigma)$ , so there exists a universal constant  $C > 0$  such that for all  $n \geq 2$ ,

$$s \leq C(\lambda \cdot \sigma \cdot n)^{C\mu}.$$

Plugging this bound into Equation (154) and using the choice of  $q, d, m'$  from Definition 10.22, we get that there exist universal constants  $a > 1$  and  $0 < b < 1$  such that

$$\begin{aligned}\delta_3^{1/2} &= O\left(\left(\delta_{\text{LD}}(\varepsilon, q, m', d, m' + 6)^{1/2} + \frac{m'd}{q}\right)^{1/4}\right) \\ &\leq a\gamma^a((\lambda \cdot \sigma \cdot n)^{\mu a} \cdot \varepsilon^b + (\lambda \cdot \sigma \cdot n)^{-\mu b \gamma}).\end{aligned}$$

To show the entanglement bound, we observe that the strategy  $\mathcal{S}''$  for  $\hat{\mathcal{V}}^{\text{ORAC}}$  constructed above uses the same entangled state  $|\psi\rangle$  as the strategy  $\mathcal{S}$  for  $\hat{\mathcal{V}}^{\text{AR}}$  that we started with; the claimed bound follows.  $\square$

## 11 Parallel Repetition

In each of the transformations on verifiers presented so far (introspection, oracularization, and answer reduction), the *soundness gap* of the resulting verifier is slightly degraded: while the completeness property (i.e. the property of having a PCC strategy with success probability 1) is preserved, if the starting game  $\mathcal{V}_n$  has value at most  $1 - \varepsilon$ , the resulting game  $\mathcal{V}'_n$  has value at most  $1 - \delta(\varepsilon, n)$  for some function  $\delta$ . In order to apply the compression procedure recursively we need a way to restore the soundness gap after a sequence of transformations. We accomplish this using (a modification of) the technique of *parallel repetition*. This is a transformation of a two-player game  $\mathfrak{G}$  into another two-player game  $\mathfrak{G}^k$  where the verifier samples  $k$  independent pairs  $(x_1, y_1), \dots, (x_k, y_k)$  from the question distribution  $\mu$  of  $\mathfrak{G}$ , sends the tuple  $(x_1, \dots, x_k)$  to the first player to obtain an answer tuple  $(a_1, \dots, a_k)$ , and sends the tuple  $(y_1, \dots, y_k)$  to the second player to obtain an answer tuple  $(b_1, \dots, b_k)$ . The players win if and only if  $D(x_i, y_i, a_i, b_i) = 1$  for all  $i$ .

Intuitively, if the value of  $\mathfrak{G}$  is  $v < 1$ , then one would expect the value of  $\mathfrak{G}^k$  to decay exponentially with the number of repetitions  $k$ . It is not true in general that the value of  $\mathfrak{G}^k$  is  $v^k$ , but exponential decay bounds on the (tensor product) value of parallel-repeated games are known for specific classes of games [JPY14, DSV15, BVY17]. In particular, it was shown in [BVY17] that the class of *anchored games* satisfies exponential-decay parallel repetition, and furthermore every game can be efficiently transformed into an equivalent anchored game. Put together, this gives a general soundness amplification procedure called “anchored parallel repetition,” which we use in our compression procedure to reset the soundness gap to a fixed constant.

We describe the “anchoring” transformation of a game  $\mathfrak{G}$  in more detail. The anchoring of  $\mathfrak{G}$  is another game  $\mathfrak{G}_\perp$  where the verifier behaves as follows: it samples a question pair  $(x, y)$  from the question distribution  $\mu$  of  $\mathfrak{G}$ , and then for each player independently changes their question to a special symbol  $\perp$  (not part of the original game  $\mathfrak{G}$ ) with probability  $1/2$  to obtain new questions  $(x', y')$ . The first player receives question  $x'$  and responds with answer  $a$ ; the second player receives question  $y'$  and responds with answer  $b$ . If neither player receives the question  $\perp$ , then they win if and only if  $D(x', y', a, b) = 1$  where  $D$  is the decision predicate of  $\mathfrak{G}$ . Otherwise, the players win if and only if the players receiving the question  $\perp$  output a specific, fixed answer (e.g. 0).

It is easy to see that  $\text{val}^*(\mathfrak{G}_\perp) = \frac{3}{4} + \frac{1}{4}\text{val}^*(\mathfrak{G})$ . In particular, we have that  $\text{val}^*(\mathfrak{G}_\perp) = 1$  if and only if  $\text{val}^*(\mathfrak{G}) = 1$ . Bavarian, Vidick and Yuen proved the following bound on the entanglement requirements of  $\mathfrak{G}_\perp^k$ , which is the  $k$ -fold repetition of  $\mathfrak{G}_\perp$ .

**Theorem 11.1** (Parallel repetition of anchored games, Theorem 6.1 of [BVY17]). *There exists a universal constant  $c > 0$  such that for all two-player games  $\mathfrak{G}$  and for all positive integers  $k$ , for all  $0 < \varepsilon \leq 1$ , for all  $p$  satisfying*

$$p > \frac{4}{\varepsilon} \exp\left(-\frac{c \varepsilon^{17} k}{s}\right)$$

where  $s$  denotes the bit-length of the players’ answers in the game  $\mathfrak{G}$ , we have

$$\mathcal{E}(\mathfrak{G}_\perp^k, p) \geq \mathcal{E}(\mathfrak{G}, 1 - \varepsilon).$$

We note that a corollary of Theorem 11.1 is that if  $\text{val}^*(\mathfrak{G}) < 1 - \varepsilon$ , then  $\text{val}^*(\mathfrak{G}_\perp^k) \leq p$  for  $p = \frac{4}{\varepsilon} \exp\left(-\frac{c \varepsilon^{17} k}{s}\right)$ . This is because if  $\text{val}^*(\mathfrak{G}_\perp^k)$  were larger than  $p$ , there would be a finite upper bound on  $\mathcal{E}(\mathfrak{G}_\perp^k, p)$ , but on the other hand  $\mathcal{E}(\mathfrak{G}, 1 - \varepsilon)$  is by definition infinite, a contradiction.

In this section, we present the anchoring and parallel repetition transformations on normal form verifiers; starting with a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ , we first apply the anchoring transformation to obtain a

normal form verifier  $\mathcal{V}^{\text{ANCH}} = (\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$ , then apply parallel repetition to obtain a normal form verifier  $\mathcal{V}^{\text{REP}} = (\mathcal{S}^{\text{REP}}, \mathcal{D}^{\text{REP}})$ . The main theorem of the section, Theorem 11.4, establishes the properties of the verifier  $\mathcal{V}^{\text{REP}}$ .

**Remark 11.2.** *The parallel repetition theorems of [DSV15, Yue16] are also applicable for the purpose of amplification of the soundness gap, but are not sufficient for us. This is because the (proofs of the) parallel repetition theorems of [DSV15, Yue16] only imply that  $\mathcal{E}(\mathfrak{G}^k, v) \geq \log \mathcal{E}(\mathfrak{G}, 1 - \epsilon)$ , which is not sufficient for the recursive compression application; we need a larger entanglement lower bound on  $\mathfrak{G}^k$ . The weaker entanglement bound is due to the use of the “quantum correlated sampling lemma” of [DSV15]; it is an open question whether there is an improved analysis that obtains a better entanglement lower bound.*

## 11.1 The anchoring transformation

We present the anchoring transformation on normal form verifiers  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ , which produces another normal form verifier  $\mathcal{V}^{\text{ANCH}} = (\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$ . We first define a typed verifier  $\hat{\mathcal{V}}^{\text{ANCH}} = (\hat{\mathcal{S}}^{\text{ANCH}}, \hat{\mathcal{D}}^{\text{ANCH}})$ , and then detype  $\hat{\mathcal{V}}^{\text{ANCH}}$  using Lemma 6.18 to obtain  $\mathcal{V}^{\text{ANCH}}$ .

Define the type set  $\mathcal{T}^{\text{ANCH}} = \{\text{GAME}, \text{ANCHOR}\}$  and type graph  $G^{\text{ANCH}}$  to be the complete graph over  $\mathcal{T}^{\text{ANCH}}$  along with self-loops at each vertex. The Turing machine  $\hat{\mathcal{S}}^{\text{ANCH}}$  is defined as follows:

1. On input  $(n, \text{DIMENSION})$ , it returns  $\mathcal{S}(n, \text{DIMENSION})$ .
2. On input  $(n, w, \text{MARGINAL}, j, z, t)$ , it returns  $\mathcal{S}(n, w, \text{MARGINAL}, j)$  if  $t = \text{GAME}$ , and otherwise returns the binary representation of the zero vector in  $\mathbb{F}_2^s$  where  $s = \mathcal{S}(n, \text{DIMENSION})$ .
3. On input  $(n, w, \text{LINEAR}, j, u, y, t)$ , it returns  $\mathcal{S}(n, w, \text{LINEAR}, j, u, y)$  if  $t = \text{GAME}$ , and otherwise returns the binary representation of the zero vector in  $\mathbb{F}_2^s$  where  $s = \mathcal{S}(n, \text{DIMENSION})$ .
4. On input  $(n, w, \text{FACTOR}, j, u, t)$ , it returns  $\mathcal{S}(n, w, \text{FACTOR}, j, u)$  if  $t = \text{GAME}$ , and otherwise returns the binary representation of the zero vector in  $\mathbb{F}_2^s$  where  $s = \mathcal{S}(n, \text{DIMENSION})$ .

Define the Turing machine  $\hat{\mathcal{D}}^{\text{ANCH}}$  that, on input  $(n, t_A, x_A, t_B, x_B, a_A, a_B)$ , if either  $t_A$  or  $t_B$  is equal to the type ANCHOR, then the decider accepts as long as the players receiving the ANCHOR-type answer with 0 (a player who receives the GAME type can respond with any answer). Otherwise, it accepts only if  $\mathcal{D}(n, x_A, x_B, a_A, a_B)$  accepts. Finally, define the Turing machines  $(\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$  to be the result of the detyping transformation  $\text{Detype}(\hat{\mathcal{V}}^{\text{ANCH}})$  from Lemma 6.18 on the pair  $\hat{\mathcal{V}}^{\text{ANCH}} = (\hat{\mathcal{S}}^{\text{ANCH}}, \hat{\mathcal{D}}^{\text{ANCH}})$ .

Note that this transformation is well-defined for *any* pair of Turing machines  $(\mathcal{S}, \mathcal{D})$ , even those that don't correspond to normal form verifiers! This transformation can be performed in time that is polynomial in the length of the descriptions of  $\mathcal{S}$  and  $\mathcal{D}$ , and always outputs a pair of Turing machines  $(\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$ . However, the following proposition establishes that if  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is furthermore a normal form verifier, then so is  $\mathcal{V}^{\text{ANCH}} = (\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$  with certain completeness, soundness, and complexity properties.

**Proposition 11.3.** *If  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is a normal form verifier, then  $\mathcal{V}^{\text{ANCH}} = (\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$  is a normal form verifier satisfying the following for all  $n \in \mathbb{N}$ .*

1. (**Completeness**) *If there is a value-1 PCC strategy for  $\mathcal{V}_n$ , then there is a value-1 PCC strategy for  $\mathcal{V}_n^{\text{ANCH}}$ .*
2. (**Soundness**) *For all  $\epsilon > 0$ , we have  $\mathcal{E}(\mathcal{V}_n^{\text{ANCH}}, 1 - \epsilon) \geq \mathcal{E}(\mathcal{V}_n, 1 - (4 \cdot 16^2)\epsilon)$ .*

3. (**Complexity**) The time complexities of the verifier  $\mathcal{V}^{\text{ANCH}}$  satisfy

$$\begin{aligned}\text{TIME}_{\mathcal{S}^{\text{ANCH}}}(n) &= \text{poly}(\text{TIME}_{\mathcal{S}}(n)), \\ \text{TIME}_{\mathcal{D}^{\text{ANCH}}}(n) &= \text{poly}(\text{TIME}_{\mathcal{D}}(n)).\end{aligned}$$

Furthermore the number of levels of  $\mathcal{S}^{\text{ANCH}}$  is  $\ell + 2$  and the dimension is  $s(n) + 8$ , where the number of levels and dimension of sampler  $\mathcal{S}$  is  $\ell$  and  $s(n)$  respectively.

4. (**Efficient computability**) The descriptions of  $\mathcal{S}^{\text{ANCH}}$  and  $\mathcal{D}^{\text{ANCH}}$  can be computed in polynomial time from the descriptions of  $\mathcal{S}$  and  $\mathcal{D}$ , respectively. In particular, the sampler  $\mathcal{S}^{\text{ANCH}}$  only depends on the sampler  $\mathcal{S}$ .

*Proof.* If  $\mathcal{S}$  is an  $\ell$ -level sampler with dimension  $s(n)$ , then by construction  $\hat{\mathcal{S}}^{\text{ANCH}}$  is a  $(\mathcal{T}^{\text{ANCH}}, \mathcal{G}^{\text{ANCH}})$ -typed sampler, with field size  $q(n) = 2$  and the same dimension  $s(n)$ . It has the following CL functions. Fix an integer  $n \in \mathbb{N}$ . Let  $V = \mathbb{F}_2^{s(n)}$  denote the ambient space of  $\mathcal{S}$  on index  $n$ . Let  $L^A, L^B : V \rightarrow V$  denote the CL functions of  $\mathcal{S}$  on index  $n$ . For  $w \in \{A, B\}$  the associated CL functions  $\{L_t^{\text{ANCH}, w}\}$  of  $\hat{\mathcal{S}}^{\text{ANCH}}$  are

$$L_t^{\text{ANCH}, w} = \begin{cases} L^w & \text{if } t = \text{GAME}, \\ 0 & \text{if } t = \text{ANCHOR}. \end{cases}$$

Intuitively, when the type  $t$  sampled for player  $w$  is GAME, then they receive a question  $L^w(z)$  as they would according to  $\mathcal{S}$ . Otherwise if  $t = \text{ANCHOR}$ , then their question is the zero string. Thus if  $L^w$  is an  $\ell$ -level CL function, then  $L_{\text{GAME}}^{\text{ANCH}, w}$  is also an  $\ell$ -level CL function, and  $L_{\text{ANCHOR}}^{\text{ANCH}, w}$  is a 0-level CL function.

We analyze the completeness, soundness, and complexity properties of the typed verifier  $\hat{\mathcal{V}}^{\text{ANCH}}$ ; the corresponding properties of the detyped verifier  $\mathcal{V}^{\text{ANCH}}$  follow from Lemma 6.18 and the fact that the type set  $\mathcal{T}^{\text{ANCH}}$  has size 2.

Fix an index  $n \in \mathbb{N}$ . For the completeness property, let  $\mathcal{S}$  be a value-1 PCC strategy for  $\mathcal{V}_n$ . We define a value-1 PCC strategy  $\mathcal{S}^{\text{ANCH}}$  for  $\hat{\mathcal{V}}_n^{\text{ANCH}}$ : whenever a player receives the ANCHOR type as a question type, they perform the trivial measurement (i.e. measure the identity operator), and output 0. Otherwise, the player performs the same measurement as in  $\mathcal{S}$ . This is clearly value-1 and PCC. Item 1 follows from this and Lemma 6.18.

For the soundness property, we observe that if a strategy  $\mathcal{S}^{\text{ANCH}}$  has value  $1 - \varepsilon$  in  $\hat{\mathcal{V}}_n^{\text{ANCH}}$ , then

$$1 - \varepsilon = \frac{3}{4} + \frac{p}{4}$$

where  $p$  is the value of  $\mathcal{S}^{\text{ANCH}}$  in the game  $\mathcal{V}_n$ , and  $1/4$  is the probability that neither player receives the question type ANCHOR; this follows from the distribution associated with the typed sampler  $\hat{\mathcal{S}}^{\text{ANCH}}$ . Solving for  $p$ , this implies that  $\mathcal{E}(\hat{\mathcal{V}}_n^{\text{ANCH}}, 1 - \varepsilon) \geq \mathcal{E}(\mathcal{V}_n, 1 - 4\varepsilon)$ . The soundness property (Item 2) then follows from Lemma 6.18.

Items 3 and 4 are straightforward and also follow from Lemma 6.18.  $\square$

## 11.2 The parallel repetition transformation

Starting with the anchored verifier  $\mathcal{V}^{\text{ANCH}} = (\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$ , we then perform the parallel repetition transformation.

Fix integers  $\lambda, \tau \in \mathbb{N}$ , and let  $k(n) = (\lambda n)^{(1+c')\tau}$  where  $c' > 0$  is the universal constant such that  $\text{TIME}_{\mathcal{D}^{\text{ANCH}}}(n) \leq c'(\text{TIME}_{\mathcal{D}}(n))^{c'}$  for all  $n \geq 1$ . Define the Turing machine  $\mathcal{S}^{\text{REP}}$  (depending on  $\lambda, \tau$ ) as follows.



1. On input  $(n, \text{DIMENSION})$ , it computes  $s' = \mathcal{S}^{\text{ANCH}}(n, \text{DIMENSION})$  and  $k = k(n)$ , interprets  $s'$  as a positive integer and outputs  $k \cdot s'$ .
2. On input  $(n, w, \text{MARGINAL}, j, z)$ , it computes  $s' = \mathcal{S}^{\text{ANCH}}(n, \text{DIMENSION})$  and  $k = k(n)$ , parses  $z = (z_1, \dots, z_k) \in (\mathbb{F}_2^{s'})^k$ , and then outputs

$$\left( \mathcal{S}^{\text{ANCH}}(n, w, \text{MARGINAL}, j, z_1), \dots, \mathcal{S}^{\text{ANCH}}(n, w, \text{MARGINAL}, j, z_k) \right).$$

3. On input  $(n, w, \text{LINEAR}, j, u, y)$ , it computes  $s' = \mathcal{S}^{\text{ANCH}}(n, \text{DIMENSION})$  and  $k = k(n)$ , parses  $u = (u_1, \dots, u_k), y = (y_1, \dots, y_k) \in (\mathbb{F}_2^{s'})^k$ , and then outputs

$$\left( \mathcal{S}^{\text{ANCH}}(n, w, \text{LINEAR}, j, u_1, y_1), \dots, \mathcal{S}^{\text{ANCH}}(n, w, \text{LINEAR}, j, u_k, y_k) \right).$$

4. On input  $(n, w, \text{FACTOR}, j, u)$ , it computes  $s' = \mathcal{S}^{\text{ANCH}}(n, \text{DIMENSION})$  and  $k = k(n)$ , parses  $u = (u_1, \dots, u_k) \in (\mathbb{F}_2^{s'})^k$ , and then outputs

$$\left( \mathcal{S}^{\text{ANCH}}(n, w, \text{FACTOR}, j, u_1), \dots, \mathcal{S}^{\text{ANCH}}(n, w, \text{FACTOR}, j, u_k) \right).$$

Intuitively, the Turing machine  $\mathcal{D}^{\text{REP}}$  behaves as follows. On input  $(n, x, y, a, b)$ , Parse  $x, y, a, b$  as  $k(n)$ -tuples of questions and answers, respectively, and accept if and only if  $\mathcal{D}^{\text{ANCH}}(n, x_i, y_i, a_i, b_i)$  accepts for all  $i \in \{1, \dots, k(n)\}$ . One potential issue is that, depending on how this parsing is implemented, the time complexity of the decider could be unbounded (for example, if  $x$  is not properly formatted as a  $k(n)$ -tuple, then  $\mathcal{D}^{\text{REP}}$  could take time that grows with the length of  $x$ , which could be much greater than  $\text{TIME}_{\mathcal{D}^{\text{ANCH}}}(n)$ ). To avoid this issue, the decider  $\mathcal{D}^{\text{REP}}$  checks if any component of the tuples  $x, y, a, b$  have length larger than  $(\lambda n)^\tau$  bits, and if so then it rejects (this includes the case that  $x, y, a, b$  are not properly formatted as  $k(n)$ -tuples). Thus, as long as the time complexity  $\text{TIME}_{\mathcal{D}^{\text{ANCH}}}(n)$  is at most  $(\lambda n)^{\tau c'}$ , then the strings  $x, y, a, b$  never need to be longer than  $k(n) \cdot (\lambda n)^\tau = (\lambda n)^{(1+2c')\tau}$  bits long in order to be accepted by the decider.

Note again that this transformation is well-defined for all Turing machines  $(\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$ . The next theorem shows that, if  $(\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$  is the anchoring of a normal form verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$ , then  $\mathcal{V}^{\text{REP}} = (\mathcal{S}^{\text{REP}}, \mathcal{D}^{\text{REP}})$  is also a normal form verifier with certain completeness, soundness, and complexity properties. This is the main result of the section.

**Theorem 11.4** (Anchored parallel repetition of normal form verifiers). *There exist universal constants  $c, c' > 0$  and a polynomial-time Turing machine  $\text{ComputeRepeatedVerifier}$  that given input a tuple  $(\mathcal{V}, \lambda, \tau)$  where  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is a pair of Turing machines and  $\lambda, \tau$  are positive integers, outputs a pair of Turing machines  $\mathcal{V}^{\text{REP}} = (\mathcal{S}^{\text{REP}}, \mathcal{D}^{\text{REP}})$  such that the following holds. If  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is an  $\ell$ -level normal form verifier, then the output  $\mathcal{V}^{\text{REP}}$  is a normal form verifier satisfying, for all  $n \in \mathbb{N}$ , letting  $k(n) = (\lambda n)^{(1+c')\tau}$ ,*

1. (**Completeness**) *If  $\mathcal{V}_n$  has a value-1 PCC strategy and  $\text{TIME}_{\mathcal{D}}(n) \leq (\lambda n)^\tau$ , then  $\mathcal{V}_n^{\text{REP}}$  has a value-1 PCC strategy.*
2. (**Soundness**) *For all  $\varepsilon > 0$ , for all*

$$p > \frac{4}{\varepsilon} \exp\left(-c \varepsilon^{17} k(n) / (\lambda n)^{\tau c'}\right),$$

*we have  $\mathcal{E}(\mathcal{V}_n^{\text{REP}}, p) \geq \mathcal{E}(\mathcal{V}_n, 1 - \varepsilon)$ .*

3. (**Complexity**) The repeated verifier  $\mathcal{V}^{\text{REP}}$  is  $(\ell + 2)$ -level and has time complexities

$$\begin{aligned}\text{TIME}_{\mathcal{S}^{\text{REP}}}(n) &= O(k(n) \cdot \text{TIME}_{\mathcal{S}}(n)), \\ \text{TIME}_{\mathcal{D}^{\text{REP}}}(n) &= O\left(k(n) \cdot \max\left(\text{TIME}_{\mathcal{D}}(n), (\lambda n)^\tau\right)\right).\end{aligned}$$

Furthermore, the repeated sampler  $\mathcal{S}^{\text{REP}}$  only depends on  $\mathcal{S}$  and the parameters  $\lambda, \tau$ .

*Proof.* The Turing machine `ComputeRepeatedVerifier`, given input  $(\mathcal{V}, \lambda, \tau)$ , first computes the description of the Turing machines  $(\hat{\mathcal{S}}^{\text{ANCH}}, \hat{\mathcal{D}}^{\text{ANCH}})$  and then their detyping  $(\mathcal{S}^{\text{ANCH}}, \mathcal{D}^{\text{ANCH}})$  as described in Section 11.1. Then, it computes the pair of Turing machines  $(\mathcal{S}^{\text{REP}}, \mathcal{D}^{\text{REP}})$  as described above. Clearly `ComputeRepeatedVerifier` runs in polynomial time.

Suppose that  $\mathcal{S}$  is an  $\ell$ -level sampler with dimension  $s(n)$ . Then by Proposition 11.3, the number of levels of  $\mathcal{S}^{\text{ANCH}}$  is  $\ell' = \ell + 2$ , and its dimension is  $s'(n) = s(n) + 8$ . Then by construction the Turing machine  $\mathcal{S}^{\text{REP}}$  is a sampler with the following properties. It has field size  $q(n) = 2$  and it has dimension  $s^{\text{REP}}(n) = k(n)s'(n)$ . We treat the ambient space  $V^{\text{REP}}$  of  $\mathcal{S}^{\text{REP}}$  on index  $n$  as the  $k(n)$ -fold direct sum of the ambient space  $V^{\text{ANCH}}$  of  $\mathcal{S}^{\text{ANCH}}$  on index  $n$ . For all integers  $n \in \mathbb{N}$ ,  $w \in \{A, B\}$ , the CL functions  $L^{\text{REP}, w} : V^{\text{REP}} \rightarrow V^{\text{REP}}$  of  $\mathcal{S}^{\text{REP}}$  are as follows:

$$L^{\text{REP}, w} = \bigoplus_{i=1}^{k(n)} L^w,$$

where  $L^A, L^B : V^{\text{ANCH}} \rightarrow V^{\text{ANCH}}$  are the CL functions of the sampler  $\mathcal{S}^{\text{ANCH}}$  on index  $n$ . The CL functions  $L^{\text{REP}, w}$  are  $\ell$ -level; the  $j$ -th factor spaces  $V_{j, u}^{\text{REP}, w}$  of  $L^{\text{REP}, w}$  are defined as

$$V_{j, u}^{\text{REP}, w} = \bigoplus_{i=1}^{k(n)} V_{j, u_i}^w$$

for all  $u = (u_1, \dots, u_{k(n)})$  where  $V_{j, u_i}^w$  is the  $j$ -th factor space of  $L^w$  with prefix  $u_i \in V^{\text{ANCH}}$ .

The number of levels of  $\mathcal{S}^{\text{REP}}$  is the same as  $\mathcal{S}^{\text{ANCH}}$ , which is  $\ell' = \ell + 2$ .

Thus, assuming that  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is a normal form verifier, we obtain that  $\mathcal{V}^{\text{REP}} = (\mathcal{S}^{\text{REP}}, \mathcal{D}^{\text{REP}})$  is a normal form verifier, and thus defines an infinite family of games  $(\mathcal{V}_n^{\text{REP}})_{n \in \mathbb{N}}$ .

Item 1 follows from the following straightforward observation: if  $\mathcal{S} = (\psi, A, B)$  is a value-1 PCC strategy for  $\mathcal{V}_n$  and  $\text{TIME}_{\mathcal{D}}(n) \leq (\lambda n)^\tau$ , then it must be that the question and answer lengths in the strategy  $\mathcal{S}$  are at most  $(\lambda n)^\tau$ . Now consider the strategy where the players share  $k(n)$  copies of  $|\psi\rangle$ , and for the  $i$ -th instance of the game  $\mathcal{V}_n^{\text{ANCH}}$ , the players use strategy  $\mathcal{S}$  on the  $i$ -th copy of  $|\psi\rangle$  (and performing the identity measurement whenever they receive the ANCHOR type). The total length of the questions and answers in the repeated game are at most  $k(n) \cdot (\lambda n)^\tau$ . It is straightforward to check that this strategy has value 1 and is PCC.

To show Item 2 we apply Theorem 11.1 and Proposition 11.3. The exponential decay bound on the value of  $\mathcal{V}_n^{\text{REP}}$  presented in Theorem 11.1 depends on the answer length of the players in the game  $\mathcal{V}_n^{\text{ANCH}}$ . By Definition 5.33, this answer length is at most  $\text{TIME}_{\mathcal{D}^{\text{ANCH}}}(n) = c'(\text{TIME}_{\mathcal{D}}(n))^{c'} \leq (\lambda n)^{\tau c'}$  for some universal constant  $c' > 0$  so the claimed bound follows.

Item 3 follows from the fact that computing the direct sum of  $k(n)$  CL functions and factor spaces of the “single-copy” sampler  $\mathcal{S}^{\text{ANCH}}$  requires  $k(n)$  times the complexity of the “single-copy” sampler, and the complexity of  $\mathcal{S}^{\text{ANCH}}$  follows from Proposition 11.3. The time complexity of the decider follows from the

repeated decider having to run  $k(n)$  instances of the decider  $\mathcal{D}^{\text{ANCH}}$  and take the logical AND of the  $k(n)$  decider outputs, as well as the fact that the decider rejects (and thus halts) in the case that the question and answer tuples are improperly formatted. The complexity of  $\mathcal{D}^{\text{ANCH}}$  follows from Proposition 11.3 which in turn runs an instance of the original decider  $\mathcal{D}$ . Since the CL functions of the sampler  $\mathcal{S}^{\text{ANCH}}$  are  $(\ell + 2)$ -level (by Proposition 11.3), and taking the direct sum of CL functions does not increase the number of levels (by Lemma 4.8), the CL functions  $L^{\text{REP}, w}$  are  $(\ell + 2)$ -level.

The “Furthermore” part of the theorem can be easily verified by inspecting the construction of the samplers  $\mathcal{S}^{\text{REP}}$  and  $\mathcal{S}^{\text{ANCH}}$ . □

## 12 Gap-Preserving Compression

We combine the transformations from the previous sections (the introspection games, answer reduction, and parallel repetition) to obtain our main technical result, a *gap-preserving compression theorem* (Theorem 12.1) for normal form verifiers.

Recall from Definition 5.32 that a verifier  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is  $\lambda$ -bounded if the description length  $|\overline{\mathcal{V}}| = \max\{|\mathcal{S}|, |\mathcal{D}|\}$  is at most  $\lambda$  and the time complexity bounds  $\text{TIME}_{\mathcal{S}}(n)$  and  $\text{TIME}_{\mathcal{D}}(n)$  are at most  $n^\lambda$  for all  $n \geq 2$ . The compression theorem states the existence of an efficient “compression procedure” `Compress` that achieves the following. Given as input a  $\lambda$ -bounded verifier  $\mathcal{V}$  and the parameter  $\lambda$ , the procedure returns a “compressed verifier”  $\mathcal{V}^{\text{COMPR}}$  such that the  $n$ -th game  $\mathcal{V}_n^{\text{COMPR}}$  simulates (in a sense to be made precise shortly) the  $N$ -th game  $\mathcal{V}_N$  for  $N = 2^n$ , and furthermore both the sampler time complexity and the decider time complexity of  $\mathcal{V}_n^{\text{COMPR}}$  can be exponentially smaller than those of  $\mathcal{V}_N$ , which can be as large as  $N^\lambda$ . We note that the time complexity bounds of the compressed verifier hold without the assumption that the input verifier is  $\lambda$ -bounded. The completeness and soundness bounds of the theorem require the  $\lambda$ -boundedness of the input verifier  $\mathcal{V}$ .

**Theorem 12.1** (Compression theorem). *There exists a universal constant  $C_0 > 0$  and a polynomial-time Turing machine `Compress` that takes as input a tuple  $(\mathcal{V}, \lambda)$  where  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is a pair of Turing machines and  $\lambda > 0$  is an integer, and outputs a 9-level normal form verifier  $\mathcal{V}^{\text{COMPR}} = (\mathcal{S}^{\text{COMPR}}, \mathcal{D}^{\text{COMPR}})$ . The time complexities of  $\mathcal{S}^{\text{COMPR}}, \mathcal{D}^{\text{COMPR}}$  satisfy*

$$\begin{aligned}\text{TIME}_{\mathcal{S}^{\text{COMPR}}}(n) &= \text{poly}(n, \lambda), \\ \text{TIME}_{\mathcal{D}^{\text{COMPR}}}(n) &= \text{poly}(n, \lambda).\end{aligned}$$

*The description of  $\mathcal{S}^{\text{COMPR}}$  is independent of  $\mathcal{V}$  and is computable from the binary representation of  $\lambda$  in time  $\text{polylog}(\lambda)$ . If furthermore  $\mathcal{V}$  is a  $\lambda$ -bounded, 9-level normal form verifier, then  $\mathcal{V}^{\text{COMPR}}$  satisfies the following: for all  $n \geq C_0$  and  $N = 2^n$ ,*

1. (**Completeness**) *If  $\mathcal{V}_N$  has a value-1 PCC strategy, then  $\mathcal{V}_n^{\text{COMPR}}$  has a value-1 PCC strategy.*
2. (**Soundness**)  $\mathcal{E}(\mathcal{V}_n^{\text{COMPR}}, \frac{1}{2}) \geq \max \left\{ \mathcal{E}(\mathcal{V}_N, \frac{1}{2}), 2^{N^\lambda - 1} \right\}$ .

### 12.1 Proof of Theorem 12.1

Recall the following Turing machines.

1. `ComputeIntroVerifier` takes as input a tuple  $(\mathcal{V}, \lambda, \ell)$  and returns in polynomial time a description of the introspective verifier  $\mathcal{V}^{\text{INTRO}}$  corresponding to the verifier  $\mathcal{V}$  and parameters  $(\lambda, \ell)$ . (See Theorem 8.3.)
2. `ComputeAnsRedVerifier` takes input  $(\mathcal{V}, \lambda, \mu, \gamma)$  and returns in polynomial time a description of the answer reduced verifier  $\mathcal{V}^{\text{AR}}$  corresponding to  $\mathcal{V}$  and parameters  $(\lambda, \mu, \gamma)$ . (See Theorem 10.27.)
3. `ComputeRepeatedVerifier` takes input  $(\mathcal{V}, \lambda, \tau)$  and returns in polynomial time a description of the anchored repeated verifier  $\mathcal{V}^{\text{REP}}$  corresponding to  $\mathcal{V}$  and a number of repetitions  $k(n) = (\lambda n)^\tau$ . (See Theorem 11.4.)

We specify the Turing machine Compress in Fig. 15. The universal constants  $\mu$  and  $\gamma$  are specified in Eq. (158), and the universal constant  $\tau$  is specified in Eq. (163).

Input:  $(\mathcal{V}, \lambda)$  where  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  is a pair of Turing machines and  $\lambda$  is an integer.

1. Compute  $\mathcal{V}^{(1)} = \text{ComputeIntroVerifier}(\mathcal{V}, \lambda, 9)$ .
2. Compute  $\mathcal{V}^{(2)} = \text{ComputeAnsRedVerifier}(\mathcal{V}^{(1)}, \lambda, \mu, \gamma)$ .
3. Compute  $\mathcal{V}^{(3)} = \text{ComputeRepeatedVerifier}(\mathcal{V}^{(2)}, \lambda, \tau)$ .
4. Return  $\mathcal{V}^{\text{COMPR}} = \mathcal{V}^{(3)}$ .

Figure 15: The definition of the Turing machine Compress, where the universal constants  $\mu$  and  $\gamma$  are specified in Eq. (158), and the universal constant  $\tau$  is specified in Eq. (163).

**Lemma 12.2.** *Let Compress be the Turing machine specified in Fig. 15. For all pairs of Turing machines  $\mathcal{V} = (\mathcal{S}, \mathcal{D})$  and integer  $\lambda$ , the output of  $\text{Compress}(\mathcal{V}, \lambda)$  is a normal form verifier  $\mathcal{V}^{\text{COMPR}} = (\mathcal{S}^{\text{COMPR}}, \mathcal{D}^{\text{COMPR}})$  such that  $\mathcal{S}^{\text{COMPR}}$  does not depend on  $\mathcal{V}$  but only on the parameter  $\lambda$ . Furthermore, there is a Turing machine ComputeSampler that on input  $\lambda$  outputs the description of the compressed sampler  $\mathcal{S}^{\text{COMPR}}$  in time  $\text{polylog}(\lambda)$ .*

*Proof.* The verifier  $\mathcal{V}^{\text{COMPR}} = \mathcal{V}^{(3)}$  is a normal form verifier because the introspective verifier  $\mathcal{V}^{(1)}$  is a normal form verifier (even if the input  $\mathcal{V}$  is not; see Remark 8.7), and the verifier transformations ComputeAnsRedVerifier, ComputeRepeatedVerifier preserve the normal form (by Theorems 10.27 and 11.4).

The introspective verifier computed in Step 1 of Figure 15 has a sampler  $\mathcal{S}^{(1)}$  that only depends on the parameter  $\lambda$ , as implied by Lemma 8.1. The answer reduced verifier from Step 2 has a sampler  $\mathcal{S}^{(2)}$  that, according to Theorem 10.27, only depends on the sampler  $\mathcal{S}^{(1)}$ , the parameter tuple  $(\lambda, \mu, \gamma)$ , and the description length  $|\mathcal{D}^{(1)}|$ . Note that  $\mathcal{S}^{(1)}$  only depends on  $\lambda$ , the parameters  $(\mu, \gamma)$  are universal constants, and the description length of  $\mathcal{D}^{(1)}$  is at most the time complexity of ComputeIntroVerifier on input  $(\mathcal{V}^{(0)}, \lambda, 9)$ , which is polynomial in  $\lambda$ . Thus  $\mathcal{S}^{(2)}$  only depends on  $\lambda$ . Similarly, the sampler  $\mathcal{S}^{(3)}$  of the repeated verifier from Step 3 only depends on the sampler  $\mathcal{S}^{(2)}$ , the parameter  $\lambda$ , and the universal constant  $\tau$ , as implied by Theorem 11.4. Therefore the final sampler  $\mathcal{S}^{\text{COMPR}} = \mathcal{S}^{(3)}$  only depends on  $\lambda$ .

Let  $\mathcal{V}^*$  be the pair of trivial Turing machines  $(0, 0)$ . Define ComputeSampler as the Turing machine that on input  $\lambda$  computes  $(\mathcal{S}, \mathcal{D}^{\text{DUMMY}}) = \text{Compress}(\mathcal{V}^*, \lambda)$  and returns the description of  $\mathcal{S}$  as the output. By the above discussion, the description of  $\mathcal{S}$  is the same as that of  $\mathcal{S}^{\text{COMPR}}$  and the time complexity of ComputeSampler is  $\text{polylog}(\lambda)$  as Compress runs in polynomial time: this follows from the fact that the Turing machines ComputeIntroVerifier, ComputeAnsRedVerifier, and ComputeRepeatedVerifier all run in polynomial time.  $\square$

*Proof of Theorem 12.1.* We evaluate the parameters of the verifiers generated in each step of the Compress procedure. These parameters are summarized in Table 2 and are explained in the following. Let  $(\mathcal{V}, \lambda)$  be the input to the Turing machine Compress. We write  $N$  to denote  $2^n$ .

**Introspection.** The verifier  $\mathcal{V}^{(1)} = (\mathcal{S}^{(1)}, \mathcal{D}^{(1)})$  is the introspective verifier corresponding to  $\mathcal{V}$  and parameters  $\lambda$  and  $\ell = 9$  computed by ComputeIntroVerifier. We state its parameters and properties in the second row of Table 2 in terms of the index  $n \in \mathbb{N}$ .

	Time complexity		Level	Completeness	Soundness
	Sampler	Decider			
$\mathcal{V}^{(1)}$	$\text{poly}(n, \lambda)$	$\text{poly}(N^\lambda)$	5	value-1 PCC	$\mathcal{E}(\mathcal{V}_n^{(1)}, 1 - \varepsilon_1) \geq \max \{ \mathcal{E}(\mathcal{V}_N, \frac{1}{2}), 2^{N^\lambda - 1} \}$
$\mathcal{V}^{(2)}$	$\text{poly}(n, \lambda)$	$\text{poly}(n, \lambda)$	7	value-1 PCC	$\mathcal{E}(\mathcal{V}_n^{(2)}, 1 - \varepsilon_2) \geq \mathcal{E}(\mathcal{V}_n^{(1)}, 1 - \varepsilon_1)$
$\mathcal{V}^{(3)}$	$\text{poly}(n, \lambda)$	$\text{poly}(n, \lambda)$	9	value-1 PCC	$\mathcal{E}(\mathcal{V}_n^{(3)}, \frac{1}{2}) \geq \mathcal{E}(\mathcal{V}_n^{(2)}, 1 - \varepsilon_2)$

Table 2: Parameters and properties of the input verifier  $\mathcal{V}$ , introspective verifier  $\mathcal{V}^{(1)}$ , answer reduced verifier  $\mathcal{V}^{(2)}$ , and parallel repeated verifier  $\mathcal{V}^{(3)}$  computed by Compress. The quantities  $\varepsilon_1, \varepsilon_2$  (which are functions of  $n$ ) are specified below in the main text.

The complexity bounds and the number of levels of the introspective verifier are given by Theorem 8.3. Using that the argument  $\ell$  in `ComputeIntroVerifier` is set to 9, there exists a universal constant  $C_{\text{INTRO}} \geq 1$  such that for all  $\lambda \in \mathbb{N}$  and for all  $n \geq C_{\text{INTRO}}$ , we have

$$\begin{aligned}
\text{TIME}_{\mathcal{S}^{(1)}}(n) &\leq (\lambda \cdot n)^{C_{\text{INTRO}}}, \\
\text{TIME}_{\mathcal{D}^{(1)}}(n) &\leq 2^{C_{\text{INTRO}} \lambda n}, \\
|\mathcal{D}^{(1)}| &\leq C_{\text{INTRO}} \lambda^{C_{\text{INTRO}}}.
\end{aligned} \tag{155}$$

By Theorem 8.3 these bounds hold for all inputs  $(\mathcal{V}, \lambda, 9)$  (in particular these bounds do not assume that the input  $\mathcal{V}$  is  $\lambda$ -bounded or is even a normal form verifier).

The statements in the completeness and soundness entries of the row assume that the input verifier  $\mathcal{V}$  is 9-level and  $\lambda$ -bounded. The completeness entry follows from Theorem 8.3 because in the completeness case we assume the verifier  $\mathcal{V}$  has a value-1 PCC strategy, and thus the resulting introspective verifier has a value-1 PCC strategy. We deduce the soundness entry as follows. Let the function  $\delta(\varepsilon, n)$  from Theorem 8.3 be denoted by  $\delta_1(\varepsilon_1, n)$ . Let  $a_1 > 0, 0 < b_1 < 1$  be the universal constants such that

$$\delta_1(\varepsilon_1, n) = a_1 \left( (\lambda n)^{a_1} \cdot \varepsilon_1^{b_1} + (\lambda n)^{-b_1} \right) \tag{156}$$

By setting

$$\varepsilon_1 = \left( \frac{1}{8a_1(\lambda n)^{a_1}} \right)^{1/b_1}, \quad C_1 = (4a_1)^{1/b_1} \tag{157}$$

we get for all  $n \geq C_1$ ,

$$\begin{aligned}
\delta_1(\varepsilon_1, n) &= a_1 \left( (\lambda n)^{a_1} \cdot (\varepsilon_1)^{b_1} + (\lambda n)^{-b_1} \right) \\
&\leq a_1 \left( (\lambda n)^{a_1} \cdot (\varepsilon_1)^{b_1} \right) + \frac{1}{4} \\
&\leq a_1 \left( \frac{1}{8a_1} \right) + \frac{1}{4} \\
&< \frac{1}{2},
\end{aligned}$$

where we used the fact that  $\lambda \geq 1$  in the second line. Thus by the soundness guarantee of Theorem 8.3 we get that

$$\mathcal{E}(\mathcal{V}_n^{(1)}, 1 - \varepsilon_1) \geq \max \left\{ \mathcal{E}(\mathcal{V}_N, \frac{1}{2}), 2^{N^\lambda - 1} \right\}$$

for all  $n \geq C_1$  as desired.

**Answer reduction.** Let  $\mathcal{V}^{(2)} = (\mathcal{S}^{(2)}, \mathcal{D}^{(2)})$  denote the answer reduced verifier corresponding to  $\mathcal{V}^{(1)}$  and parameters  $(\lambda, \mu, \gamma)$  computed by `ComputeAnsRedVerifier` where we define  $\mu$  and  $\gamma$  as

$$\mu = \lceil C_{\text{INTRO}} \rceil, \quad \gamma = \left\lceil \frac{2a_1}{b_1 b_2} \right\rceil \quad (158)$$

where  $a_1, b_1$  are the universal constants defining  $\delta_1$  in Eq. (156), and  $b_2$  is defined below in Eq. (161). The third row in Table 2 gives the properties of the answer reduced verifier.

The bounds on the number of levels and the time complexities follow from Theorem 10.27: the number of levels of the introspective verifier  $\mathcal{V}^{(1)}$  is 5, and therefore the number of levels of the sampler  $\mathcal{S}^{(2)}$  is 7. Furthermore, using the bounds on the complexities of  $\mathcal{S}^{(1)}$ ,  $\mathcal{D}^{(1)}$ , and the bound on the description length  $|\mathcal{D}^{(1)}|$  from Eq. (155), we get that assumption (134) is satisfied and thus by Theorem 10.27 there exists a universal constant  $C_{\text{AR}} \geq 1$  such that for all  $n \geq C_{\text{AR}}$ ,

$$\max\left\{\text{TIME}_{\mathcal{S}^{(2)}}(n), \text{TIME}_{\mathcal{D}^{(2)}}(n)\right\} \leq (\lambda \cdot n)^{C_{\text{AR}}}. \quad (159)$$

The constant  $C_{\text{AR}}$  depends on the universal constants  $\mu, \gamma$ . Again, these time complexities do not depend on whether the starting verifier  $\mathcal{V}$  is  $\lambda$ -bounded.

For the soundness and completeness entries, we assume that the starting verifier  $\mathcal{V}$  is  $\lambda$ -bounded. If we further assume that  $\mathcal{V}$  has a value-1 PCC strategy, then  $\mathcal{V}^{(1)}$  has a value-1 PCC strategy (by the completeness guarantee of the introspective verifier), and then the completeness part of Theorem 10.27 guarantees that  $\mathcal{V}^{(2)}$  also has a value-1 PCC strategy.

The soundness entry is argued as follows. Define  $\delta_2(\varepsilon_2, n)$  to be the function  $\delta(\varepsilon, n)$  from Theorem 10.27; write it as

$$\delta_2(\varepsilon_2, n) = a' \cdot \gamma^{a'} \cdot \left( (\lambda \cdot |\mathcal{D}^{(1)}| \cdot n)^{\mu a'} \cdot \varepsilon_2^{b'} + (\lambda \cdot |\mathcal{D}^{(1)}| \cdot n)^{-\mu b' \gamma} \right) \quad (160)$$

for universal constants  $a' > 0$  and  $0 < b' < 1$ . By our bound on the description length of  $\mathcal{D}^{(1)}$  from Eq. (155), we have that  $\delta_2$  can be bounded by

$$\delta_2(\varepsilon_2, n) \leq a_2 \left( (\lambda n)^{a_2} \cdot \varepsilon_2^{b_2} + (\lambda n)^{-c_2} \right)$$

where we set

$$a_2 = \max\{a' \gamma^{a'}, a' \gamma^{a'} (C_{\text{INTRO}})^{\mu a'}, (1 + C_{\text{INTRO}}) \mu a'\}, \quad b_2 = b', \quad c_2 = \mu b' \gamma. \quad (161)$$

Since  $a', b', \mu, \gamma$  and  $C_{\text{INTRO}}$  are universal constants, so are  $a_2, b_2, c_2$ . By setting

$$\varepsilon_2 = \left( \frac{\varepsilon_1}{8a_2(\lambda n)^{a_2}} \right)^{1/b_2}, \quad C_2 = (4a_2)^{b_1/a_1} (8a_1)^{1/a_1}, \quad (162)$$

we get for all  $n \geq C_2$ ,

$$\begin{aligned}
\delta_2(\varepsilon_2, n) &\leq a_2 \left( (\lambda n)^{a_2} \cdot \varepsilon_2^{b_2} + (\lambda n)^{-b_2 \gamma} \right) && (\mu \geq 1) \\
&= a_2 \cdot (\lambda n)^{a_2} \cdot \varepsilon_2^{b_2} + a_2 \cdot (\lambda n)^{-b_2 \gamma} \varepsilon_1^{-1} \cdot \varepsilon_1 \\
&= a_2 \cdot (\lambda n)^{a_2} \cdot \varepsilon_2^{b_2} + a_2 \cdot (\lambda n)^{-(b_2 \gamma - a_1/b_1)} (8a_1)^{1/b_1} \cdot \varepsilon_1 && \text{(Definition of } \varepsilon_1) \\
&\leq a_2 \cdot (\lambda n)^{a_2} \cdot \varepsilon_2^{b_2} + a_2 \cdot (\lambda n)^{-a_1/b_1} (8a_1)^{1/b_1} \cdot \varepsilon_1 && \text{(Definition of } \gamma) \\
&\leq a_2 \left( \frac{\varepsilon_1}{8a_2} \right) + \frac{\varepsilon_1}{4} && (n \geq C_2, \lambda \geq 1, \text{ def. of } \varepsilon_2) \\
&< \varepsilon_1.
\end{aligned}$$

Thus by the soundness guarantee of Theorem 10.27 we get that

$$\mathcal{E}(\mathcal{V}_n^{(2)}, 1 - \varepsilon_2) \geq \mathcal{E}(\mathcal{V}_n^{(1)}, 1 - \varepsilon_1)$$

for all  $n \geq C_2$  as desired.

**Parallel repetition.** Let  $\mathcal{V}^{(3)} = (\mathcal{S}^{(3)}, \mathcal{D}^{(3)})$  denote the anchored repetition (see Section 11.2) of  $\mathcal{V}^{(2)}$  computed by ComputeRepeatedVerifier, with parameters  $\lambda$  and  $\tau$  defined below in Eq. (163). We state the parameters and properties of  $\mathcal{V}^{(3)}$  in the fourth row of Table 2.

The number of levels of  $\mathcal{S}^{(3)}$  is 9 by Item 3 of Theorem 11.4, because the number of levels of  $\mathcal{V}^{(2)}$  is 7. The time complexities of  $\mathcal{S}^{(3)}$  and  $\mathcal{D}^{(3)}$  follow from Item 3 of Theorem 11.4 and the complexity upper bounds on  $\mathcal{S}^{(2)}$  and  $\mathcal{D}^{(2)}$  specified in the third row of Table 2; in particular this uses the fact that  $\text{TIME}_{\mathcal{D}^{(2)}}(n) \leq (\lambda \cdot n)^{C_{\text{AR}}} \leq (\lambda \cdot n)^\tau$  by the way we set  $\tau$  in Eq. (163). These bounds only depend on the bounds stated for verifier  $\mathcal{V}^{(2)}$  in the third row of Table 2, and again do not depend on whether the input verifier  $\mathcal{V}$  is  $\lambda$ -bounded.

For the soundness and completeness entries, we assume that the starting verifier  $\mathcal{V}$  is  $\lambda$ -bounded. The statement in the completeness entry follows from Item 1 of Theorem 11.4, where we assume that  $\mathcal{V}$  and thus  $\mathcal{V}^{(1)}$  and thus  $\mathcal{V}^{(2)}$  all have value-1 PCC strategies; therefore  $\mathcal{V}^{(3)}$  would have a value-1 PCC strategy as well.

The soundness entry is argued as follows. Let  $c_3, c'_3 > 0$  denote the universal constants  $c, c'$  from Item 2 of Theorem 11.4. Recalling that  $\varepsilon_2$  (defined in Eq. (162)) is a function of  $n$  and  $\lambda$ , define  $\tau$  to be the minimum integer satisfying  $\tau \geq C_{\text{AR}}$  and

$$(\lambda n)^\tau \geq \frac{1}{c_3 \varepsilon_2^{17}} \ln \frac{8}{\varepsilon_2} \tag{163}$$

for all  $n \geq \tau$  and for all integer  $\lambda \geq 1$ . The integer  $\tau$  is well-defined because the right-hand side is upper-bounded by a fixed polynomial in  $\lambda$ . Setting  $k(n) = (\lambda n)^{(1+c'_3)\tau}$ , we have that for all  $n \geq \max\{\tau, C_{\text{AR}}\}$

$$\frac{4}{\varepsilon_2} \exp\left(-c_3 \varepsilon_2^{17} k(n) / (\lambda n)^{\tau c_3}\right) = \frac{4}{\varepsilon_2} \exp\left(-c_3 \varepsilon_2^{17} (\lambda n)^\tau\right) < \frac{1}{2}$$

where we used our definition of  $k(n)$  and the definition of  $\tau$ . Thus by the soundness guarantee of Theorem 11.4 we get that

$$\mathcal{E}(\mathcal{V}_n^{(3)}, \frac{1}{2}) \geq \mathcal{E}(\mathcal{V}_n^{(2)}, 1 - \varepsilon_2)$$

for all  $n \geq \max\{\tau, C_{\text{AR}}\}$  as desired.



**Putting everything together.** The verifier  $\mathcal{V}^{\text{COMPR}}$  is  $\mathcal{V}^{(3)}$ . We now put together the bounds and parameters from Table 2 to obtain the conclusions stated in Theorem 12.1.

The claim that the Turing machine Compress is polynomial time follows from the fact that the Turing machines ComputeIntroVerifier, ComputeAnsRedVerifier, and ComputeRepeatedVerifier all run in polynomial time.

The claimed number of levels and time complexity of the sampler  $\mathcal{S}^{\text{COMPR}}$  follows from the complexity bounds in Table 2, which in turn only depend on the parameter  $\lambda$ . In particular the sampler  $\mathcal{S}^{\text{COMPR}}$  is independent of the input verifier  $\mathcal{V}$ , as shown in Lemma 12.2. Finally, the claimed time complexity to compute the description of  $\mathcal{S}^{\text{COMPR}}$  also follows from Lemma 12.2.

The claimed time complexity of  $\mathcal{D}^{\text{COMPR}}$  in the theorem statement also follows from Table 2.

We now establish the completeness and soundness properties of  $\mathcal{V}^{\text{COMPR}}$ . Assume that the input verifier  $\mathcal{V}$  is 9-level and  $\lambda$ -bounded. The completeness property (Item 1 in Theorem 12.1) follows from chaining together the completeness properties of  $\mathcal{V}_n^{(3)}$ ,  $\mathcal{V}_n^{(2)}$ ,  $\mathcal{V}_n^{(1)}$  and the assumption that  $\mathcal{V}_N$  has a value-1 PCC strategy.

We now establish the soundness property (Item 2 in Theorem 12.1). Set

$$C_0 = \max\{C_1, C_2, C_{\text{AR}}, \tau\}. \quad (164)$$

Since  $C_1, C_2, C_{\text{AR}}, \tau$  are all universal constants,  $C_0$  is also a universal constant. By chaining together the soundness properties of  $\mathcal{V}_n^{(3)}$ ,  $\mathcal{V}_n^{(2)}$ ,  $\mathcal{V}_n^{(1)}$ , we get that

$$\mathcal{E}(\mathcal{V}_n^{(3)}, \frac{1}{2}) \geq \mathcal{E}(\mathcal{V}_n^{(2)}, 1 - \varepsilon_2) \geq \mathcal{E}(\mathcal{V}_n^{(1)}, 1 - \varepsilon_1) \geq \max\{\mathcal{E}(\mathcal{V}_N, \frac{1}{2}), 2^{N^\lambda - 1}\}$$

for all  $n \geq C_0$  as desired. This concludes the proof of the theorem.  $\square$

## 12.2 An MIP\* protocol for the Halting problem

For every Turing machine  $\mathcal{M}$ , we construct a nonlocal game  $\mathfrak{G}$  such that  $\text{val}^*(\mathfrak{G}) = 1$  if  $\mathcal{M}$  halts and  $\text{val}^*(\mathfrak{G}) \leq \frac{1}{2}$  otherwise. In what follows, to explicitly disambiguate between a Turing machine (which is a tuple consisting of a set of states, transition rules, etc) and its description (which is a binary string), we use calligraphic letters to denote the Turing machine  $\mathcal{M}$  and an overline  $\overline{\mathcal{M}}$  to denote a *description* of  $\mathcal{M}$  (see Section 3.1 for a discussion of Turing machines and their descriptions).

First, we define a Turing machine  $\mathcal{F}$  as in Fig. 16. For all Turing machines  $\mathcal{M}$  and parameters  $\lambda \in \mathbb{N}$  we define the decider  $\mathcal{D}_{\mathcal{M}, \lambda}^{\text{HALT}}$  to be the 5-input Turing machine

$$\mathcal{D}_{\mathcal{M}, \lambda}^{\text{HALT}}(n, x, y, a, b) = \mathcal{F}(\overline{\mathcal{F}}, \overline{\mathcal{M}}, \lambda, n, x, y, a, b)$$

i.e., on input  $(n, x, y, a, b)$  runs  $\mathcal{F}$  with input  $(\overline{\mathcal{F}}, \overline{\mathcal{M}}, \lambda, n, x, y, a, b)$ . We generally omit the subscripts  $\mathcal{M}, \lambda$  for notational simplicity and denote the decider as  $\mathcal{D}^{\text{HALT}}$ . Define  $\mathcal{S}^{\text{HALT}} = \text{ComputeSampler}(\lambda)$  and  $\mathcal{V}^{\text{HALT}} = (\mathcal{S}^{\text{HALT}}, \mathcal{D}^{\text{HALT}})$ . By inspecting the definition of the Turing machine  $\mathcal{F}$  in Fig. 16, one can see that the description  $\overline{\mathcal{D}}$  computed by  $\mathcal{D}^{\text{HALT}}$  at Step 2 is the description of  $\mathcal{D}^{\text{HALT}}$  itself.<sup>35</sup> Therefore, the input to the Compress Turing machine at Step 4 is the description of  $\mathcal{V}^{\text{HALT}}$  and  $\lambda$ .

We note that for all Turing machines  $\mathcal{M}$  and integers  $\lambda$ , the Turing machine  $\mathcal{D}^{\text{HALT}}$  halts on all inputs. This is because each step of  $\mathcal{F}$  terminates in some finite time. This uses that the procedures ComputeSampler

<sup>35</sup>Alternatively, we can use Kleene's recursion theorem [Kle54] here so that  $\mathcal{D}^{\text{HALT}}$  is the fixed point of a Turing machine similarly defined as  $\mathcal{F}$ . This approach was taken in an earlier version of the paper. We choose to present the proof without resorting to Kleene's recursion theorem for the sake of simplicity.

Input:  $(\overline{\mathcal{R}}, \overline{\mathcal{M}}, \lambda, n, x, y, a, b)$  where  $\overline{\mathcal{R}}$  is the description of an 8-input Turing machine and  $\overline{\mathcal{M}}$  is a description of a Turing machine  $\mathcal{M}$ .

1. Run  $\mathcal{M}$  on the blank input for  $n$  steps. Accept if it halts and continue otherwise.
2. Compute the description  $\overline{\mathcal{D}}$  of the 5-input Turing machine  $\mathcal{D}$  defined as follows:

$$\mathcal{D}(n', x', y', a', b') = \mathcal{R}(\overline{\mathcal{R}}, \overline{\mathcal{M}}, \lambda, n', x', y', a', b')$$

i.e., on input  $(n', x', y', a', b')$  runs  $\mathcal{R}$  with input  $(\overline{\mathcal{R}}, \overline{\mathcal{M}}, \lambda, n', x', y', a', b')$ .

3. Compute the description  $\overline{\mathcal{S}} = \text{ComputeSampler}(\lambda)$ .
4. Compute the description  $\overline{\mathcal{V}^{\text{COMPR}}} = \text{Compress}(\overline{\mathcal{V}}, \lambda)$  where  $\overline{\mathcal{V}} = (\overline{\mathcal{S}}, \overline{\mathcal{D}})$ .
5. Accepts if the decider  $\mathcal{D}^{\text{COMPR}}$  of verifier  $\mathcal{V}^{\text{COMPR}}$  accepts  $(n, x, y, a, b)$ .

Figure 16: Description of Turing machine  $\mathcal{F}$ . The Turing machines `ComputeSampler` and `Compress` are given in Lemma 12.2 and Theorem 12.1.

and `Compress` both terminate in finite time, and the decider  $\mathcal{D}^{\text{COMPR}}$  terminates in finite time (as given by Theorem 12.1), regardless of what input is given to `Compress`. Furthermore, the output of `ComputeSampler` is always a sampler, which means that  $\mathcal{V}^{\text{HALT}} = (\mathcal{S}^{\text{HALT}}, \mathcal{D}^{\text{HALT}})$  is by construction a normal form verifier. We have the following lemma for  $\mathcal{V}^{\text{HALT}}$ .

**Lemma 12.3.** *Let  $\mathcal{M}$  be a Turing machine and  $\lambda$  integer. Then the verifier  $\mathcal{V}^{\text{HALT}}$  corresponding to  $\mathcal{M}$  and  $\lambda$  has the following properties. For all  $n \in \mathbb{N}$ ,*

1. *If  $\mathcal{M}$  halts in  $n$  steps then  $\text{val}^*(\mathcal{V}_n^{\text{HALT}}) = 1$ . In this case, there is a value-1 PCC strategy for the game  $\mathcal{V}_n^{\text{HALT}}$ .*
2. *If  $\mathcal{M}$  does not halt in  $n$  steps then  $\mathcal{V}_n^{\text{HALT}}$  has a value-1 PCC strategy if and only if  $\mathcal{V}_n^{\text{COMPR}}$  does. Furthermore, under the same assumption (that  $\mathcal{M}$  does not halt in  $n$  steps) it holds that*

$$\mathcal{E}(\mathcal{V}_n^{\text{HALT}}, \frac{1}{2}) = \mathcal{E}(\mathcal{V}_n^{\text{COMPR}}, \frac{1}{2}).$$

*Proof.* Suppose  $\mathcal{M}$  halts in  $n$  steps. Then from the definition of  $\mathcal{F}$  and  $\mathcal{D}^{\text{HALT}}$ , it follows that  $\mathcal{D}^{\text{HALT}}$  accepts on input  $(n, x, y, a, b)$  for all  $x, y, a, b$ , and hence  $\text{val}^*(\mathcal{V}_n^{\text{HALT}}) = 1$ . A value-1 PCC strategy for  $\mathcal{V}_n^{\text{HALT}}$  is the following trivial strategy: for all questions, the players always return a fixed answer (e.g. 0).

If  $\mathcal{M}$  does not halt in  $n$  steps, then  $\mathcal{D}^{\text{HALT}}$  accepts on input  $(n, x, y, a, b)$  if and only if  $\mathcal{D}^{\text{COMPR}}$  accepts on  $(n, x, y, a, b)$ . Since  $\mathcal{V}_n^{\text{HALT}}$  and  $\mathcal{V}_n^{\text{COMPR}}$  use the same sampler  $\mathcal{S}^{\text{HALT}} = \text{ComputeSampler}(\lambda)$  by construction, any strategy for  $\mathcal{V}_n^{\text{HALT}}$  is a strategy (with the same winning probability) for  $\mathcal{V}_n^{\text{COMPR}}$  and vice versa. This implies that there is a value-1 PCC strategy for  $\mathcal{V}_n^{\text{HALT}}$  if and only if there is a value-1 PCC strategy for  $\mathcal{V}_n^{\text{COMPR}}$ . The ‘‘Furthermore’’ statement also follows directly from this.  $\square$

We show in Lemma 12.5 below that for all Turing machine  $\mathcal{M}$  there is a choice of  $\lambda$  such that the corresponding  $\mathcal{V}^{\text{HALT}} = (\mathcal{S}^{\text{HALT}}, \mathcal{D}^{\text{HALT}})$  is  $\lambda$ -bounded. We first show a technical lemma.

**Lemma 12.4.** *Suppose  $C, C' \in \mathbb{N}$  are integers. Then for all  $\lambda \geq 4 \max\{(4C)^{8C}, C \log C'\}$  and  $n \geq 2$ ,*

$$C(C' \cdot \lambda n)^C \leq n^\lambda.$$

*Proof.* We aim to find a function  $f(C, C')$  such that for all  $\lambda \geq f(C, C')$ , it holds that  $C(C' \cdot \lambda n)^C \leq n^\lambda$  for all  $n \geq 2$ . For this it suffices that

$$\lambda \geq 4 \max\{\log C, C, C \log C', C \log \lambda\} \quad \text{for all } \lambda \geq f(C, C') \quad (165)$$

This is because (165) implies that  $\lambda \geq \log C + C + C \log C' + C \log \lambda$  which, by exponentiating both sides, gives

$$2^\lambda \geq 2^{\log C + C + C \log C' + C \log \lambda} = C \cdot 2^C \cdot (C' \lambda)^C.$$

Since  $n^{\lambda-C} \geq 2^{\lambda-C}$  for all  $n \geq 2$ , we get that  $n^{\lambda-C} \geq C(C' \lambda)^C$  which is the desired inequality. It is easy to verify that for all  $\lambda \geq (4C)^{8C}$ , we have  $\lambda \geq 4C \log \lambda$ . Thus, setting

$$f(C, C') = 4 \max\{(4C)^{8C}, C \log C'\},$$

we obtain (165), which concludes the lemma.  $\square$

**Lemma 12.5.** *There is an integer parameter  $\lambda$ , polynomial-time computable from the description of the Turing machine  $\mathcal{M}$ , and scaling as  $\text{poly}(|\overline{\mathcal{M}}|)$ , such that the verifier  $\mathcal{V}^{\text{HALT}}$  corresponding to  $\mathcal{M}$  and parameter  $\lambda$  is  $\lambda$ -bounded and moreover  $\text{TIME}_{\mathcal{D}^{\text{HALT}}}(n), \text{TIME}_{\mathcal{S}^{\text{HALT}}}(n) = \text{poly}(n, |\overline{\mathcal{M}}|)$ .*

*Proof.* We first do an accounting of the time complexity of  $\mathcal{D}^{\text{HALT}}$ . By definition, for all  $\mathcal{M}$  and  $\lambda$ , the decider  $\mathcal{D}^{\text{HALT}}$  runs  $\mathcal{F}$  on input  $(\overline{\mathcal{F}}, \overline{\mathcal{M}}, \lambda, n, x, y, a, b)$ . Its running time is therefore a polynomial in the time bounds of the following steps.

1. Simulating  $\mathcal{M}$ : this takes  $\text{poly}(|\overline{\mathcal{M}}|, n)$  time.
2. Computing the description  $\overline{\mathcal{D}}$ : this takes  $\text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$  time.
3. Computing the description  $\overline{\mathcal{S}^{\text{COMPR}}}$ : this takes  $\text{polylog}(\lambda)$  time (by Lemma 12.2).
4. Computing the description  $\overline{\mathcal{D}^{\text{COMPR}}}$ : this takes  $\text{poly}(|\overline{\mathcal{V}}|, \log \lambda)$  time (by Theorem 12.1).
5. Executing  $\mathcal{D}^{\text{COMPR}}$  on input  $(n, x, y, a, b)$ : this takes  $\text{poly}(n, \lambda)$  time (by Theorem 12.1).

As  $\overline{\mathcal{D}}$  is computed in time  $\text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$ , we have that the description length of  $\mathcal{D}$  is bounded by  $|\overline{\mathcal{D}}| \leq \text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$ . Similarly, the description length of the sampler is bounded by  $|\overline{\mathcal{S}}| \leq \text{polylog}(\lambda)$  as  $\text{ComputeSampler}$  runs in polynomial time in the bit-length of  $\lambda$ . Therefore, the size of the verifier  $|\overline{\mathcal{V}}|$  is bounded by  $\text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$ . So the time complexity bound in Item 4 is at most  $\text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$ . We emphasize that we did not use the  $\lambda$ -boundedness of the verifier  $\mathcal{V}$ , as this is what we want to prove here! Overall the total time complexity of  $\mathcal{D}^{\text{HALT}}$  can be bounded by

$$\text{TIME}_{\mathcal{D}^{\text{HALT}}}(n) \leq C_D (|\overline{\mathcal{M}}| \cdot |\overline{\mathcal{F}}| \cdot \lambda \cdot n)^{C_D}, \quad (166)$$

for some universal constant  $C_D > 0$ . By Lemma 12.4, we have for all  $\lambda \geq \lambda_0 := 4 \max\{(4C_D)^{8C_D}, C_D \log(|\overline{\mathcal{M}}| \cdot |\overline{\mathcal{F}}|)\}$  and  $n \geq 2$ , we have  $\text{TIME}_{\mathcal{D}^{\text{HALT}}}(n) \leq n^\lambda$ .

Similarly, by Theorem 12.1 the sampler  $\mathcal{S}^{\text{HALT}}$  has running time  $\text{poly}(n, \lambda)$  and therefore there is a universal constant  $C_S > 0$  such that

$$\text{TIME}_{\mathcal{S}^{\text{HALT}}}(n) \leq C_S(\lambda \cdot n)^{C_S}, \quad (167)$$

which, again by Lemma 12.4, is at most  $n^\lambda$  for  $\lambda \geq \lambda_1 := 4 \cdot (4C_S)^{8C_S}$  and  $n \geq 2$ .

As we have shown  $|\mathcal{V}| = \text{poly}(|\overline{\mathcal{M}}|, |\overline{\mathcal{F}}|, \log \lambda)$  and the description of the verifier  $\mathcal{V}$  is the same as that of  $\mathcal{V}^{\text{HALT}}$ , there exists an integer  $C_V > 0$  such that

$$|\mathcal{V}^{\text{HALT}}| \leq C_V(|\overline{\mathcal{M}}| \cdot |\overline{\mathcal{F}}| \cdot \log \lambda)^{C_V}.$$

Define  $\lambda_2 := (|\overline{\mathcal{M}}| \cdot |\overline{\mathcal{F}}|)^{2C_V} \cdot 2^{2C_V}$ . One can verify that for all  $\lambda \geq \lambda_2$ , we have that  $|\mathcal{V}^{\text{HALT}}| \leq \lambda$ .

Finally, taking  $\lambda = \lceil \max\{\lambda_0, \lambda_1, \lambda_2\} \rceil$  completes the proof. The quantity  $|\overline{\mathcal{F}}|$  is a universal constant, because the Turing machine  $\mathcal{F}$  does not depend on  $\lambda$  or  $\mathcal{M}$ . The quantities  $C_D, C_S, C_V$  are universal constants; therefore  $\lambda_0, \lambda_1, \lambda_2$  and thus  $\lambda$  can be computed from  $|\overline{\mathcal{M}}|$ . It is clear that  $\lambda = \text{poly}(|\overline{\mathcal{M}}|)$ . Finally, for this value of  $\lambda$  one immediately verifies from (166) and (166) that  $\text{TIME}_{\mathcal{D}^{\text{HALT}}}(n), \text{TIME}_{\mathcal{S}^{\text{HALT}}}(n) = \text{poly}(n, |\overline{\mathcal{M}}|)$ , as claimed.  $\square$

Putting things together we obtain the following result.

**Theorem 12.6.** *There exists a polynomial-time computable map from Turing machines  $\mathcal{M}$  to games  $\mathfrak{G}$  that satisfies the “efficiency” requirement from Definition 5.29 and is such that*

1. *If  $\mathcal{M}$  halts on the empty input, then  $\text{val}^*(\mathfrak{G}) = 1$ . Moreover, in this case there is a (finite-dimensional) PCC strategy for the players in  $\mathfrak{G}$  that wins with certainty.*
2. *If  $\mathcal{M}$  does not halt on the empty input, then  $\text{val}^*(\mathfrak{G}) \leq \frac{1}{2}$ .*

*Proof.* The map from Turing machines to games is computed as follows. Given a Turing machine  $\mathcal{M}$ , first compute the parameter  $\lambda$  as given by Lemma 12.5. This takes time at most  $\text{poly}(|\overline{\mathcal{M}}|)$ , and furthermore  $\lambda = \text{poly}(|\overline{\mathcal{M}}|)$ . Then, compute the description of the verifier  $\mathcal{V}^{\text{HALT}} = (\mathcal{S}^{\text{HALT}}, \mathcal{D}^{\text{HALT}})$  corresponding to  $\mathcal{M}$  and  $\lambda$  by first computing the description of  $\mathcal{S}^{\text{HALT}} = \text{ComputeSampler}(\lambda)$  and then computing the description of the Turing machine  $\mathcal{D}^{\text{HALT}}$  that computes

$$\mathcal{D}^{\text{HALT}}(n, x, y, a, b) = \mathcal{F}(\overline{\mathcal{F}}, \overline{\mathcal{M}}, n, x, y, a, b).$$

Computing the description of  $\mathcal{V}^{\text{HALT}}$  thus takes time  $\text{polylog}(\lambda) + \text{poly}(|\overline{\mathcal{M}}|)$  which is  $\text{poly}(|\overline{\mathcal{M}}|)$ . Finally, output the description of the game  $\mathfrak{G} = \mathcal{V}_{C_0}^{\text{HALT}}$  where  $C_0$  is the universal constant given by Theorem 12.1. Clearly this computation takes  $\text{poly}(|\overline{\mathcal{M}}|)$  time in total and, according to the “moreover” part of Lemma 12.5, returns a sampler and decider for  $\mathfrak{G}$  that run in time  $\text{poly}(|\overline{\mathcal{M}}|)$ , as required.

Now fix a Turing machine  $\mathcal{M}$  and let  $\mathfrak{G}_n$  denote the  $n$ -th game  $\mathcal{V}_n^{\text{HALT}}$  of the verifier  $\mathcal{V}^{\text{HALT}}$  computed by the aforementioned map (so in particular the output game  $\mathfrak{G}$  is  $\mathfrak{G}_{C_0}$ ). Suppose that  $\mathcal{M}$  halts on the empty input; let  $T$  be the minimum number of time steps required for  $\mathcal{M}$  to halt on the empty input. Observe that for all  $n \geq T$ , by Lemma 12.3 it holds that  $\mathfrak{G}_n$  has a value-1 PCC strategy. We will use this to show inductively that  $\mathfrak{G}_n$  also has a value-1 PCC strategy, for all  $C_0 \leq n < T$ .

**Claim 12.7.** *Let  $C_0 \leq n < T$ . Suppose that  $\mathfrak{G}_m$  has a value 1 PCC strategy for all  $m > n$ . Then  $\mathfrak{G}_n$  also has a value-1 PCC strategy.*

*Proof.* Since by assumption  $\mathcal{M}$  does not halt in  $n$  steps, by Lemma 12.3 it holds that  $\mathfrak{G}_n$  has a value-1 PCC strategy if  $\mathfrak{G}_n^{\text{COMPR}}$  does. Since  $\mathcal{V}^{\text{HALT}}$  is  $\lambda$ -bounded (by Lemma 12.5) and  $n \geq C_0$ , by Theorem 12.1 it follows that  $\mathfrak{G}_n^{\text{COMPR}}$  has a value-1 PCC strategy if  $\mathfrak{G}_{2^n}$  does. Since  $2^n > n$ , this is true by the hypothesis of the claim. Thus,  $\mathfrak{G}_n$  has a value-1 PCC strategy as claimed.  $\square$

By Claim 12.7 and downwards induction on  $n$  (with the base case  $n = T$ ), we have that  $\mathfrak{G}_n$  has a value-1 PCC strategy for all  $n \geq C_0$ . In particular, we have  $\text{val}^*(\mathfrak{G}) = \text{val}^*(\mathfrak{G}_{C_0}) = 1$ . This shows the first item in the theorem statement.

Now suppose that  $\mathcal{M}$  does not halt on the empty input. We have that for all  $n \in \mathbb{N}$ :

$$\mathcal{E}(\mathfrak{G}_n, \frac{1}{2}) = \mathcal{E}(\mathfrak{G}_n^{\text{COMPR}}, \frac{1}{2}) \geq \mathcal{E}(\mathfrak{G}_N, \frac{1}{2}),$$

where the equality follows from Lemma 12.3 and the inequality follows from Theorem 12.1 (again using the  $\lambda$ -bounded property of  $\mathcal{V}^{\text{HALT}}$ ). By induction, we get that for all  $k \in \mathbb{N}$ ,

$$\mathcal{E}(\mathfrak{G}, \frac{1}{2}) = \mathcal{E}(\mathfrak{G}_{C_0}, \frac{1}{2}) \geq \mathcal{E}(\mathfrak{G}_{g^{(k)}(C_0)}, \frac{1}{2}) = \mathcal{E}(\mathfrak{G}_{g^{(k)}(C_0)}^{\text{COMPR}}, \frac{1}{2}) \geq \frac{1}{2} 2^{(g^{(k+1)}(C_0))^\lambda},$$

where  $g^{(k)}(\cdot)$  is the  $k$ -fold composition of the function  $g(n) = 2^n$  and the second inequality follows from Theorem 12.1 again. Since  $g(\cdot)$  is a strictly monotonically increasing function this implies that there is no finite upper bound on  $\mathcal{E}(\mathfrak{G}, \frac{1}{2})$  and therefore every finite dimensional strategy for the game  $\mathfrak{G}$  must have success probability at most  $\frac{1}{2}$ .  $\square$

Recall the definition of the complexity class RE, which stands for the set of *recursively enumerable* languages (also called *Turing-recognizable* languages). Precisely, a language  $L \subseteq \{0, 1\}^*$  is in RE if and only if there exists a Turing machine  $\mathcal{M}$  such that if  $x \in L$ , then  $\mathcal{M}(x)$  halts and outputs 1, and if  $x \notin L$ , then either  $\mathcal{M}(x)$  outputs 0 or it does not halt. The Halting Problem is the language that contains descriptions of Turing machines that halt on the empty input. The following well-known lemma shows that the Halting Problem is complete for RE, meaning that every language  $L \in \text{RE}$  can be polynomial-time reduced to the Halting Problem. We include the simple proof for convenience.

**Lemma 12.8.** *The Halting Problem is complete for RE via Karp reductions.*<sup>36</sup>

*Proof.* To see that the Halting Problem is in RE, define  $\mathcal{M}$  to take as input an  $x$  that represents a Turing machine  $\mathcal{N} = [x]$ , and runs the universal Turing machine to simulate  $\mathcal{N}$  on the empty input; if  $\mathcal{N}$  halts with a 1 then so does  $\mathcal{M}$ .

To show that the Halting problem is complete for RE, let  $L \in \text{RE}$  and  $\mathcal{M}$  a Turing machine such that if  $x \in L$ , then  $\mathcal{M}(x)$  halts and outputs 1. For an input  $x$ , let  $\mathcal{N}_x$  be the following Turing machine.  $\mathcal{N}_x$  first runs  $\mathcal{M}$  on input  $x$ . If  $\mathcal{M}$  accepts, then  $\mathcal{N}_x$  accepts. On all other outcomes,  $\mathcal{N}_x$  goes into an infinite loop. Thus  $\mathcal{N}_x$  halts if and only if  $x \in L$ .  $\square$

**Corollary 12.9.**  $\text{MIP}^* = \text{RE}$ .

*Proof.* Since the Halting Problem is complete for RE, and by Theorem 12.6 is contained in  $\text{MIP}_{1,1/2}^* \subseteq \text{MIP}^*$  (see Definition 5.29), we have the inclusion  $\text{RE} \subseteq \text{MIP}^*$ . The reverse inclusion  $\text{MIP}^* \subseteq \text{RE}$  follows from the following observation. Let  $L \in \text{MIP}^*$ . From the definition of  $\text{MIP}^*$  (see e.g. [VW16, Section

<sup>36</sup>A Karp reduction between languages  $L$  and  $K$  is a polynomial-time Turing machine  $R$  such that  $x \in L$  if and only if  $R(x) \in K$  for all instances  $x \in \{0, 1\}^*$ .

6.1], from which we borrow the terminology used here) it follows that there exists a polynomial-time Turing machine  $\mathcal{R}$  such that for all  $x \in \{0, 1\}^*$ ,  $\mathcal{R}(x)$  is the description of an  $m$ -turn verifier  $V_x$  interacting with  $k$  provers, where  $m$  and  $k$  are both polynomial functions of  $|x|$  and such that

$$\begin{aligned} \text{val}^*(V_x) &\geq 2/3 && \text{if } x \in L \\ \text{val}^*(V_x) &\leq 1/3 && \text{if } x \notin L \end{aligned}$$

Consider the following Turing machine  $\mathcal{A}$ : on input  $x$ , it computes  $V_x = \mathcal{R}(x)$ , and then exhaustively searches over tensor-product strategies of increasing dimension and increasing accuracy to evaluate a lower bound on  $\text{val}^*(V_x)$ . If  $\text{val}^*(V_x) \geq 2/3$ , then for arbitrarily small  $\delta$  there exists a finite dimensional tensor-product strategy for the players that achieves value  $2/3 - \delta > 1/3$ . When the Turing machine  $\mathcal{A}$  identifies such a strategy it terminates, outputting 1. If there is no such strategy, then  $\mathcal{A}$  never halts. This implies that  $L \in \text{RE}$ .  $\square$

### 12.3 An explicit separation

As discussed in Section 1.3, Theorem 12.6 implies that  $C_{qa}$ , the set of approximately finite-dimensional correlations, is a strict subset of  $C_{qc}$ , the set of commuting-operator correlations. This is because if  $C_{qa} = C_{qc}$ , then there exists an algorithm to approximate the entangled value of a given nonlocal game  $\mathfrak{G}$  to arbitrary accuracy. On the other hand, Theorem 12.6 shows that deciding whether a game has entangled value 1 or at most  $1/2$ , promised that one is the case, is undecidable. Therefore the correlation sets must be different.

In fact, Theorem 12.6 implies that there is an infinite family  $\mathcal{M}$  of Turing machines that do not halt on an empty input such that for all  $\mathcal{M} \in \mathcal{M}$ , the corresponding game  $\mathfrak{G}_{\mathcal{M}}$  has  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) < \text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}})$ , where recall that  $\text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}})$  denotes the *commuting-operator value* of  $\mathfrak{G}_{\mathcal{M}}$ , which is the supremum of success probabilities over all commuting-operator strategies for  $\mathfrak{G}_{\mathcal{M}}$ .<sup>37</sup> However, it is not immediately clear, given a *specific* non-halting Turing machine  $\mathcal{M}$ , whether the associated game  $\mathfrak{G}_{\mathcal{M}}$  separates the commuting-operator model from the tensor product model of strategies. While Theorem 12.6 implies that  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) \leq \frac{1}{2}$ , it could also be the case that  $\text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}}) = \text{val}^*(\mathfrak{G}_{\mathcal{M}})$  in that particular instance. We conjecture that  $\text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}}) = 1$  for *all* non-halting Turing machines  $\mathcal{M}$ , but it appears to be difficult to identify an explicit value-1 commuting operator strategy that demonstrates this.

In this section we identify an explicit game  $\mathfrak{G}$  that separates the tensor product model from the commuting-operator model; we show that  $\text{val}^*(\mathfrak{G}) \leq \frac{1}{2}$  but  $\text{val}^{\text{co}}(\mathfrak{G}) = 1$ . Interestingly, the proof does not exhibit an explicit value-1 commuting-operator strategy for  $\mathfrak{G}$ .

We construct the separating game in a similar manner to the games constructed in Section 12.2. Let  $\mathcal{A}$  denote the following 1-input Turing machine: it takes as input a description of a nonlocal game  $\mathfrak{G}$  and runs the semidefinite programming hierarchy of [NPA08, DLTW08] to compute a non-increasing sequence  $\alpha_1, \alpha_2, \dots$  of upper bounds on  $\text{val}^{\text{co}}(\mathfrak{G})$  such that  $\lim_{n \rightarrow \infty} \alpha_n = \text{val}^{\text{co}}(\mathfrak{G})$ . The Turing machine  $\mathcal{A}$  halts if it obtains a bound  $\alpha_n < 1$ . Thus this algorithm eventually halts whenever  $\text{val}^{\text{co}}(\mathfrak{G}) < 1$ , and otherwise it runs forever.

Consider the Turing machine  $\mathcal{N}$  in Figure 17. We follow the same steps as in Section 12.2. Let  $\mathcal{D}_{\lambda}^{\text{SEP}}$  be the decider that on input  $(n, x, y, a, b)$ , runs  $\mathcal{N}$  on input  $(\overline{\mathcal{N}}, \lambda, n, x, y, a, b)$ . We sometimes leave the

---

<sup>37</sup>To see why this holds, observe that if it were the case that  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) = \text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}})$  for all but finitely many non-halting  $\mathcal{M}$  then we could construct an algorithm  $\mathcal{A}$  to decide the Halting problem as follows. On input  $\mathcal{M}$ ,  $\mathcal{A}$  first checks if  $\mathcal{M}$  is one of the finitely many Turing machines for which  $\text{val}^*(\mathfrak{G}_{\mathcal{M}}) < \text{val}^{\text{co}}(\mathfrak{G}_{\mathcal{M}})$ ; if so, then it outputs a hard-coded answer for whether  $\mathcal{M}$  halts on the empty tape or not. Otherwise,  $\mathcal{A}$  computes the nonlocal game  $\mathfrak{G}_{\mathcal{M}}$  and executes the aforementioned algorithm for approximating the entangled value of games assuming that  $C_{qa} = C_{qc}$ .

parameter  $\lambda$  implicit. Define  $\mathcal{S}^{\text{SEP}} = \text{ComputeSampler}(\lambda)$  and  $\mathcal{V}^{\text{SEP}} = (\mathcal{S}^{\text{SEP}}, \mathcal{D}^{\text{SEP}})$ . By the definition of Turing machine  $\mathcal{N}$  in Fig. 17, the description  $\overline{\mathcal{V}}$  computed by  $\mathcal{D}^{\text{SEP}}$  is the description of  $\mathcal{V}^{\text{SEP}}$  itself and the input to  $\text{Compress}$  is the description of  $\mathcal{V}^{\text{SEP}}$  and  $\lambda$ . Define the game  $\mathfrak{G}^{\text{SEP}} = \mathcal{V}_{C_0}^{\text{SEP}}$  where  $C_0$  is the constant from Theorem 12.1. Then  $\mathfrak{G}^{\text{SEP}}$  is the game  $\mathfrak{G}$  computed in  $\mathcal{D}^{\text{SEP}}$ .

A similar argument to that of Lemma 12.5 shows that there exist choices of  $\lambda$  (computable from the description of  $\mathcal{A}$ ) such that  $\mathcal{V}^{\text{SEP}}$  is  $\lambda$ -bounded.

Input:  $(\overline{\mathcal{R}}, \lambda, n, x, y, a, b)$  where  $\overline{\mathcal{R}}$  is the description of a 7-input Turing machine.

1. Compute the description  $\overline{\mathcal{S}} = \text{ComputeSampler}(\lambda)$ .
2. Compute the description  $\overline{\mathcal{D}}$  of the 5-input Turing machine  $\mathcal{D}$  that on input  $(n', x', y', a', b')$  runs  $\mathcal{R}$  with input  $(\overline{\mathcal{R}}, \lambda, n', x', y', a', b')$ . Let  $\overline{\mathcal{V}} = (\overline{\mathcal{S}}, \overline{\mathcal{D}})$ .
3. Compute the description of the game  $\mathfrak{G} = \mathcal{V}_{C_0}$  where  $C_0$  is the integer from Theorem 12.1.
4. Simulate  $\mathcal{A}$  on input  $\mathfrak{G}$  for  $n$  steps and accept if  $\mathcal{A}$  halts.
5. Compute the description  $\overline{\mathcal{V}^{\text{COMPR}}} = \text{Compress}(\overline{\mathcal{V}}, \lambda)$ .
6. Accept if the decider  $\mathcal{D}^{\text{COMPR}}$  of verifier  $\mathcal{V}^{\text{COMPR}}$  accepts  $(n, x, y, a, b)$ .

Figure 17: Description of Turing machine  $\mathcal{N}$ . The Turing machines  $\text{ComputeSampler}$  and  $\text{Compress}$  are given in Lemma 12.2 and Theorem 12.1.

**Theorem 12.10.** *For the game  $\mathfrak{G}^{\text{SEP}} = \mathcal{V}_{C_0}^{\text{SEP}}$  it holds that  $\text{val}^*(\mathfrak{G}^{\text{SEP}}) \leq \frac{1}{2}$  and  $\text{val}^{\text{co}}(\mathfrak{G}^{\text{SEP}}) = 1$ .*

*Proof.* Suppose that  $\text{val}^{\text{co}}(\mathfrak{G}^{\text{SEP}}) = 1$ . The invocation of the Turing machine  $\mathcal{A}$  on input  $\mathfrak{G}^{\text{SEP}}$  never halts, and therefore the decider  $\mathcal{D}^{\text{SEP}}$  never accepts in Step 4 of Figure 17. Applying Theorem 12.1, we get that  $\mathcal{E}(\mathcal{V}_n^{\text{SEP}}, \frac{1}{2}) \geq \mathcal{E}(\mathcal{V}_{2^n}^{\text{SEP}}, \frac{1}{2})$  and

$$\mathcal{E}\left(\mathcal{V}_n^{\text{SEP}}, \frac{1}{2}\right) \geq 2^{2^n - 1}$$

for all  $n \geq C_0$ . An inductive argument implies that there is no finite upper bound on  $\mathcal{E}(\mathcal{V}_n^{\text{SEP}}, \frac{1}{2})$ , and thus  $\text{val}^*(\mathfrak{G}^{\text{SEP}}) = \text{val}^*(\mathcal{V}_{C_0}^{\text{SEP}}) \leq \frac{1}{2}$ , which implies the theorem.

On the other hand, suppose that  $\text{val}^{\text{co}}(\mathfrak{G}^{\text{SEP}}) < 1$ . Then there exists some  $m \geq C_0$  such that  $\mathcal{A}$  halts on input  $\mathfrak{G}^{\text{SEP}}$  after  $m$  steps, so  $\mathcal{V}_n^{\text{SEP}}$  has a value-1 PCC strategy for all  $n \geq m$  (i.e. the players do not respond with any answers). Thus by the completeness statement of Theorem 12.1 and an induction argument analogous to the one in the proof of Theorem 12.6, we have that  $\mathcal{V}_n^{\text{SEP}}$  has a value-1 PCC strategy for all  $n \geq C_0$ , which implies that  $\text{val}^*(\mathfrak{G}^{\text{SEP}}) = 1$ , a contradiction because of  $\text{val}^*(\mathfrak{G}^{\text{SEP}}) \leq \text{val}^{\text{co}}(\mathfrak{G}^{\text{SEP}})$ . This completes the theorem.  $\square$

## A Analysis of the Pauli basis test

In this appendix we give a proof of Theorem 7.14. For convenience, we repeat the statement of the theorem here.

**Theorem A.1.** *There exists a function*

$$\delta_{\text{QLD}}(\varepsilon, m, d, q) = a(md)^a(\varepsilon^b + q^{-b} + 2^{-bmd})$$

for universal constants  $a \geq 1$  and  $0 < b < 1$  such that the following holds. For all admissible parameter tuples  $\text{qldparams} = (q, m, d)$  and for all strategies  $\mathcal{S} = (|\psi\rangle, A, B)$  for the game  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  that succeed with probability at least  $1 - \varepsilon$ , there exist local isometries  $\phi_A : \mathcal{H}_A \rightarrow \mathcal{H}_{A'} \otimes \mathcal{H}_{A''}$ ,  $\phi_B : \mathcal{H}_B \rightarrow \mathcal{H}_{B'} \otimes \mathcal{H}_{B''}$  (where  $|\psi\rangle \in \mathcal{H}_A \otimes \mathcal{H}_B$  and  $\mathcal{H}_{A''}, \mathcal{H}_{B''} \cong (\mathbb{C}^q)^{\otimes M}$  with  $M = 2^m$ ) and a state  $|\text{AUX}\rangle \in \mathcal{H}_{A'} \otimes \mathcal{H}_{B'}$  such that

1.  $\|\phi_A \otimes \phi_B |\psi\rangle - |\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}\| \leq \delta_{\text{QLD}}(\varepsilon, m, d, q)$ ,
2. Letting  $\tilde{A}_a^x = \phi_A A_a^x \phi_A^\dagger$  and  $\tilde{B}_b^y = \phi_B B_b^y \phi_B^\dagger$ , we have for  $W \in \{X, Z\}$

$$\begin{aligned} \tilde{A}_u^{(\text{PAULI}, W)} \otimes I_{B'B''} &\approx_{\delta_{\text{QLD}}} (\tau_u^W)_{A''} \otimes I_{A'B'B''} \\ I_{A'A''} \otimes \tilde{B}_u^{(\text{PAULI}, W)} &\approx_{\delta_{\text{QLD}}} I_{A'A''B'} \otimes (\tau_u^W)_{B''}, \end{aligned}$$

where the  $\approx_{\delta_{\text{QLD}}}$  statement holds with respect to the state  $|\text{AUX}\rangle_{A'B'} \otimes |\text{EPR}_q\rangle_{A''B''}^{\otimes M}$  and the answer summation is over  $u \in \mathbb{F}_q^M$ .

We also recall the meaning of the parameters:  $q$  is the field size,  $m$  the number of variables,  $d$  the degree of the low-degree code, and  $M = 2^m$  is the number of  $|\text{EPR}_q\rangle$  states being tested for.

This appendix is structured as follows. First, we establish some preliminary facts and definitions in Appendix A.1. In the remainder of the appendix, we show Theorem A.1 by showing how to construct a representation of the Pauli group on the provers' Hilbert space, starting from a strategy  $\mathcal{S}$  that succeeds in  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  with high probability. This is done in several stages: in Appendix A.2, we use the strategy measurements to define a set of observables that correspond to a subset of the Pauli  $X$  and  $Z$  observables, and show that these approximately satisfy the associated Pauli group relations. In Appendix A.3, we adjoin ancillary registers containing EPR states to the provers' state  $|\psi\rangle$ , and define a new set of approximately-commuting observables acting on the expanded state. In Appendix A.4, we combine the approximately-commuting observables into a strategy for the *classical* low-degree test, and in Appendix A.5, we apply the soundness theorem of the classical low-degree test to construct a single projective measurement that simultaneously measures all of the approximately-commuting observables. Finally, in Appendix A.6, we use this single simultaneous measurement to construct an exact representation of the Pauli group, show that it is close to the original strategy observables on the expanded state, and deduce that the game  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  is a self-test for the strategy  $\mathcal{S}^{\text{PAULI}}$ .

### A.1 Preliminaries

**Definition A.2.** Given integer  $d, m$  and a prime power  $q$  we denote by  $\text{ideg}_{d,m}(\mathbb{F}_q)$  the set of polynomials in  $m$  variables with *individual* degree at most  $d$  in each variable. When the field is clear from context we write  $\text{ideg}_{d,m}$ . When the number of variables is clear from context we write  $\text{ideg}_d$ . We denote by  $\text{deg}_d$  the set of polynomials with *total* degree at most  $d$ .



Recall the low-degree code defined in Section 3.4, in which vectors  $h \in \mathbb{F}_q^M$  get mapped to polynomials  $g_h \in \text{ideg}_{\mathcal{G}_{d,m}}(\mathbb{F}_q)$  such that for every  $u \in \{0,1\}^m$ ,  $g_h(u) = h_u$  where the coordinates of  $h$  are indexed by binary strings of length  $m$ . For  $x \in \mathbb{F}_q^m$ , recall the definition of the vector  $\text{ind}_m(x) \in \mathbb{F}_q^M$  from Section 3.4. Recall also the finite field trace  $\text{tr}(\cdot) = \text{tr}_{q \rightarrow 2}(\cdot) : \mathbb{F}_q \rightarrow \mathbb{F}_2$ , where  $q = 2^k$  for odd  $k$ , defined in Section 3.3.1.

Call a tuple  $\omega = (u_X, u_Z, r_X, r_Z) \in (\mathbb{F}_q^m)^2 \times (\mathbb{F}_q)^2$  *anticommuting* if the quantity

$$\gamma = \text{tr}((r_X \cdot \text{ind}_m(u_X)) \cdot (r_Z \cdot \text{ind}_m(u_Z)))$$

introduced in Equation (54) is not 0, and *commuting* if  $\gamma = 0$ .

**Fact A.3.** *The probability that a tuple  $\omega = (u_X, u_Z, r_X, r_Z)$  sampled uniformly at random from  $(\mathbb{F}_q^m)^2 \times (\mathbb{F}_q)^2$  is anti-commuting is at least*

$$(1 - q^{-m}) \cdot (1 - q^{-1}) \cdot \left(1 - \frac{md}{q}\right) \cdot \frac{1}{2} \geq \left(1 - \frac{3md}{q}\right) \cdot \frac{1}{2}.$$

*The probability that  $\omega$  is commuting is also at least  $\left(1 - \frac{3md}{q}\right) \cdot \frac{1}{2}$ .*

*Proof.* First, note that  $\text{ind}_m(u_X) = 0$  if and only if  $u_X = 0$ . If  $u_X = 0$  then  $\omega$  is commuting. Otherwise, if  $u_X \neq 0$ , then observe that for all  $r_X, r_Z, u_Z$ ,

$$\begin{aligned} \text{tr}((r_X \cdot \text{ind}_m(u_X)) \cdot (r_Z \cdot \text{ind}_m(u_Z))) &= \text{tr}(r_X r_Z (\text{ind}_m(u_X) \cdot \text{ind}_m(u_Z))) \\ &= \text{tr}((r_X r_Z) \cdot g_{\text{ind}_m(u_X)}(u_Z)), \end{aligned}$$

where  $g_{\text{ind}_m(u_X)}$  is a nonzero polynomial in  $\text{ideg}_{\mathcal{G}_{d,m}}(\mathbb{F}_q)$ .

If  $r_X = 0$ , then  $\omega$  is commuting. Condition on  $r_X \neq 0$ . Since  $g_{\text{ind}_m(u_X)} \neq 0$  has total degree at most  $md$ , by the Schwartz-Zippel lemma the probability that  $g_{\text{ind}_m(u_X)}(u_Z)$  is zero over a uniformly random choice of  $u_Z$  is at most  $md/q$ . Conditioned on  $g_{\text{ind}_m(u_X)}(u_Z) \neq 0$ , the product  $(r_X r_Z) \cdot g_{\text{ind}_m(u_X)}(u_Z)$  is a uniformly random element of  $\mathbb{F}_q$  when  $r_Z$  is chosen uniformly at random.

The probability that  $u_X \neq 0, r_X \neq 0, g_{\text{ind}_m(u_X)}(u_Z) \neq 0$  is at least  $(1 - q^{-m}) \cdot (1 - q^{-1}) \cdot (1 - md/q)$  since  $u_X, r_X, u_Z$  are chosen independently.

Since  $q = 2^t$  is an admissible field size, we have that for all  $a \in \mathbb{F}_q$ , the trace of  $a$  is equal to  $\sum_i a_i$  where  $(a_1, \dots, a_t) = \kappa(a) \in \mathbb{F}_2^t$  are defined in Section 3.3.1. Thus the probability that a uniformly random element of  $\mathbb{F}_q$  has trace 0 is exactly  $1/2$ .  $\square$

In this appendix we often use the “ $\approx_\delta$ ” notation to specify closeness of general operators (not necessarily POVMs) that are indexed by questions but not answers: given sets of operators  $\{A^x\}$  and  $\{B^x\}$  indexed by questions  $x \in \mathcal{X}$ , we write  $A^x \approx_\delta B^x$  to denote

$$\mathbb{E}_{x \sim \mu} \langle \psi | (A^x - B^x)^\dagger (A^x - B^x) | \psi \rangle \leq O(\delta)$$

where  $\mu$  is a question distribution and  $|\psi\rangle$  is a state that has been fixed by context. This will be useful for expressing closeness of observables, for example. The following two lemmas relate the closeness between sets of operators and weighted sums of those operators.

**Lemma A.4.** *Let  $\{A^x\}$  and  $\{B^x\}$  be operators indexed by some finite set  $\mathcal{X}$ , and let  $\mu$  be a probability distribution over  $\mathcal{X}$ . Let  $\{\alpha_x\}$  be a set of complex numbers such that  $|\alpha_x| \leq 1$ . Then if  $A^x \approx_\delta B^x$  on average over  $x$  drawn from  $\mu$ , then  $A \approx_\delta B$ , where  $A = \mathbb{E}_x \alpha_x A^x$  and  $B = \mathbb{E}_x \alpha_x B^x$ .*

*Proof.* Expand

$$\begin{aligned} \left\| \mathbb{E}_x \alpha_x (A^x - B^x) |\psi\rangle \right\|^2 &\leq \left( \mathbb{E}_x \|(A^x - B^x)|\psi\rangle\| \right)^2 \\ &\leq \mathbb{E}_x \|(A^x - B^x)|\psi\rangle\|^2 \\ &\leq \delta. \end{aligned}$$

The first inequality is the triangle inequality and the second inequality is Jensen's inequality.  $\square$

**Lemma A.5.** *Let  $\mathcal{A}$  be a finite set. Let  $\{A_a^x\}_{a \in \mathcal{A}}$  and  $\{B_a^x\}_{a \in \mathcal{A}}$  be POVMs and let  $\{\alpha_a\}_{a \in \mathcal{A}}$  be a collection of complex numbers on the unit circle. Let  $A^x = \sum_a \alpha_a A_a^x$  and  $B^x = \sum_a \alpha_a B_a^x$ . Then*

$$A_a^x \approx_\delta B_a^x \quad \text{implies} \quad A^x \approx_{\delta'} B^x$$

for  $\delta' = |\mathcal{A}| \cdot \delta$ .

*Proof.* Expand

$$\begin{aligned} \mathbb{E}_x \|(A^x - B^x)|\psi\rangle\|^2 &= \mathbb{E}_x \left\| \sum_a \alpha_a (A_a^x - B_a^x) |\psi\rangle \right\|^2 \\ &\leq \mathbb{E}_x \left( \sum_a \|(A_a^x - B_a^x)|\psi\rangle\| \right)^2 \\ &\leq \mathbb{E}_x |\mathcal{A}| \cdot \sum_a \|(A_a^x - B_a^x)|\psi\rangle\|^2 \\ &\leq |\mathcal{A}| \cdot \delta. \end{aligned}$$

The first inequality follows from the triangle inequality, the second inequality follows from Cauchy-Schwarz, and the last inequality follows from the assumption that  $A_a^x \approx_\delta B_a^x$ .  $\square$

The following lemma shows that POVMs that are approximately projective (in the sense of being self-consistent) are close to exact projective measurements. The lemma first appears in [KV11]; for a self-contained proof, see [JNV<sup>+</sup>20].

**Lemma A.6** (Orthonormalization lemma). *Let  $0 \leq \delta \leq 1$ . There exists a function  $\eta_{\text{ortho}}(\delta) = O(\delta^{1/4})$  such that the following holds. Let  $|\psi\rangle$  be a state on  $\mathcal{H}_A \otimes \mathcal{H}_B$  where  $\mathcal{H}_A, \mathcal{H}_B$  are finite dimensional. Let  $\{Q_a\}$  and  $\{R_a\}$  be POVM on  $\mathcal{H}_A$  and  $\mathcal{H}_B$  respectively such that*

$$Q_a \otimes I_B \simeq_\delta I_A \otimes R_a. \quad (168)$$

*Then there exists a projective measurement  $\{P_a\}$  such that*

$$P_a \otimes I_B \approx_{\eta_{\text{ortho}}} Q_a \otimes I_B. \quad (169)$$

## A.2 Strategies

Let  $\text{qldparams} = (q, m, d)$  be an admissible parameter tuple (Definition 7.12) and let  $\mathcal{S} = (|\psi\rangle, M)$  be a strategy for  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  that succeeds with probability at least  $1 - \varepsilon$ , where  $|\psi\rangle$  is a bipartite state on registers A and B, with associated Hilbert spaces  $\mathcal{H}_A$  and  $\mathcal{H}_B$  respectively. For notational convenience we use  $M$  to

denote the operators for both players; it will be clear from context which player's operators we are referring to (for example, we write  $M_a^{(\text{POINT},W),u} \otimes I_B$  to indicate that  $M_a^{(\text{POINT},W),u}$  is viewed as an operator acting on register A).

Using Naimark's theorem (as formulated in [JNV<sup>+</sup>20, Theorem 5.1]), at the cost of extending the state  $|\psi\rangle$  by adding sufficiently many qubits initialized in the  $|0\rangle$  state we may assume that the measurements in  $\mathcal{S}$  are projective. Let

$$|\psi'\rangle = (|\psi\rangle \otimes |0 \cdots 0\rangle \otimes |0 \cdots 0\rangle)_{AB}$$

be the extended state. In the remainder of this section, we will work with the projective strategy on the extended state, which we continue labeling  $|\psi\rangle$  (instead of  $|\psi'\rangle$ ) for notational convenience.

We introduce several additional notational shorthands. By definition the strategy  $\mathcal{S}$  contains measurement operators for each of the possible question types and content summarized in Fig. 8, such as  $\{M_a^{(\text{POINT},W),y}\}_{a \in \mathbb{F}_q}$  for all  $y \in \mathbb{F}_q^m$ , etc. For "line" questions (of type ALINE or DLINE) we will use LINE as a formal placeholder for either type ALINE or DLINE; which type is meant will be clear from the nature of the line. Moreover, we will use  $\ell$  to denote the description of either an axis-parallel line (given by a pair  $\ell = (u_0, s)$ ) or a diagonal line (given by a triple  $\ell = (u_0, s, v)$ ). We also introduce the following collection of observables constructed from the measurement operators in the strategy. For every  $u \in \mathbb{F}_q^m$ ,  $r \in \mathbb{F}_q$ , and  $W \in \{X, Z\}$ , define

$$W^r(u) = \sum_{a \in \mathbb{F}_q} (-1)^{\text{tr}(ar)} M_a^{(\text{POINT},W),u}. \quad (170)$$

The observable  $W^r(u)$  acts on register A or B depending on whether  $M_a^{(\text{POINT},W),u}$  is viewed as being an operator acting on register A or B; this will be made clear from context.

Recall the definitions of (anti-)commuting tuple given at the start of Section A.1.

**Lemma A.7.** *Assume that  $\mathcal{S}$  succeeds in  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  with probability at least  $1 - \varepsilon$ , for some  $\varepsilon \geq 0$ . Then the following hold:*

1. **(Consistency check)** *On average over  $(t, x)$  sampled from the question distribution of  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$ , we have that  $M_a^{t,x} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{t,x}$ .*
2. **(Low-degree check)** *For all  $W \in \{X, Z\}$ , on average over  $(\ell, u)$  drawn from the line-point distribution (see Definition 7.6), we have  $M_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{(\text{POINT},W),u}$ , where the answer summation is over  $a \in \mathbb{F}_q$ .*
3. **(Pauli basis consistency check)** *For all  $W \in \{X, Z\}$ , on average over  $u \in \mathbb{F}_q^m$ , we have  $M_a^{(\text{POINT},W),u} \otimes I_B \simeq_\varepsilon I_A \otimes M_{[g_h(u)=a]}^{(\text{PAULI},W)}$  where  $M_{[g_h(u)=a]}^{(\text{PAULI},W)} = \sum_{h \in \mathbb{F}_q^m: g_h(u)=a} M_h^{(\text{PAULI},W)}$  and  $g_h$  is the low-degree encoding of  $h$ .*
4. **(Commutation check)** *For all  $W \in \{X, Z\}$ , on average over commuting  $\omega = (u_X, u_Z, r_X, r_Z)$ , we have  $M_a^{(\text{PAIR},W),\omega} \otimes I_B \simeq_\varepsilon I_A \otimes M_{[\beta_W=a]}^{\text{PAIR},\omega}$ , where the answer summation is over  $a \in \mathbb{F}_2$ .*
5. **(Commutation consistency check)** *For all  $W \in \{X, Z\}$ , on average over commuting  $\omega = (u_X, u_Z, r_X, r_Z)$ , we have  $M_{[\text{tr}(\cdot r_W)=a]}^{(\text{POINT},W),u_W} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{(\text{PAIR},W),\omega}$ , where the answer summation is over  $a \in \mathbb{F}_2$ .*

6. (**Magic square check**) For any anticommuting  $\omega = (u_X, u_Z, r_X, r_Z)$  let  $\Lambda_\omega$  be the value of the strategy  $(|\psi\rangle, \{M_{\alpha_1, \alpha_2, \alpha_3}^{\text{CONSTRAINT}_i, \omega}\} \cup \{M_a^{\text{VARIABLE}_j, \omega}\})$  in the game  $\mathfrak{G}^{\text{MS}}$ . Then

$$\mathbb{E}_{\omega: \text{tr}((r_X \cdot u_X) \cdot (r_Z \cdot u_Z)) \neq 0} \Lambda_\omega = 1 - O(\varepsilon).$$

7. (**Magic square consistency check**) For all  $W \in \{X, Z\}$ , on average over anticommuting  $\omega = (u_X, u_Z, r_X, r_Z)$ ,

$$M_{[\text{tr}(\cdot r_X)=a]}^{(\text{POINT}, X), u_X} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{\text{VARIABLE}_1, \omega}, \quad (171)$$

$$M_{[\text{tr}(\cdot r_Z)=a]}^{(\text{POINT}, Z), u_Z} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{\text{VARIABLE}_5, \omega}, \quad (172)$$

where the answer summations are over  $\mathbb{F}_2$ .

Furthermore, all of the approximations above hold with “ $\simeq_\varepsilon$ ” instead of “ $\simeq_\varepsilon$ ”, and they also hold with the tensor factors interchanged.

*Proof.* Since the question types are sampled according to the type graph  $G^{\text{PAULI}}$  of finite size in Fig. 7 in the game  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$ , the probability of any of the subtests of  $\mathcal{D}_{\text{qldparams}}^{\text{PAULI}}$  (described in Fig. 8) being executed is at least some universal constant. Thus the consistency conditions expressed in Items 1, 2, 3 of the lemma follow directly from the corresponding tests in Fig. 8 (where Item 2 follows from the consistency test that is a subtest of  $\mathcal{D}_{\text{ldparams}}^{\text{LD}}$ ) and the fact that the strategy  $\mathcal{S}$  is assumed to succeed with probability at least  $1 - \varepsilon$  in  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$ .

For the remaining items, recall that by Fact A.3 the probability that a tuple  $\omega = (u_X, u_Z, r_X, r_Z)$  sampled as question content for a question of type PAIR, (PAIR,  $W$ ), CONSTRAINT $_i$  or VARIABLE $_j$  is commuting is at least  $(1 - 3md/q) \cdot (1/2)$ , which is at least  $1/4$  assuming  $6md \leq q$ . We claim this assumption holds without loss of generality. Indeed, observe that the function  $\delta$  in Theorem A.1 is monotonically increasing in  $a$ , so without loss of generality we may assume that  $a \geq 6$ . Then it follows that we may assume that  $6md \leq q$  without loss of generality, since otherwise,  $\delta \geq amd/q \geq 1$  and the conclusion of the theorem holds trivially. The same lower bound holds for the probability that  $\omega$  is anticommuting. Therefore, the consistency conditions in Items 4, 5 and 7 of the lemma also follow directly from the corresponding tests in Fig. 8. Finally, Item 6 in the lemma follows from the fact that the Magic Square check in Fig. 8 is executed with constant probability over the choice of a pair of questions.

That all of the approximations in the Lemma statement also hold with  $\simeq_\varepsilon$  replaced by  $\approx_\varepsilon$  is immediate from Fact 5.18.  $\square$

**Lemma A.8.** *The following hold for the observables defined in (170).*

- For all  $W \in \{X, Z\}$ ,  $r \in \mathbb{F}_q$ , on average over uniformly random  $u \in \mathbb{F}_q^m$  we have

$$W^r(u) \otimes I_B \approx_\varepsilon I_A \otimes W^r(u). \quad (173)$$

- On average over uniformly random  $\omega = (u_X, u_Z, r_X, r_Z)$ ,

$$\begin{aligned} X^{r_X}(u_X) Z^{r_Z}(u_Z) \otimes I_B \\ \approx_{\sqrt{\varepsilon}} (-1)^{\text{tr}((r_X \cdot \text{ind}_m(u_X)) \cdot (r_Z \cdot \text{ind}_m(u_Z)))} Z^{r_Z}(u_Z) X^{r_X}(u_X) \otimes I_B. \end{aligned} \quad (174)$$

Furthermore, all approximations above hold with the tensor factors interchanged.

*Proof.* We first establish (173). Fix a  $W \in \{X, Z\}$  and  $r \in \mathbb{F}_q$ . From item 1 of Lemma A.7 and the data processing inequality (Fact 5.24), using that the question type (POINT,  $W$ ) has constant probability of being sampled by  $\mathcal{S}^{\text{PAULI}}$ , we get that on average over  $u \in \mathbb{F}_q$ ,

$$M_{[\text{tr}(\cdot r)=a]}^{(\text{POINT}, W), u} \otimes I_B \approx_\varepsilon I_A \otimes M_{[\text{tr}(\cdot r)=a]}^{(\text{POINT}, W), u},$$

where the answer summation is over  $a \in \mathbb{F}_2$ . Equation (173) follows directly from Lemma A.5.

We now establish (174). We have that on average over commuting  $\omega = (u_X, u_Z, r_X, r_Z)$  each of the approximate consistency relations hold, for any  $W \in \{X, Z\}$ :

$$M_{[\text{tr}(\cdot r_W)=a]}^{(\text{POINT}, W), u_W} \otimes I_B \simeq_\varepsilon I_A \otimes M_a^{(\text{PAIR}, W), \omega} \quad (\text{Item 5 of Lemma A.7})$$

$$\simeq_\varepsilon M_{[\beta_W=a]}^{\text{PAIR}, \omega} \otimes I_B \quad (\text{Item 4})$$

$$\simeq_\varepsilon I_A \otimes M_{[\beta_W=a]}^{\text{PAIR}, \omega}. \quad (\text{Item 1 and Fact 5.24})$$

Fact 5.18 then implies that on average over commuting  $\omega$ ,

$$M_{[\text{tr}(\cdot r_W)=a]}^{(\text{POINT}, W), u_W} \otimes I_B \approx_\varepsilon I_A \otimes M_{[\beta_W=a]}^{\text{PAIR}, \omega}. \quad (175)$$

Since  $\{M_{\beta_X, \beta_Z}^{\text{PAIR}, \omega}\}$  is projective, Equation (175) puts us in a position to apply Lemma 5.25, setting  $A_{a,b}^x = M_{[\text{tr}(\cdot r_X)=b]}^{(\text{POINT}, X), u_X}$ ,  $C_{a,c}^x = M_{[\text{tr}(\cdot r_Z)=c]}^{(\text{POINT}, Z), u_Z}$ , and  $B_{a,b,c}^x = M_{b,c}^{\text{PAIR}, \omega}$ . It follows that

$$M_{[\text{tr}(\cdot r_X)=b]}^{(\text{POINT}, X), u_X} M_{[\text{tr}(\cdot r_Z)=c]}^{(\text{POINT}, Z), u_Z} \otimes I_B \approx_\varepsilon M_{[\text{tr}(\cdot r_Z)=c]}^{(\text{POINT}, Z), u_Z} M_{[\text{tr}(\cdot r_X)=b]}^{(\text{POINT}, X), u_X} \otimes I_B \quad (176)$$

on average over commuting  $\omega$ . Then for every  $\omega = (u_X, u_Z, r_X, r_Z)$ , since the measurements  $\{M_a^{t,x}\}$  are projective we get that

$$\begin{aligned} X^{r_X}(u_X) Z^{r_Z}(u_Z) &= \left( I - 2M_{[\text{tr}(\cdot r_X)=1]}^{(\text{POINT}, X), u_X} \right) \left( I - 2M_{[\text{tr}(\cdot r_Z)=1]}^{(\text{POINT}, Z), u_Z} \right) \\ &= I - 2M_{[\text{tr}(\cdot r_X)=1]}^{(\text{POINT}, X), u_X} - 2M_{[\text{tr}(\cdot r_Z)=1]}^{(\text{POINT}, Z), u_Z} + 4M_{[\text{tr}(\cdot r_X)=1]}^{(\text{POINT}, X), u_X} M_{[\text{tr}(\cdot r_Z)=1]}^{(\text{POINT}, Z), u_Z}. \end{aligned} \quad (177)$$

Applying (176) with  $b = c = 1$  to commute the measurement operators in the the fourth term in (177), and performing the same steps in reverse, we deduce that on average over commuting  $\omega$ ,

$$X^{r_X}(u_X) Z^{r_Z}(u_Z) \approx_\varepsilon Z^{r_Z}(u_Z) X^{r_X}(u_X).$$

This shows the ‘‘commuting’’ part of (174).

Now we consider the case that  $\omega$  is anticommuting. Item 7 of Lemma A.7, combined with Lemma A.5, implies that the following two approximations hold on average over anticommuting  $\omega$ :

$$X^{r_X}(u_X) \otimes I_B \approx_\varepsilon I_A \otimes M^{\text{VARIABLE}_1, \omega}, \quad (178)$$

$$Z^{r_Z}(u_Z) \otimes I_B \approx_\varepsilon I_A \otimes M^{\text{VARIABLE}_5, \omega}, \quad (179)$$

where for  $j \in \{1, 5\}$ ,  $M^{\text{VARIABLE}_j, \omega} = M_0^{\text{VARIABLE}_j, \omega} - M_1^{\text{VARIABLE}_j, \omega}$ .

For all anticommuting  $\omega$ , let  $\varepsilon_\omega = 1 - \Lambda_\omega$ , where  $\Lambda_\omega$  is defined in item 6 of Lemma A.7. From item 6 of Lemma A.7 we get that  $\mathbb{E}_{\omega \text{ anticom.}} \varepsilon_\omega \leq O(\varepsilon)$ . For each anticommuting  $\omega$ , Theorem 7.10 implies that

$$I_A \otimes M^{\text{VARIABLE}_{E_1, \omega}} M^{\text{VARIABLE}_{E_5, \omega}} \approx_{\sqrt{\varepsilon_\omega}} -I_A \otimes M^{\text{VARIABLE}_{E_5, \omega}} M^{\text{VARIABLE}_{E_1, \omega}} .$$

Thus, averaging over anticommuting  $\omega$ , and applying Jensen's inequality to the definition of  $\approx_\varepsilon$ , we obtain the same anticommutation relation with error  $\mathbb{E}_{\omega \text{ anticom.}} \varepsilon_\omega^{1/2} \leq (\mathbb{E}_{\omega \text{ anticom.}} \varepsilon_\omega)^{1/2} = O(\varepsilon^{1/2})$ . In summary, on average over anticommuting  $\omega$ ,

$$I_A \otimes M^{\text{VARIABLE}_{E_1, \omega}} M^{\text{VARIABLE}_{E_5, \omega}} \approx_{\sqrt{\varepsilon}} -I_A \otimes M^{\text{VARIABLE}_{E_5, \omega}} M^{\text{VARIABLE}_{E_1, \omega}} . \quad (180)$$

Using 5.19 twice we get from (178) and (179) that on average over anticommuting  $\omega$ ,

$$X^{rx}(u_X) Z^{rz}(u_Z) \otimes I_B \approx_\varepsilon I_A \otimes M^{\text{VARIABLE}_{E_1, \omega}} M^{\text{VARIABLE}_{E_5, \omega}} , \quad (181)$$

$$Z^{rx}(u_X) X^{rz}(u_Z) \otimes I_B \approx_\varepsilon I_A \otimes M^{\text{VARIABLE}_{E_5, \omega}} M^{\text{VARIABLE}_{E_1, \omega}} . \quad (182)$$

Using (180) we get that on average over anticommuting  $\omega$ , we have

$$X^{rx}(u_X) Z^{rz}(u_Z) \otimes I_B \approx_{\sqrt{\varepsilon}} -Z^{rz}(u_Z) X^{rx}(u_X) \otimes I_B .$$

This shows the ‘‘anticommuting’’ part of (174).  $\square$

### A.3 Expanding the Hilbert space and defining commuting observables

**Expanded state.** We introduce registers  $A', A'', B', B''$  that are each isomorphic to  $(\mathbb{C}^q)^{\otimes M}$ . Define the state

$$|\hat{\psi}\rangle_{AA'A''BB'B''} = (|\psi\rangle)_{AB} \otimes |\text{EPR}_q\rangle_{A'A''}^{\otimes M} \otimes |\text{EPR}_q\rangle_{B'B''}^{\otimes M} , \quad (183)$$

where we have added maximally entangled states in registers  $A'A''$  and  $B'B''$ ; recall also that the state  $|\psi\rangle$  is already implicitly appended with sufficiently many ancilla qubits initialized in the  $|0\rangle$  state. For the remainder of the Appendix all approximations will be evaluated with respect to this state.

**Expanded observables.** For all  $W \in \{X, Z\}$ ,  $r \in \mathbb{F}_q$ , and  $u \in \mathbb{F}_q^m$  define the following observable  $\hat{W}^r(u)$ :

$$\hat{W}^r(u) = W^r(u) \otimes \tau^W(r \cdot \text{ind}_m(u)) , \quad (184)$$

where  $W^r(u)$  is the observable defined in Equation (170) and  $\tau^W(r \cdot \text{ind}_m(u))$  is the generalized Pauli observable (see Section 3.7) acting on  $(\mathbb{C}^q)^{\otimes M}$ .

For  $a \in \mathbb{F}_q$  define corresponding projections

$$\hat{M}_a^{(\text{POINT}, W), u} = \mathbb{E}_{r \in \mathbb{F}_q} (-1)^{\text{tr}(ar)} \hat{W}^r(u) .$$

Note that  $\hat{M}_a^{(\text{POINT}, W), u}$  can be equivalently written as a sum of projectors by applying Fourier identities. Specifically, for  $W \in \{X, Z\}$ ,  $u \in \mathbb{F}_q^m$ , and  $a \in \mathbb{F}_q$ , define the projector

$$\tau_a^{W, u} = \tau_{[g \cdot (u)=a]}^W = \sum_{h \in \mathbb{F}_q^M : h \cdot \text{ind}_m(u) = a} \tau_h^W . \quad (185)$$

Then we can write  $\hat{M}_a^{(\text{POINT},W),u}$  as follows:

$$\begin{aligned}
\hat{M}_a^{(\text{POINT},W),u} &= \mathbb{E}_{r \in \mathbb{F}_q} (-1)^{\text{tr}(ar)} W^r(u) \otimes \tau^W(r \cdot \text{ind}_m(u)) \\
&= \mathbb{E}_{r \in \mathbb{F}_q} (-1)^{\text{tr}(ar)} \left( \sum_{a' \in \mathbb{F}_q} (-1)^{\text{tr}(a'r)} M_{a'}^{(\text{POINT},W),u} \right) \otimes \left( \sum_{h \in \mathbb{F}_q^n} (-1)^{\text{tr}(r(h \cdot \text{ind}_m(u)))} \tau_h^W \right) \\
&= \mathbb{E}_{r \in \mathbb{F}_q} (-1)^{\text{tr}(ar)} \left( \sum_{a' \in \mathbb{F}_q} (-1)^{\text{tr}(a'r)} M_{a'}^{(\text{POINT},W),u} \right) \otimes \left( \sum_{a'' \in \mathbb{F}_q} (-1)^{\text{tr}(a''r)} \tau_{a''}^{W,u} \right) \\
&= \sum_{a', a'' \in \mathbb{F}_q} \mathbb{E}_{r \in \mathbb{F}_q} (-1)^{\text{tr}((a+a'+a'')r)} M_{a'}^{(\text{POINT},W),u} \otimes \tau_{a''}^{W,u} \\
&= \sum_{\substack{a', a'' \in \mathbb{F}_q \\ a'+a''=a}} M_{a'}^{(\text{POINT},W),u} \otimes \tau_{a''}^{W,u},
\end{aligned}$$

where in the second line we have used Equation (170) on the first tensor factor and Equation (20) on the second, and in the third line we have used Equation (185). For a fixed  $u \in \mathbb{F}_q^m$ ,  $r \in \mathbb{F}_q$  and  $b \in \mathbb{F}_2$ , define a projection

$$\hat{M}_b^{(\text{POINT},W),u,r} = \sum_{a \in \mathbb{F}_q: \text{tr}(ar)=b} \hat{M}_a^{(\text{POINT},W),u}, \quad (186)$$

which can be equivalently written as

$$\hat{M}_b^{(\text{POINT},W),u,r} = \frac{1}{2} \left( I + (-1)^b \hat{W}^r(u) \right). \quad (187)$$

The registers that the operators  $\hat{W}^r(u)$ ,  $\hat{M}_a^{(\text{POINT},W),u}$ , and  $\hat{M}_b^{(\text{POINT},W),u,r}$  act on depend on context. For example if  $W^r(u)$  is viewed as an operator acting on register A, then we will view  $\hat{W}^r(u)$  as acting on registers AA' (with the  $\tau^W$  observable acting on register A'). We generally indicate via subscripts which registers the operators act on (for example we will write  $(\hat{W}^r(u))_{AA'}$  or  $(\hat{W}^r(u))_{BA''}$ ).

**Partitioning the registers, and symmetries** In the remainder of this section as well as Appendices A.4 and A.5, we will partition the six registers AA'A''BB'B'' into two ‘‘parties’’ in two different ways. The first is to consider AA' together to form a single party, and BA'' to form the second party. The second is to consider BB' to form the first party, and AB'' to form the second party. Thanks to the symmetry of the Pauli basis test under exchanging the two players, as well as the symmetry of the ancilla states in registers A'A'' and B'B'', every ‘‘bipartite’’ consistency relation we derive between two operators  $M, N$  will hold for both of the two bipartitioning schemes, and with the first and second party interchanged. That is, whenever we derive a bipartite relation of the form

$$M_{AA'} \approx_\delta N_{BA''},$$

the same steps, with the registers appropriately changed, will also yield

$$N_{AA'} \approx_\delta M_{BA''},$$

$$M_{BB'} \approx_\delta N_{AB''},$$

$$N_{BB'} \approx_\delta M_{AB''}.$$

Similarly, whenever we derive a single-party relation of the form  $M_{AA'} \approx N_{AA'}$ , the same steps with the registers appropriately changed will also yield the same relation on BA'', BB' and AB''. We refer to these relations as *symmetric equivalents*.

**Lemma A.9.** *There exists a function  $\delta_{\text{ANTICOM}}(\varepsilon) = \text{poly}(\varepsilon)$  such that the following holds.*

1. (**Self-consistency**) *For all  $W \in \{X, Z\}$ , on average over  $r \in \mathbb{F}_q$  and  $u \in \mathbb{F}_q^m$ , we have*

$$(\hat{M}_a^{(\text{POINT}, W), u})_{AA'} \approx_\varepsilon (\hat{M}_a^{(\text{POINT}, W), u})_{BA''}.$$

2. (**Approximate commutation**) *On average over uniformly random  $\omega = (u_X, u_Z, r_X, r_Z)$ , we have*

$$(\hat{M}_b^{(\text{POINT}, X), u_X, r_X} \hat{M}_{b'}^{(\text{POINT}, Z), u_Z, r_Z})_{AA'} \approx_{\delta_{\text{ANTICOM}}} (\hat{M}_{b'}^{(\text{POINT}, Z), u_Z, r_Z} \hat{M}_b^{(\text{POINT}, X), u_X, r_X})_{AA'}.$$

Furthermore, all symmetric equivalents of these approximations also hold.

*Proof.* Item 1, the self-consistency of  $\{\hat{M}_a^{(\text{POINT}, W), u}\}$ , follows from the fact that the original points measurements  $\{M_a^{(\text{POINT}, W), u}\}$  are approximately self-consistent between registers A and B in  $|\hat{\psi}\rangle$  (item 1 in Lemma A.7), the Pauli projections  $\{\tau_a^W(\text{ind}_m(u))\}$  are perfectly self-consistent between registers A' and A'', and Fact 5.24.

We now establish the approximate commutation relations. On average over uniformly random  $\omega = (u_X, u_Z, r_X, r_Z)$ , we have

$$\begin{aligned} & \hat{M}_b^{(\text{POINT}, X), u_X, r_X} \hat{M}_{b'}^{(\text{POINT}, Z), u_Z, r_Z} \\ &= \frac{1}{4} \left( I + (-1)^b \hat{X}^{r_X}(u_X) \right) \left( I + (-1)^{b'} \hat{Z}^{r_Z}(u_Z) \right) \\ &= \frac{1}{4} \left( I + (-1)^b \hat{X}^{r_X}(u_X) + (-1)^{b'} \hat{Z}^{r_Z}(u_Z) + (-1)^{b+b'} \hat{X}^{r_X}(u_X) \hat{Z}^{r_Z}(u_Z) \right), \end{aligned}$$

where the first equality follows from (187). Note that

$$\begin{aligned} & \hat{X}^{r_X}(u_X) \hat{Z}^{r_Z}(u_Z) \\ &= X^{r_X}(u_X) Z^{r_Z}(u_Z) \otimes \tau^X(r_X \cdot \text{ind}_m(u_X)) \tau^Z(r_Z \cdot \text{ind}_m(u_Z)) \\ &\approx_{\sqrt{\varepsilon}} Z^{r_Z}(u_Z) X^{r_X}(u_X) \otimes \tau^Z(r_Z \cdot \text{ind}_m(u_Z)) \tau^X(r_X \cdot \text{ind}_m(u_X)) \\ &= \hat{Z}^{r_Z}(u_Z) \hat{X}^{r_X}(u_X), \end{aligned}$$

where the first equality follows from (184) and for the second line we used the approximate (anti-)commutation relation (174) for the  $X^{r_X}(u_X)$  and  $Z^{r_Z}(u_Z)$  observables, as well as the exact (anti-)commutation relations for the  $\tau^Z(r_Z \cdot \text{ind}_m(u_Z))$  and  $\tau^X(r_X \cdot \text{ind}_m(u_X))$  observables. This shows the desired approximate commutation relation, with  $\delta_{\text{ANTICOM}} = \sqrt{\varepsilon}$ .  $\square$

**Lemma A.10.** *There exists a function  $\delta_{\text{LINE}}(\varepsilon) = \text{poly}(\varepsilon)$  such that the following holds. For all  $W \in \{X, Z\}$  and lines  $\ell$  there exists a projective measurement  $\{\hat{M}_f^{(\text{LINE}, W), \ell}\}_{f \in \text{deg}_{\text{md}}(\ell)}$ , acting on registers AA' or BA'', such that:*

1. (**Self-consistency**) *For all  $W \in \{X, Z\}$  and  $r \in \mathbb{F}_q$ , on average over a uniformly random pair  $(\ell, u)$  drawn from the line-point distribution (Definition 7.6),*

$$(\hat{M}_f^{(\text{LINE}, W), \ell})_{AA'} \approx_{\delta_{\text{LINE}}} (\hat{M}_f^{(\text{LINE}, W), \ell})_{BA''}.$$



2. (**Consistency with points measurements I**) On average over a uniformly random pair  $(\ell, u)$  drawn from the line-point distribution,

$$(\hat{M}_f^{(\text{LINE}, W), \ell})_{AA'} \approx_{\delta_{\text{LINE}}} (\hat{M}_f^{(\text{LINE}, W), \ell})_{AA'} \otimes (\hat{M}_{f(u)}^{(\text{POINT}, W), u})_{BA''}.$$

3. (**Consistency with points measurements II**) On average over a uniformly random pair  $(\ell, u)$  drawn from the line-point distribution,

$$(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE}, W), \ell})_{AA'} \approx_{\delta_{\text{LINE}}} (\hat{M}_a^{(\text{POINT}, W), u})_{BA''}.$$

Furthermore, all symmetric equivalents of these approximations also hold.

*Proof.* Define the operator

$$\hat{M}_f^{(\text{LINE}, W), \ell} = \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell},$$

where the Pauli operator  $\tau_{f''}^{W, \ell}$  acting on register  $A'$  is obtained by measuring all  $M$  qudits in the basis  $W$  to obtain an outcome  $h \in \mathbb{F}_q^M$  and then returning the restriction  $f''$  of  $g_h : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  to the line  $\ell$ . Here,  $g_h$  is the degree- $d$  low-degree extension of  $h$  defined in (12).

Item 1, the self-consistency of  $\{\hat{M}_f^{(\text{LINE}, W), \ell}\}$ , follows from the fact that the original lines measurement  $\{M_f^{(\text{LINE}, W), \ell}\}$  is self-consistent between registers  $A$  and  $B$  in  $|\hat{\psi}\rangle$  (Item 1 in Lemma A.7), the Pauli projections  $\{\tau_{f''}^{W, \ell}\}$  are perfectly self-consistent on between registers  $A'$  and  $A''$ , and Fact 5.24.

Item 2, the consistency with the points measurements, follows because on average over  $(\ell, u)$  drawn from the line-point distribution we have

$$(\hat{M}_f^{(\text{LINE}, W), \ell})_{AA'} \tag{188}$$

$$= \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} (M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell})_{AA'} \tag{189}$$

$$= \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} ((M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell}) \cdot (M_{[\text{eval}_u(\cdot)=f'(u)]}^{(\text{LINE}, W), \ell}))_{AA'} \tag{190}$$

$$\approx_{\varepsilon} \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} (M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell})_{AA'} \otimes (M_{f'(u)}^{(\text{POINT}, W), u})_{BA''} \tag{191}$$

$$= \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} (M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell})_{AA'} \otimes (M_{f'(u)}^{(\text{POINT}, W), u} \otimes \tau_{f''(u)}^{W, u})_{BA''} \tag{192}$$

$$= \sum_{\substack{f', f'' \in \text{deg}_{\text{md}}(\ell): \\ f' + f'' = f}} (M_{f'}^{(\text{LINE}, W), \ell} \otimes \tau_{f''}^{W, \ell})_{AA'} \otimes \left( \sum_{\substack{a', a'': \\ a' + a'' = f'(u) + f''(u)}} M_{a'}^{(\text{POINT}, W), u} \otimes \tau_{a''}^{W, u} \right)_{BA''} \tag{193}$$

$$= (\hat{M}_f^{(\text{LINE}, W), \ell})_{AA'} \otimes (\hat{M}_{f(u)}^{(\text{POINT}, W), u})_{BA''}. \tag{194}$$

Equation (190) follows from the projectivity of  $\{\hat{M}_f^{(\text{LINE},W),\ell}\}$ . Equation (191) follows from the consistency between the line and point measurements (Item 2 in Lemma A.7) together with Fact 5.20. Equation (192) follows from the exact consistency between the  $\{\tau_{f''}^{W,\ell}\}$  and  $\{\tau_{f''(u)}^{W,u}\}$  measurements on the  $A'$  and  $A''$  registers. Equation (193) follows because the consistency of the  $\tau$  measurements force  $a' = f'(u)$  and  $a'' = f''(u)$ . This establishes Item 2, with approximation error  $\varepsilon$ .

To show Item 3 we would like to first apply Fact 5.24 to Equation (194), to sum over all functions  $f$  that evaluate to the same value at a given point  $u$ . Unfortunately, we cannot use the Fact as written, since the bound in Item 2 is in terms of  $\approx$ , not  $\simeq$ , and moreover the right-hand side does not have the form  $I \otimes B$ . Instead, we show the desired bound by direct calculation from Equation (194)

$$\varepsilon \geq \mathbb{E}_\omega \sum_f \langle \hat{\psi} | (\hat{M}_f^{(\text{LINE},W),\ell})^2 \otimes (I - \hat{M}_{f(u)}^{(\text{POINT},W),u})^2 | \hat{\psi} \rangle \quad (195)$$

$$= \mathbb{E}_\omega \sum_f \langle \hat{\psi} | \hat{M}_f^{(\text{LINE},W),\ell} \otimes (I - \hat{M}_{f(u)}^{(\text{POINT},W),u}) | \hat{\psi} \rangle \quad (196)$$

$$= \mathbb{E}_\omega \sum_a \sum_{f:f(u)=a} \langle \hat{\psi} | \hat{M}_f^{(\text{LINE},W),\ell} \otimes (I - \hat{M}_a^{(\text{POINT},W),u}) | \hat{\psi} \rangle \quad (197)$$

$$= \mathbb{E}_\omega \sum_a \langle \hat{\psi} | \hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell} \otimes (I - \hat{M}_a^{(\text{POINT},W),u}) | \hat{\psi} \rangle, \quad (198)$$

where in going to the second line we have used the projectivity of the measurements  $\{\hat{M}_f^{(\text{LINE},W),\ell}\}$  and  $\{\hat{M}_a^{(\text{POINT},W),u}\}$ . Thus, we have deduced that

$$(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \approx_\varepsilon (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes (\hat{M}_a^{(\text{POINT},W),u})_{BA''}. \quad (199)$$

Since both the  $\{\hat{M}_f^{(\text{LINE},W),\ell}\}$  and  $\{\hat{M}_a^{(\text{POINT},W),u}\}$  measurements are projective, we also get

$$\begin{aligned} I &= \sum_a (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes I_{BA''} \\ &= \sum_a \left( (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes I_{BA''} - (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes (\hat{M}_a^{(\text{POINT},W),u})_{BA''} \right) \\ &\quad + \sum_a (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes (\hat{M}_a^{(\text{POINT},W),u})_{BA''}. \end{aligned} \quad (200)$$

For projective sub-measurements  $\{A_i\}$  and  $\{B_i\}$  and a state  $|\varphi\rangle$ ,

$$\begin{aligned} \sum_i \langle \varphi | (A_i - B_i) | \varphi \rangle &= \sum_i \langle \varphi | A_i (A_i - B_i) | \varphi \rangle + \sum_i \langle \varphi | (A_i - B_i) B_i | \varphi \rangle \\ &\leq \left( \sum_i \langle \varphi | A_i^2 | \varphi \rangle \right)^{1/2} \left( \sum_i \langle \varphi | (A_i - B_i)^2 | \varphi \rangle \right)^{1/2} \\ &\quad + \left( \sum_i \langle \varphi | (A_i - B_i)^2 | \varphi \rangle \right)^{1/2} \left( \sum_i \langle \varphi | B_i^2 | \varphi \rangle \right)^{1/2} \\ &\leq 2 \left( \sum_i \langle \varphi | (A_i - B_i)^2 | \varphi \rangle \right)^{1/2}. \end{aligned}$$

Applying this fact to the projective sub-measurements  $\{(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'}\}_a$  and  $\{(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes (\hat{M}_a^{(\text{POINT},W),\mu})_{BA''}\}$  in (200) and using (199) gives

$$I \approx_{\sqrt{\varepsilon}} \sum_a (\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \otimes (\hat{M}_a^{(\text{POINT},W),\mu})_{BA''}. \quad (201)$$

To go from (201) to Item 3 of the lemma we use another fact about projective measurements. For families of projective measurements  $\{A_i^x\}$  and  $\{B_i^x\}$  acting on separate registers of a bipartite state  $|\phi\rangle$ , suppose  $I \approx_{\delta} \sum_i A_i^x \otimes B_i^x$ . Then

$$\begin{aligned} \delta &\geq \mathbb{E}_x \left\| \left( I - \sum_i A_i^x \otimes B_i^x \right) |\phi\rangle \right\|^2 \\ &= \mathbb{E}_x \left( 1 + \langle \phi | \left( \sum_i A_i^x \otimes B_i^x \right)^2 | \phi \rangle - 2 \sum_i \langle \phi | A_i^x \otimes B_i^x | \phi \rangle \right) \\ &= 1 - \mathbb{E}_x \sum_i \langle \phi | A_i^x \otimes B_i^x | \phi \rangle, \end{aligned}$$

where we used the projectivity in going from the second to the third line, to remove the square. Thus

$$\mathbb{E}_x \sum_{i \neq j} \langle \phi | A_i^x \otimes B_j^x | \phi \rangle \leq \delta,$$

and  $A_i^x \otimes I \simeq_{\delta} I \otimes B_i^x$ . By Item 1 of Fact 5.18, this in turn implies that  $A_i^x \otimes I \approx_{\delta} I \otimes B_i^x$ . Applying this fact to the projective measurements  $\{(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'}\}_a$  and  $\{(\hat{M}_a^{(\text{POINT},W),\mu})_{BA''}\}$  in (201) gives

$$(\hat{M}_{[\text{eval}_u(\cdot)=a]}^{(\text{LINE},W),\ell})_{AA'} \approx_{\sqrt{\varepsilon}} (\hat{M}_a^{(\text{POINT},W),\mu})_{BA''}.$$

Taking the function  $\delta_{\text{LINE}}$  in the lemma to be  $\sqrt{\varepsilon}$ , this yields the conclusion of Item 3 of the lemma.  $\square$

#### A.4 Combining the X and Z measurements

In this section our goal is to create a set of combined measurements that simultaneously measure the approximately-commuting measurements constructed in the previous section. We do this first for the points. Recall that  $\mathcal{H}_A$  and  $\mathcal{H}_B$  denote the Hilbert spaces corresponding to registers A and B, respectively.

**Lemma A.11.** *There exists a function  $\delta_Q(\varepsilon) = \text{poly}(\varepsilon)$  such that the following holds. For every  $x, z \in \mathbb{F}_q^m$  and  $\mathcal{H} \in \{\mathcal{H}_A, \mathcal{H}_B\}$  there are projective measurements  $\{\hat{Q}_{a,b}^{x,z}\}_{a,b \in \mathbb{F}_q}$  acting on  $\mathcal{H} \otimes (\mathbb{C}^q)^{\otimes M}$  such that:*

1. (**Self-consistency**) *On average over uniformly random  $x, z \in \mathbb{F}_q^m$ ,*

$$(\hat{Q}_{a,b}^{x,z})_{AA'} \approx_{\delta_Q} (\hat{Q}_{a,b}^{x,z})_{BA''}. \quad (202)$$

2. (**Consistency with  $\hat{M}$** ) *On average over uniformly random  $x, z \in \mathbb{F}_q^m$ ,*

$$(\hat{Q}_{a,b}^{x,z})_{AA'} \approx_{\delta_Q} (\hat{M}_a^{(\text{POINT},X),x} \hat{M}_b^{(\text{POINT},Z),z})_{BA''}, \quad (203)$$

$$(\hat{Q}_{a,b}^{x,z})_{AA'} \approx_{\delta_Q} (\hat{M}_b^{(\text{POINT},Z),z} \hat{M}_a^{(\text{POINT},X),x})_{BA''}. \quad (204)$$

Furthermore, all symmetric equivalents of these approximations also hold.

The construction of the measurements  $\{\hat{Q}_{a,b}^{x,z}\}$  whose existence is guaranteed in the lemma proceeds in three steps. In the first step, for every  $(r, s) \in \mathbb{F}_q^2$  we combine the binary-outcome measurements  $\hat{M}_a^{(\text{POINT}, X), x, r}$  and  $\hat{M}_b^{(\text{POINT}, X), z, s}$ ; this is made possible by their approximate commutation as shown in Lemma A.9. In a second step, we show that the measurements thus constructed have a particular ‘‘approximate linearity’’ property, when seen as a function of the pair  $(r, s)$ . This allows us, in the third step, to combine them into a single measurement  $\{\hat{Q}_{a,b}^{x,z}\}$  with outcomes in  $\mathbb{F}_q^2 \cong \mathbb{F}_2^{2t}$  by applying the following result from [NV17]:

**Theorem A.12** (Quantum linearity test). *Let  $0 \leq \delta \leq 1$ , let  $t$  be a positive integer, and let  $|\psi\rangle$  be a state in a Hilbert space  $\mathcal{H}_A \otimes \mathcal{H}_B$ . Let  $\{\mathcal{O}^u\}$  be a set of observables indexed by  $u \in \mathbb{F}_2^t$  that act on  $\mathcal{H}_A$ . Suppose that on average over  $u, u'$  chosen uniformly from  $\mathbb{F}_2^t$ , we have the following ‘‘approximate linearity’’ holds:*

$$(\mathcal{O}^u \mathcal{O}^{u'})_A \approx_\delta (\mathcal{O}^{u+u'})_A$$

where the  $\approx$  statement is taken with respect to the state  $|\psi\rangle$ . Then there exists an extended state  $|\psi'\rangle_{AA'B} = |\psi\rangle_{AB} \otimes |0\rangle_{A'}$  in an extended space  $\mathcal{H}_A \otimes \mathcal{H}_B \otimes \mathcal{H}_{A'}$  and observables  $\{\mathcal{L}^u\}$  acting on  $\mathcal{H}_A \otimes \mathcal{H}_{A'}$  such that for all  $u, u' \in \mathbb{F}_2^t$ ,

$$\mathcal{L}^u \mathcal{L}^{u'} = \mathcal{L}^{u+u'} \quad \text{and} \quad (\mathcal{L}^u)_{AA'} \approx_\delta (\mathcal{O}^u)_A.$$

Furthermore, the observable  $\mathcal{L}^u = I$  when  $u = 0$ .

We now give the proof of Lemma A.11.

*Proof of Lemma A.11.* For convenience in the proof we slightly abuse the notation  $\approx_\delta$  to mean  $\approx_{\text{poly}(\varepsilon)}$  for some polynomial function  $\delta = \text{poly}(\varepsilon)$  that may differ each time the notation is used.

**First step: construction of projective binary measurements  $\{\hat{R}_{a,b}^\omega\}$ .** For every  $\omega = (x, z, r, s)$  and  $a, b \in \mathbb{F}_2$ , define a measurement operator

$$\hat{R}_{a,b}^\omega = \hat{M}_b^{(\text{POINT}, Z), z, s} \cdot \hat{M}_a^{(\text{POINT}, X), x, r} \cdot \hat{M}_b^{(\text{POINT}, Z), z, s}.$$

It is clear that  $\{\hat{R}_{a,b}^\omega\}_{a,b \in \mathbb{F}_2}$  is a POVM. Observe that on average over uniformly random  $\omega$ ,

$$\hat{R}_{a,b}^\omega \approx_{\sqrt{\delta_{\text{ANTICOM}}}} \hat{M}_b^{(\text{POINT}, Z), z, s} \cdot \hat{M}_a^{(\text{POINT}, X), x, r}, \quad (205)$$

which follows from the approximate commutation relation of Lemma A.9 and the projectivity of  $\{\hat{M}_b^{(\text{POINT}, Z), z, s}\}$ , together with Fact 5.19. Next we show that  $\{\hat{R}_{a,b}^\omega\}$  is approximately self-consistent: we aim to show that on average over  $\omega$ , it holds that  $(\hat{R}_{a,b}^\omega)_{AA'} \simeq_\delta (\hat{R}_{a,b}^\omega)_{BA''}$  for some polynomial function  $\delta = \text{poly}(\varepsilon)$ . To show

this, we perform the following calculation: (recall the state  $|\hat{\psi}\rangle$  defined in (183))

$$\begin{aligned} & \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{R}_{a,b}^{\omega})_{AA'} \otimes (\hat{R}_{a,b}^{\omega})_{BA''} | \hat{\psi} \rangle \\ & \approx_{\delta_{\text{ANTICOM}}^{1/4}} \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes (\hat{R}_{a,b}^{\omega})_{BA''} | \hat{\psi} \rangle \end{aligned} \quad (206)$$

$$\approx_{\delta_{\text{ANTICOM}}^{1/4}} \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{BA''} | \hat{\psi} \rangle \quad (207)$$

$$\approx_{\sqrt{\varepsilon}} \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s})_{AA'} \otimes (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{BA''} | \hat{\psi} \rangle \quad (208)$$

$$= \mathbb{E}_{\omega} \sum_b \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s})_{AA'} \otimes (\hat{M}_b^{(\text{POINT,Z}),z,s})_{BA''} | \hat{\psi} \rangle$$

$$\approx_{\varepsilon} \mathbb{E}_{\omega} \sum_b \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s})_{AA'} | \hat{\psi} \rangle \quad (209)$$

$$= 1.$$

The approximation in (206) is derived by bounding the magnitude of the difference:

$$\begin{aligned} & \left| \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{R}_{a,b}^{\omega} - \hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes (\hat{R}_{a,b}^{\omega})_{BA''} | \hat{\psi} \rangle \right| \\ & \leq \sqrt{\mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{R}_{a,b}^{\omega} - \hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'}^2 | \hat{\psi} \rangle} \cdot \sqrt{\mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{R}_{a,b}^{\omega})_{BA''}^2 | \hat{\psi} \rangle} \\ & \leq \sqrt{\delta_{\text{ANTICOM}}^{1/2}} \cdot \sqrt{1}, \end{aligned}$$

where the second line follows from Cauchy-Schwarz, and the third line follows from (205) and the fact that  $\sum_{a,b} (\hat{R}_{a,b}^{\omega})^2 \leq I$ . The approximation in (207) follows from a similar calculation. The approximation in (208) is derived by bounding the magnitude of the difference:

$$\begin{aligned} & \left| \mathbb{E}_{\omega} \sum_{a,b} \langle \hat{\psi} | (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot (I - \hat{M}_a^{(\text{POINT,X}),x,r}))_{AA'} \otimes (\hat{M}_b^{(\text{POINT,Z}),z,s} \cdot \hat{M}_a^{(\text{POINT,X}),x,r})_{BA''} | \hat{\psi} \rangle \right| \\ & = \left| \mathbb{E}_{\omega} \langle \hat{\psi} | \left( \sum_b \hat{M}_b^{(\text{POINT,Z}),z,s} \otimes \hat{M}_b^{(\text{POINT,Z}),z,s} \right) \cdot \left( \sum_a (I - \hat{M}_a^{(\text{POINT,X}),x,r}) \otimes \hat{M}_a^{(\text{POINT,X}),x,r} \right) | \hat{\psi} \rangle \right| \end{aligned} \quad (210)$$

$$\leq \sqrt{\mathbb{E}_{\omega} \sum_a \langle \hat{\psi} | (I - \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'}^2 \otimes (\hat{M}_a^{(\text{POINT,X}),x,r})_{BA''}^2 | \hat{\psi} \rangle} \quad (211)$$

$$= \sqrt{\mathbb{E}_{\omega} \sum_a \langle \hat{\psi} | (I - \hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes (\hat{M}_a^{(\text{POINT,X}),x,r})_{BA''} | \hat{\psi} \rangle} \quad (212)$$

$$= \sqrt{1 - \mathbb{E}_{\omega} \sum_a \langle \hat{\psi} | (\hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes (\hat{M}_a^{(\text{POINT,X}),x,r})_{BA''} | \hat{\psi} \rangle} \quad (213)$$

$$= \sqrt{\frac{1}{2} \mathbb{E}_{\omega} \sum_a \langle \hat{\psi} | ((\hat{M}_a^{(\text{POINT,X}),x,r})_{AA'} \otimes I_{BA''} - I_{AA'} \otimes (\hat{M}_a^{(\text{POINT,X}),x,r})_{BA''})^2 | \hat{\psi} \rangle} \quad (214)$$

$$\leq \sqrt{\varepsilon}. \quad (215)$$

Here Equation (211) follows from the Cauchy-Schwarz inequality, the projectivity of  $\hat{M}_a^{(\text{POINT},X),x,r}$ , and the fact that  $\sum_b \hat{M}_b^{(\text{POINT},Z),z,s} \otimes \hat{M}_b^{(\text{POINT},Z),z,s} \leq I$ . Equations (212) to (214) follow from projectivity and completeness of  $\hat{M}_a^{(\text{POINT},X),x,r}$ . Equation (215) follows from the self-consistency of  $\hat{M}_a^{(\text{POINT},X),x,r}$ , established in Lemma A.9.

The approximation in (209) follows from the self-consistency of  $\hat{M}_a^{(\text{POINT},X),x,r}$ , again from Lemma A.9, together with projectivity and Fact 5.18.

This shows that  $(\hat{R}_{a,b}^\omega)_{AA'} \simeq_\delta (\hat{R}_{a,b}^\omega)_{BA''}$ , as desired. Combined with Fact 5.18, we have

$$(\hat{R}_{a,b}^\omega)_{AA'} \approx_\delta (\hat{R}_{a,b}^\omega)_{BA''} . \quad (216)$$

Finally we apply the orthonormalization Lemma (Lemma A.6) to  $\{\hat{R}_{a,b}^\omega\}$  in order to obtain projective measurements  $\{\hat{L}_{a,b}^\omega\}$  such that on average over  $\omega$ ,

$$\hat{L}_{a,b}^\omega \approx_\delta \hat{R}_{a,b}^\omega . \quad (217)$$

**Second step: approximate linearity relations for  $\{\hat{R}_{a,b}^\omega\}$ .** We proceed in three steps. First we establish linearity relations for  $\{\hat{M}_b^{(\text{POINT},W),u,r}\}$ . For all  $W \in \{X, Z\}$ ,  $u \in \mathbb{F}_q^m$ ,  $r, r' \in \mathbb{F}_q$  and  $c \in \mathbb{F}_2$ ,

$$\begin{aligned} \sum_{\substack{b,b' \in \mathbb{F}_2: \\ b+b'=c}} \hat{M}_b^{(\text{POINT},W),u,r} \hat{M}_{b'}^{(\text{POINT},W),u,r'} &= \sum_{\substack{a,a' \in \mathbb{F}_q: \\ \text{tr}(ar+a'r')=c}} \hat{M}_a^{(\text{POINT},W),u} \cdot \hat{M}_{a'}^{(\text{POINT},W),u} \\ &= \sum_{a: \text{tr}(a(r+r'))=c} \hat{M}_a^{(\text{POINT},W),u} \\ &= \hat{M}_c^{(\text{POINT},W),u,r+r'} , \end{aligned} \quad (218)$$

where the second line follows from projectivity of  $\{\hat{M}_a^{(\text{POINT},W),u}\}$ .

Second we show approximate linearity of the  $\{\hat{R}_{a,b}^\omega\}$ . On average over  $x, z \in \mathbb{F}_q^m$  and  $r, r', s, s' \in \mathbb{F}_q$ ,

$$\begin{aligned} \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{R}_{a,b}^{x,z,r,s} \cdot \hat{R}_{a',b'}^{x,z,r',s'} \right)_{AA'} \\ \approx_\delta \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{R}_{a,b}^{x,z,r,s} \right)_{AA'} \otimes \left( \hat{R}_{a',b'}^{x,z,r',s'} \right)_{BA''} \end{aligned} \quad (219)$$

$$\approx_\delta \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{M}_b^{(\text{POINT},Z),z,s} \cdot \hat{M}_a^{(\text{POINT},X),x,r} \right)_{AA'} \otimes \left( \hat{M}_{b'}^{(\text{POINT},Z),z,s'} \cdot \hat{M}_{a'}^{(\text{POINT},X),x,r'} \right)_{BA''} \quad (220)$$

$$\approx_\delta \sum_{\substack{b,b': \\ b+b'=d}} \left( \hat{M}_b^{(\text{POINT},Z),z,s} \right)_{AA'} \otimes \left( \hat{M}_{b'}^{(\text{POINT},Z),z,s'} \cdot \hat{M}_c^{(\text{POINT},X),x,r+r'} \right)_{BA''} \quad (221)$$

$$\approx_\delta \sum_{\substack{b,b': \\ b+b'=d}} \left( \hat{M}_b^{(\text{POINT},Z),z,s} \right)_{AA'} \otimes \left( \hat{M}_c^{(\text{POINT},X),x,r+r'} \cdot \hat{M}_{b'}^{(\text{POINT},Z),z,s'} \right)_{BA''} . \quad (222)$$

In each approximation, the answer summation is over  $c, d \in \mathbb{F}_2$ . Each approximation follows by applying a previously derived approximation to part of the expression, together with Fact 5.19. Specifically, line (219) uses the self-consistency of  $\{\hat{R}_{a,b}^\omega\}$  shown in (216). Line (220) follows from using (205) twice. Line (221) follows from the self-consistency and linearity properties of  $\{\hat{M}_a^{(\text{POINT},X),x,r}\}$  shown in Lemma A.9 and (218) respectively. Line (222) follows from the approximate commutativity relation shown in Lemma A.9. Continuing,

$$(222) \approx_\delta \left( \hat{M}_c^{(\text{POINT},X),x,r+r'} \cdot \hat{M}_d^{(\text{POINT},Z),z,s+s'} \right)_{\text{BA}''} \quad (223)$$

$$\approx_\delta \left( \hat{R}_{c,d}^{x,z,r+r',s+s'} \right)_{\text{BA}''} \quad (224)$$

$$\approx_\delta \left( \hat{R}_{c,d}^{x,z,r+r',s+s'} \right)_{\text{AA}'}. \quad (225)$$

Line (223) follows from self-consistency and linearity properties of  $\{\hat{M}_b^{(\text{POINT},Z),z,s}\}$ . Line (224) follows from (205) and Lemma A.9, and line (225) follows from the self-consistency of  $\{\hat{R}_{a,b}^\omega\}$  from Eq. (216).

Third we deduce approximate linearity of the  $\{\hat{L}_{a,b}^\omega\}$ . On average over  $x, z \in \mathbb{F}_q^m$ , and  $r, r', s, s' \in \mathbb{F}_q$ , we have

$$\begin{aligned} \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{L}_{a,b}^{x,z,r,s} \hat{L}_{a',b'}^{x,z,r',s'} \right)_{\text{AA}'} &\approx_\delta \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{L}_{a,b}^{x,z,r,s} \right)_{\text{AA}'} \otimes \left( \hat{L}_{a',b'}^{x,z,r',s'} \right)_{\text{BA}''} \\ &\approx_\delta \sum_{\substack{a,b,a',b': \\ a+a'=c \\ b+b'=d}} \left( \hat{R}_{a,b}^{x,z,r,s} \right)_{\text{AA}'} \otimes \left( \hat{R}_{a',b'}^{x,z,r',s'} \right)_{\text{BA}''} \\ &\approx_\delta \left( \hat{R}_{c,d}^{x,z,r+r',s+s'} \right)_{\text{AA}'} \\ &\approx_\delta \left( \hat{L}_{c,d}^{x,z,r+r',s+s'} \right)_{\text{AA}'}. \end{aligned} \quad (226)$$

The first two approximations follow from the closeness of the  $\{\hat{R}_{a,b}^\omega\}$  and  $\{\hat{L}_{a,b}^\omega\}$  guaranteed by (217) as well as the self-consistency of  $\{\hat{R}_{a,b}^\omega\}$  shown in (216). The third approximation follows from the linearity properties of  $\{\hat{R}_{a,b}^\omega\}$  in (222).

**Third step: construction of  $\{\hat{\mathcal{O}}_{a,b}^{x,z}\}$ .** For each  $\omega$  define an observable  $\mathcal{O}^\omega = \sum_{a,b \in \mathbb{F}_2} (-1)^{a+b} \hat{L}_{a,b}^\omega$ . The self-consistency of  $\mathcal{O}^\omega$  between registers AA' and BA'' follows from the self-consistency of the  $\{\hat{L}_{a,b}^\omega\}$ , which is obtained by combining (216) and (217). We now verify approximate linearity. On average over  $x, z \in \mathbb{F}_q^m$ , and  $r, r', s, s' \in \mathbb{F}_q$ ,

$$\begin{aligned} \mathcal{O}^{x,z,r,s} \mathcal{O}^{x,z,r',s'} &= \sum_{a,b,a',b'} (-1)^{a+b+a'+b'} \hat{L}_{a,b}^{x,z,r,s} \hat{L}_{a',b'}^{x,z,r',s'} \\ &\approx_\delta \sum_{c,c'} (-1)^{c+c'} \hat{L}_{c,c'}^{x,z,r+r',s+s'} \\ &= \mathcal{O}^{x,z,r+r',s+s'} \end{aligned}$$

where the approximation follows from the approximate linearity properties of  $\{\hat{L}_{a,b}^\omega\}$  shown in (226). Identify the field  $\mathbb{F}_q$  with the vector space  $\mathbb{F}_2^t$ , and thus we treat pairs  $(r, s)$  as vectors in  $\mathbb{F}_2^{2t}$ . For every  $x, z \in \mathbb{F}_q^m$ ,

we can apply Theorem A.12 to the collection of observables  $\{\mathcal{O}^{x,z,r,s}\}_{r,s \in \mathbb{F}_q}$  to obtain the existence of “exactly linear” observables  $\{\mathcal{L}^{x,z,r,s}\}$  that are  $\delta$ -close to  $\{\mathcal{O}^{x,z,r,s}\}$ , on average over the choice of  $x, z, r, s$ . By assuming that  $|\hat{\psi}\rangle$  has sufficiently many ancilla zero qubits, we do not need to extend the state further.

We are ready to define the measurements  $\{\hat{Q}_{a,b}^{x,z}\}$  whose existence is claimed in the statement of the lemma. For all  $x, z \in \mathbb{F}_q^m$  and  $a, b \in \mathbb{F}_q$ , define

$$\hat{Q}_{a,b}^{x,z} = \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{L}^{x,z,r,s} .$$

This operator is a projection (and thus positive semidefinite):

$$\begin{aligned} (\hat{Q}_{a,b}^{x,z})^2 &= \mathbb{E}_{r,s,r',s'} (-1)^{\text{tr}((r+r')a+(s+s')b)} \mathcal{L}^{x,z,r,s} \mathcal{L}^{x,z,r',s'} \\ &= \mathbb{E}_{r,s,r',s'} (-1)^{\text{tr}((r+r')a+(s+s')b)} \mathcal{L}^{x,z,r+r',s+s'} \\ &= \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{L}^{x,z,r,s} . \end{aligned}$$

The second equality follows from the exact linearity of the  $\{\mathcal{L}^{x,z,r,s}\}$ . Next, the operators sum to identity:

$$\begin{aligned} \sum_{a,b} \hat{Q}_{a,b}^{x,z} &= \mathbb{E}_{r,s} \sum_{a,b} (-1)^{\text{tr}(ra+sb)} \mathcal{L}^{x,z,r,s} \\ &= \mathcal{L}^{x,z,0,0} \\ &= I . \end{aligned}$$

Thus  $\{\hat{Q}_{a,b}^{x,z}\}$  forms a projective measurement.

We establish self-consistency:

$$\begin{aligned} \left( \hat{Q}_{a,b}^{x,z} \right)_{AA'} &= \left( \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{L}^{x,z,r,s} \right)_{AA'} \\ &\approx_{\delta} \left( \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{O}^{x,z,r,s} \right)_{AA'} \\ &\approx_{\delta} \left( \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{O}^{x,z,r,s} \right)_{BA''} \\ &\approx_{\delta} \left( \hat{Q}_{a,b}^{x,z} \right)_{BA''} . \end{aligned}$$

The first approximation is due to Lemma A.4. The second approximation is due to Lemma A.4 and the self-consistency of the  $\{\mathcal{O}^{x,z,r,s}\}$  observables. The third approximation follows from Lemma A.4 again.



Finally we show consistency with the  $\{\mathcal{M}_a^{(\text{POINT},W),u}\}$ .

$$\begin{aligned}
(\hat{Q}_{a,b}^{x,z})_{AA'} &\approx_\delta \left( \mathbb{E}_{r,s} (-1)^{\text{tr}(ra+sb)} \mathcal{O}^{x,z,r,s} \right)_{BA''} \\
&= \left( \mathbb{E}_{r,s} \sum_{c,d} (-1)^{\text{tr}(ra+sb)+c+d} \hat{L}_{c,d}^{x,z,r,s} \right)_{BA''} \\
&\approx_\delta \left( \mathbb{E}_{r,s} \sum_{c,d} (-1)^{\text{tr}(ra+sb)+c+d} \hat{R}_{c,d}^{x,z,r,s} \right)_{BA''} \\
&\approx_\delta \left( \mathbb{E}_{r,s} \sum_{c,d} (-1)^{\text{tr}(ra+sb)+c+d} \hat{M}_d^{(\text{POINT},Z),z,s} \hat{M}_c^{(\text{POINT},X),x,r} \right)_{BA''} \\
&= \left( \mathbb{E}_s \sum_d (-1)^{\text{tr}(sb)+d} \hat{M}_d^{(\text{POINT},Z),z,s} \right) \cdot \left( \mathbb{E}_r \sum_c (-1)^{\text{tr}(ra)+c} \hat{M}_c^{(\text{POINT},X),x,r} \right)_{BA''} \\
&= \left( \mathbb{E}_s \sum_d \sum_{b',a'} (-1)^{\text{tr}(sb)+\text{tr}(sb')} \hat{M}_{b'}^{(\text{POINT},Z),z} \right)_{BA''} \cdot \left( \mathbb{E}_r \sum_c (-1)^{\text{tr}(ra)+\text{tr}(ra')} \hat{M}_{b'}^{(\text{POINT},X),x} \right)_{BA''} \\
&= \left( \hat{M}_b^{(\text{POINT},Z),z} \cdot \hat{M}_a^{(\text{POINT},X),x} \right)_{BA''}.
\end{aligned}$$

The second line follows from the definition of  $\{\mathcal{O}^{x,z,r,s}\}$ . The third line follows from the closeness of  $\{\hat{L}_{c,d}^\omega\}$  and  $\{\hat{R}_{c,d}^\omega\}$ . The fourth line follows from (205). The fifth line follows from (186), and the last line from Lemma 3.25.

The above calculation establishes Equation (204) of the Lemma. To obtain Equation (203), which is the same consistency relation with  $X$  and  $Z$  swapped, we may use Lemma A.9 immediately after the fourth line to swap the two operators in the product, and then proceed as above.

An analogous derivation shows the same properties of  $\{\hat{Q}_{a,b}^{x,z}\}$  with the tensor factors switched to  $BB'$  and  $AB''$ , respectively. This concludes the proof of the lemma.  $\square$

**Lemma A.13.** *There exists a function  $\delta_P(\varepsilon, m, d, q) = \text{poly}(\varepsilon, md/q)$  such that the following holds. For  $\mathcal{H} \in \{\mathcal{H}_A, \mathcal{H}_B\}$  and for every pair of lines  $\ell_X$  and  $\ell_Z$  there exists a POVM  $\{T_{f_X, f_Z}^{\ell_X, \ell_Z}\}$  acting on  $\mathcal{H} \otimes (\mathbb{C}^q)^{\otimes M}$  with answers consisting of pairs of polynomials  $f_X, f_Z$  in  $\text{deg}_{md}(\ell_X) \times \text{deg}_{md}(\ell_Z)$  such that on average over pairs  $(\ell_X, x)$  and  $(\ell_Z, z)$  independently sampled from the line-point distribution,*

$$(T_{[\text{eval}_x(\cdot)=a, \text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z})_{AA'} \simeq_{\delta_P} (\hat{Q}_{a,b}^{x,z})_{BA''},$$

as well as all symmetric equivalents of this approximation. Moreover, if  $\ell_W$  is axis-parallel, then  $f_W \in \text{deg}_d(\ell_W)$ .

*Proof.* We construct the required measurements by combining the measurements  $\hat{M}_{f_W}^{(\text{LINE},W),\ell_W}$  for  $W \in \{X, Z\}$ . For convenience in the proof we slightly abuse the notation  $\approx_\delta$  to mean  $\approx_{\text{poly}(\varepsilon)}$  for some polynomial function  $\delta = \text{poly}(\varepsilon)$  that may differ each time the notation is used.

We first argue that the measurements  $\{\hat{Q}_{a,b}^{x,z}\}$  obtained in Lemma A.11 are consistent with the  $\{\hat{M}_{f_W}^{(\text{LINE},W),\ell_W}\}$ .

Applying Fact 5.19 to (204), we get that on average over uniformly random  $x, z$ ,

$$\begin{aligned} \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \cdot \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} &= \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} \cdot \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \\ &\approx_{\delta} \left(\hat{M}_b^{(\text{POINT},Z),z} \hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} \\ &\approx_{\delta} \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'}. \end{aligned}$$

Since  $\{\hat{Q}_{a,b}^{x,z}\}$  is a projective measurement, applying Lemma 5.21 with  $(\hat{Q}_{a,b}^{x,z})_{AA'}$  playing the role of  $A_a^x$  and  $(\hat{Q}_{a,b}^{x,z})_{AA'} \cdot (\hat{M}_b^{(\text{POINT},Z),z})_{BA''}$  playing the role of  $B_a^x$ , we obtain that

$$\left(\hat{Q}_a^{x,z}\right)_{AA'} \approx_{\delta} \sum_b \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \cdot \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''}. \quad (227)$$

We now show that on average over uniformly random  $x, z$ ,

$$\left(\hat{Q}_a^{x,z}\right)_{AA'} \approx_{\delta_Q} \left(\hat{M}_a^{(\text{POINT},X),x}\right)_{BA''}. \quad (228)$$

This follows from (227) and the following calculation:

$$\begin{aligned} &\mathbb{E}_{x,z} \sum_a \left\| \sum_b \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \cdot \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} - \left(\hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} |\hat{\psi}\rangle \right\|^2 \\ &= \mathbb{E}_{x,z} \sum_a \left\| \sum_b \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} \cdot \left( \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} - \left(\hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} \right) |\hat{\psi}\rangle \right\|^2 \\ &= \mathbb{E}_{x,z} \sum_{a,b} \left\| \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} \cdot \left( \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} - \left(\hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} \right) |\hat{\psi}\rangle \right\|^2 \\ &= \mathbb{E}_{x,z} \sum_{a,b} \left\| \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''} \cdot \left( \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} - \left(\hat{M}_b^{(\text{POINT},Z),z} \cdot \hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} \right) |\hat{\psi}\rangle \right\|^2 \\ &\leq \mathbb{E}_{x,z} \sum_{a,b} \left\| \left( \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} - \left(\hat{M}_b^{(\text{POINT},Z),z} \cdot \hat{M}_a^{(\text{POINT},X),x}\right)_{BA''} \right) |\hat{\psi}\rangle \right\|^2 \\ &= O(\delta), \end{aligned}$$

where the second line uses the property  $\sum_b \hat{M}_b^{(\text{POINT},Z),z} = I$ , the third and fourth lines uses the projectivity of  $\{\hat{M}_b^{(\text{POINT},Z),z}\}$ , the fifth line uses  $\hat{M}_b^{(\text{POINT},Z),z} \leq I$ , and the last line uses Equation (204). This shows (228). A similar argument shows that

$$\left(\hat{Q}_b^{x,z}\right)_{AA'} \approx_{\delta_Q} \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''}. \quad (229)$$

Next, from Lemma A.10 we get that on average over  $(\ell_X, x)$  and  $(\ell_Z, z)$  sampled independently from the line-point distribution (Definition 7.6),

$$\left(\hat{M}_{[\text{eval}_x(\cdot)=a]}^{(\text{LINE},X),\ell_X}\right)_{BA''} \approx_{\delta} \left(\hat{M}_a^{(\text{POINT},X),x}\right)_{BA''}, \quad (230)$$

$$\left(\hat{M}_{[\text{eval}_z(\cdot)=b]}^{(\text{LINE},Z),\ell_Z}\right)_{BA''} \approx_{\delta} \left(\hat{M}_b^{(\text{POINT},Z),z}\right)_{BA''}. \quad (231)$$

where we used both the self-consistency of the line measurements (item 1 in Lemma A.10) as well as their consistency with the points measurements (item 3). To deduce both of these relations, we used Fact 5.18 to convert between  $\approx$  and  $\simeq$  distance, and Fact 5.24 to sum over outcomes to the lines measurement. Since the  $\{\hat{M}_f^{(\text{LINE},W),\ell}\}$  and  $\{\hat{Q}_{a,b}^{x,z}\}$  are projective, together with (228) and (229) by Fact 5.18 we get

$$\left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \simeq_{\delta_Q} \left(\hat{M}_{[\text{eval}_x(\cdot)=a]}^{(\text{LINE},X),\ell_X}\right)_{BA''} \quad \text{and} \quad \left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \simeq_{\delta_Q} \left(\hat{M}_{[\text{eval}_z(\cdot)=b]}^{(\text{LINE},Z),\ell_Z}\right)_{BA''}. \quad (232)$$

For all lines  $\ell_X, \ell_Z$  and  $(f_X, f_Z) \in \text{deg}_{md}(\ell_X) \times \text{deg}_{md}(\ell_Z)$  define the operator

$$T_{f_X, f_Z}^{\ell_X, \ell_Z} = \hat{M}_{f_X}^{(\text{LINE},X),\ell_X} \cdot \hat{M}_{f_Z}^{(\text{LINE},Z),\ell_Z} \cdot \hat{M}_{f_X}^{(\text{LINE},X),\ell_X}.$$

Then the collection  $\{T_{f_X, f_Z}^{\ell_X, \ell_Z}\}$  forms a valid POVM. Moreover, by the fact that  $\{\hat{M}_{f_W}^{(\text{LINE},W),\ell_W}\}$  comes from a valid strategy for the low-degree test, it holds that  $f_W \in \text{deg}_d(\ell_W)$  whenever  $\ell_W$  is axis-parallel, and thus property holds for  $T$  as well.

It only remains to bound the consistency of  $\{T_{f_X, f_Z}^{\ell_X, \ell_Z}\}$  with  $\{\hat{Q}_{a,b}^{x,z}\}$ . Using the following choices of measurements,

$$"A_{a_1, a_2}^{x, y_1, y_2}" : \hat{Q}_{a, b}^{x, z}, \quad "(G_1)_g^{x"} : \hat{M}_{f_X}^{(\text{LINE},X),\ell_X}, \quad "(G_2)_g^{x"} : \hat{M}_{f_Z}^{(\text{LINE},Z),\ell_Z}, \quad "J_{g_1, g_2}^{x}" : T_{f_X, f_Z}^{\ell_X, \ell_Z},$$

Then the hypotheses of Lemma 5.27 hold, with  $\delta$  in the lemma set to  $\delta_Q$ , and  $\eta$  in the lemma set to  $md/q$  (this follows from the Schwartz-Zippel lemma applied to the polynomials  $f_X, f_Z$ , which are of total degree at most  $md$ ). Applying the lemma, we conclude that

$$\left(\hat{Q}_{a,b}^{x,z}\right)_{AA'} \simeq_{\delta_P} \left(T_{[\text{eval}_x(\cdot)=a, \text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z}\right)_{BA''}, \quad (233)$$

where  $\delta_P = \text{poly}(\varepsilon, md/q)$  is the  $\delta_{\text{pasting}}$  as given by Lemma 5.27, conditions (42) and (43) in the lemma are given by (232) and (202) respectively, and the distribution over  $(\ell_X, \ell_Z)$  are two independently and uniformly chosen lines. This is a symmetric equivalent of the consistency relation in the conclusion of Lemma. Thus, by applying an identical argument with the appropriate modifications to the registers, all symmetric equivalents of this relation hold, establishing the conclusion of the Lemma.  $\square$

## A.5 Applying the classical low-degree test

In the previous section we showed how to combine the approximately commuting  $X$  and  $Z$  basis point and line measurements into a joint point measurement and a joint line measurement. In this section we show how these measurements can be combined in a single measurement that returns a pair of global polynomials encoding the  $X$ -basis and  $Z$ -basis information, respectively, by applying the lemma that states quantum soundness of the classical low-degree test, Theorem 7.8.

Introduce a ‘‘combining’’ map that takes two polynomials  $f, g \in \text{ideg}_{d,m}(\mathbb{F}_q)$  and returns a new polynomial  $\text{combine}_{f,g} \in \text{ideg}_{d,2m+2}(\mathbb{F}_q)$  defined by

$$\text{combine}_{f,g} \left( \underbrace{x}_{\in \mathbb{F}_q^m}, \underbrace{z}_{\in \mathbb{F}_q^m}, \underbrace{\alpha}_{\in \mathbb{F}_q}, \underbrace{\beta}_{\in \mathbb{F}_q} \right) = \alpha f(x) + \beta g(z).$$

This combining map can also be defined for polynomials restricted to lines. Let  $\ell$  be a line in  $\mathbb{F}_q^{2m+2}$  with intercept  $u = (u_X, u_Z, u_\alpha, u_\beta)$  and slope  $v = (v_X, v_Z, v_\alpha, v_\beta)$ . Let  $\ell_X$  and  $\ell_Z$  be lines in  $\mathbb{F}_q^m$  such that for

any point  $(x, z, \alpha, \beta) \in \ell$ , it holds that  $x \in \ell_X$  and  $z \in \ell_Z$ . Then for any two polynomials  $f \in \deg_{dm}(\ell_X)$  and  $g \in \deg_{dm}(\ell_Z)$ , we define  $\text{combine}_{f,g} \in \deg_{dm+1}(\ell)$  by

$$\text{combine}_{f,g}(u + t \cdot v) = (u_\alpha + t \cdot v_\alpha)f(u_X + t \cdot v_X) + (u_\beta + t \cdot v_\beta)g(u_Z + t \cdot v_Z). \quad (234)$$

In the following two lemma we define combined point and line measurements according to this definition of  $\text{combine}_{f,g}$ .

**Lemma A.14.** *For all  $x, z \in \mathbb{F}_q^m$  and  $\alpha, \beta \in \mathbb{F}_q$  and  $\mathcal{H} \in \{\mathcal{H}_A, \mathcal{H}_B\}$  there exists a projective measurement  $\{\hat{Q}_c^{x,z,\alpha,\beta}\}_{c \in \mathbb{F}_q}$  acting on  $\mathcal{H} \otimes (\mathbb{C}^q)^{\otimes M}$  such that the following holds:*

1. (**Self-consistency**) *On average over uniformly random  $(x, z, \alpha, \beta) \in \mathbb{F}_q^{2m+2}$ :*

$$(\hat{Q}_c^{x,z,\alpha,\beta})_{AA'} \approx_{\delta_Q} (\hat{Q}_c^{x,z,\alpha,\beta})_{BA''}. \quad (235)$$

2. (**Consistency with  $\hat{M}$** ) *On average over uniformly random  $(x, z, \alpha, \beta) \in \mathbb{F}_q^{2m+2}$ :*

$$(\hat{Q}_c^{x,z,\alpha,\beta})_{AA'} \approx_{\delta_Q} \left( \sum_{\substack{a,b: \\ \alpha a + \beta b = c}} \hat{M}_a^{(\text{POINT}, X), x} \hat{M}_b^{(\text{POINT}, Z), z} \right)_{BA''}, \quad (236)$$

$$(\hat{Q}_c^{x,z,\alpha,\beta})_{AA'} \approx_{\delta_Q} \left( \sum_{\substack{a,b: \\ \alpha a + \beta b = c}} \hat{M}_b^{(\text{POINT}, Z), z} \hat{M}_a^{(\text{POINT}, X), x} \right)_{BA''}. \quad (237)$$

Furthermore, all symmetric equivalents of these approximations also hold.

*Proof.* Define

$$\hat{Q}_c^{x,z,\alpha,\beta} = \sum_{\substack{a,b: \\ \alpha a + \beta b = c}} \hat{Q}_{a,b}^{x,z}.$$

The self-consistency and consistency properties of  $\{\hat{Q}_c^{x,z,\alpha,\beta}\}$  follow from the consistency properties of  $\{\hat{Q}_{a,b}^{x,z}\}$  established by Lemma A.11, and Fact 5.24.  $\square$

**Lemma A.15.** *There exists a function  $\delta_{\text{combine}}(\varepsilon, m, d, q) = \text{poly}(m^2\varepsilon, md/q)$  such that the following holds. For every line  $\ell$  in  $\mathbb{F}_q^{2m+2}$  and  $\mathcal{H} \in \{\mathcal{H}_A, \mathcal{H}_B\}$  there exists a POVM  $\{\hat{Q}_f^\ell\}$  acting on  $\mathcal{H} \otimes (\mathbb{C}^q)^{\otimes M}$  with outcomes  $f \in \deg_{md+1}(\ell)$  such that on average over a uniformly random pair  $(\ell, (x, z, \alpha, \beta))$  drawn from the line-point distribution over  $\mathbb{F}_q^{2m+2}$ ,*

$$(\hat{Q}_{[\text{eval}_{(x,z,\alpha,\beta)}(\cdot)=a]}^\ell)_{AA'} \simeq_{\delta_{\text{combine}}} (\hat{Q}_a^{x,z,\alpha,\beta})_{BA''}, \quad (238)$$

$$(\hat{Q}_{[\text{eval}_{(x,z,\alpha,\beta)}(\cdot)=a]}^\ell)_{BB'} \simeq_{\delta_{\text{combine}}} (\hat{Q}_a^{x,z,\alpha,\beta})_{AB''}, \quad (239)$$

where the answer summation is over  $a \in \mathbb{F}_q$ . Moreover, if  $\ell$  is axis-parallel, then the outcome  $f \in \deg_d(\ell)$ .

Before proving the lemma we give definitions regarding distributions over lines and points.

**Definition A.16.** For any  $i \in \{1, \dots, m\}$  the  $i$ -th restricted axis parallel line distribution  $D_{\text{ALINE},i}$  is the restriction of  $D_{\text{ALINE}}$  to pairs  $(\ell, u)$  where the line  $\ell$  is along the direction  $e_i$  (i.e. is of the form  $\ell = (u_0, e_i)$ ). Similarly, the  $i$ -th restricted diagonal line distribution  $D_{\text{DLINE},i}$  is the restriction of  $D_{\text{DLINE}}$  to pairs  $(\ell, u)$ , where the line  $\ell = (u_0, s, v)$  is such that  $\chi(s) = i$ , and coordinates  $1, \dots, i-1$  of  $v$  are 0. (Recall that  $\chi$  is defined in (50).)

Observe that  $D_{\text{ALINE}}$  is a uniform mixture over the distributions  $D_{\text{ALINE},i}$  for  $i \in \{1, \dots, m\}$ , and likewise for  $D_{\text{DLINE}}$ . As a consequence, any approximation that holds on average over the line-point distribution  $D_{\text{LINE}}$  with error  $\delta$  holds over each of the  $i$ -th restricted distributions  $D_{\text{ALINE},i}$  and  $D_{\text{DLINE},i}$  with error  $2m\delta$ . Similarly, any approximation that holds over  $D_{\text{LINE}} \times D_{\text{LINE}}$  holds over any product of the  $i$ -th restricted distributions with error  $4m^2\delta$ . For example, by Lemma A.13 for any  $t_1, t_2 \in \{\text{ALINE}, \text{DLINE}\}$  and  $i, j \in \{1, \dots, m\}$ , we have

$$\mathbb{E}_{(\ell_X, x) \sim D_{t_1, i}} \mathbb{E}_{(\ell_Z, z) \sim D_{t_2, j}} \sum_{f_X, f_Z} \langle \hat{\psi} | (T_{f_X, f_Z}^{\ell_X, \ell_Z})_{\text{AA}'} \otimes (I - (\hat{Q}_{f_X(x), f_Z(z)}^{x, z}))_{\text{BA}''} | \hat{\psi} \rangle = O(m^2 \delta_P). \quad (240)$$

**Lemma A.17.** *There exists a distribution  $D$  over tuples  $(\ell, \ell_X, \ell_Z)$  with the following properties:*

1. *The marginal distribution over  $\ell$  is the same as the marginal of the line-point distribution over  $\mathbb{F}_q^{2m+2}$ .*
2. *For any point of the form  $u = (u_X, u_Z, \alpha, \beta)$  lying on  $\ell$ , it holds that  $u_X \in \ell_X$  and  $u_Z \in \ell_Z$ . Moreover, for  $u$  chosen uniformly at random in  $\ell$  the marginal distribution over  $(\ell_X, \ell_Z, u_X)$  (resp. over  $(\ell_X, \ell_Z, u_Z)$ ) is such that the marginal over  $(\ell_X, \ell_Z)$  is a mixture of  $D_{\text{ALINE},i} \times D_{\text{ALINE},j}$  and  $D_{\text{DLINE},i} \times D_{\text{DLINE},j}$  for  $i, j \in \{1, \dots, m\}$  and moreover, conditioned on  $(\ell_X, \ell_Z)$ ,  $u_X$  is uniformly random on  $\ell_X$  (resp.  $u_Z$  uniformly random on  $\ell_Z$ ).*

*Proof.* We describe  $D$  by giving a procedure to sample from it. We start by sampling  $(\ell, u)$  from the line-point distribution over  $\mathbb{F}_q^{2m+2}$ . Write  $u = (u_X, u_Z, \alpha, \beta)$ . There are two cases to consider: the case where  $\ell$  is axis-parallel and the case where  $\ell$  is diagonal. In each case, we sample lines  $\ell_X$  and  $\ell_Z$  such that the tuples  $(\ell_X, \ell_Z, u_X)$  and  $(\ell_X, \ell_Z, u_Z)$  satisfy Property 2 of the lemma.

1. Suppose  $\ell$  is an axis-parallel line  $(v, e_j)$  where  $v = (v_X, v_Z, \alpha_0, \beta_0)$ . Then we sample  $(v'_X, i_X), (v'_Z, i_Z) \in \mathbb{F}_q^m \times \{1, 2, \dots, m\}$  as described below, and set  $\ell_X = (v'_X, e_{i_X}), \ell_Z = (v'_Z, e_{i_Z})$ .
  - (a) If  $1 \leq j \leq m$ , then set  $i_X = j$  and choose  $i_Z$  to be uniformly random in  $\{1, \dots, m\}$ .
  - (b) If  $m+1 \leq j \leq 2m$ , then set  $i_X$  to be uniformly random in  $\{1, \dots, m\}$  and  $i_Z = j - m$ .
  - (c) If  $2m+1 \leq j \leq 2m+2$ , then set  $i_X$  and  $i_Z$  to both be uniformly random in  $\{1, \dots, m\}$ .
  - (d) Let  $v'_X$  be the canonical representative  $L_{e_{i_X}}^{\text{LN}}(v_X)$ , and likewise let  $v'_Z$  be the canonical representative  $L_{e_{i_Z}}^{\text{LN}}(v_Z)$  (see Definition 7.3 for the definition of the canonical representative of a line). Note that  $v_X$  lies on the line  $\ell_X = (v'_X, e_{i_X})$ , and similarly for  $v_Z$  and  $\ell_Z = (v'_Z, e_{i_Z})$ .

We claim that this construction satisfies Property 2 of the lemma. We show this first  $(\ell_X, \ell_Z)$  and  $u_X$ , the other case being symmetric. If we are in case (a), then  $u_X$  is a uniformly random point on the line  $\ell_X = (v_X, e_j)$  and  $\ell_Z = (v_Z, e_{i_Z})$ , which is independent from  $\ell_X$ . Thus, the marginal distribution over  $(\ell_X, u_X)$  conditioned on  $j$  in this case is equal to  $D_{\text{ALINE},j}$  and furthermore even conditioned on  $(\ell_X, u_X)$ ,  $\ell_Z$  is distributed according to  $D_{\text{ALINE},i_Z}$  for a uniformly random  $i_Z$ . Cases (b) and (c) are analogous except that in this case even conditioned on  $j$ ,  $i_X$  is uniform.

2. Suppose  $\ell$  is a random diagonal line  $(v, s, w)$ , where  $v = (v_X, v_Z, \alpha, \beta)$ , and where  $w = (w_X, w_Z, \gamma, \delta)$  is a direction vector whose first nonzero coordinate is  $j = \chi(s)$  (where  $\chi$  is defined in Equation (50)). We sample  $\ell_X = (v'_X, s_X, w'_X)$  and  $\ell_Z = (v'_Z, s_Z, w'_Z)$  in the following manner:
  - (a) If  $1 \leq j \leq m$ , then  $w'_X = w_X$  and  $w'_Z = w_Z$ . We choose  $s_X$  and  $s_Z$  such that  $\chi(s_X) = j$  and  $\chi(s_Z) = 1$ .
  - (b) If  $m + 1 \leq j \leq 2m$ , then  $w'_X$  is a random diagonal line direction (and  $s_X$  is chosen accordingly), and  $w'_Z = w_Z$ , with  $s_Z$  such that  $\chi(s_Z) = j - m$ .
  - (c) If  $2m + 1 \leq j \leq 2m + 2$  then both  $w_X$ , and  $w_Z$  are uniformly random, with  $s_X$  and  $s_Z$  chosen accordingly.
  - (d) Set  $v'_X = L_{w'_X}^{L_N}(v_X)$  and  $v'_Z = L_{w'_Z}^{L_N}(v_Z)$ .

We claim that this construction satisfies Property 2 of the Lemma. We show this first  $(\ell_X, \ell_Z)$  and  $u_X$ , the other case being symmetric. If we are in case (a), then  $u_X$  is a uniformly random point on the diagonal line  $(v_X, s_X, w_X)$  which is equal to  $\ell_X$ . Thus, the marginal distribution over  $(\ell_X, u_X)$  conditioned on  $j$  in this case is equal to  $D_{\text{DLINE},j}$ . Furthermore, conditioned on  $(\ell_X, u_X)$ ,  $\ell_Z$  is distributed according to  $D_{\text{DLINE},1}$ . The other cases can be analyzed analogously, with the only difference being how the indices  $i, j$  of  $\ell_X$  and  $\ell_Z$  distributions are chosen.

Let  $D$  be the distribution over tuples  $(\ell, \ell_X, \ell_Z)$  induced by this procedure. This distribution  $D$  satisfies Property 1 in the lemma by construction, and Property 2 as argued above.  $\square$

*Proof of Lemma A.15.* We construct the measurements  $\hat{Q}_f^\ell$  out of the lines measurements  $T_{f_X, f_Z}^{\ell_X, \ell_Z}$  guaranteed by Lemma A.13. To do so we first show how to map any line-point pair  $(\ell, u)$  in the expanded space  $\mathbb{F}_q^{2m+2}$  to two pairs  $(\ell_X, u_X)$  and  $(\ell_Z, u_Z)$  in  $\mathbb{F}_q^m$ . We will define  $\hat{Q}^\ell$  in terms of  $T_{f_X, f_Z}^{\ell_X, \ell_Z}$  and show that the resulting measurement is consistent with the combined points measurement  $\hat{Q}^{x,z,\alpha,\beta}$  from Lemma A.14.

To define  $\hat{Q}^\ell$  we make use of the distribution  $D$  from Lemma A.17. Let  $D|_\ell$  denote the distribution derived from  $D$  by conditioning on the first element being  $\ell$ . Then for every line  $\ell$  in  $\mathbb{F}_q^{2m+2}$  and function  $f \in \text{deg}_{(md+1)}(\ell)$ , define

$$\hat{Q}_f^\ell = \mathbb{E}_{(\ell_X, \ell_Z) \sim D|_\ell} \sum_{\substack{f_X \in \text{deg}_{md}(\ell_X), f_Z \in \text{deg}_{md}(\ell_Z): \\ f = \text{combine}_{f_X, f_Z}}} T_{f_X, f_Z}^{\ell_X, \ell_Z},$$

where  $\text{combine}_{f_X, f_Z}$  is defined in (234). Note that  $\{\hat{Q}_f^\ell\}$  forms a valid POVM for every choice of line  $\ell$ , and moreover that  $\text{combine}_{f_X, f_Z}$  is well-defined for the lines  $\ell, \ell_X, \ell_Z$  because of Property 2 of Lemma A.17.

Observe that by Property 3 of Lemma A.17, whenever  $\ell$  is axis-parallel,  $\ell_X$  and  $\ell_Z$  are axis-parallel also. Thus, in this case, by Lemma A.13 it holds that the outcomes  $f_X$  and  $f_Z$  both have degree  $d$ . By definition of  $\text{combine}_{f_X, f_Z}$  it holds that the outcome  $f$  of  $\{\hat{Q}_f^\ell\}$  has degree  $d$  as well.

We argue the following bound:

$$\mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes (I - \hat{Q}_{f_X(x), f_Z(z)}^{x,z}) | \hat{\psi} \rangle = O(m \cdot (\epsilon^{1/4} + \delta_P^{1/4} + \delta_Q^{1/4} + \delta_{\text{LINE}}^{1/2})). \quad (241)$$

This bound will follow from chaining the following three claims.

**Claim A.18.**

$$\begin{aligned} & \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{Q}_{f_X(x), f_Z(z)}^{x, z} | \hat{\psi} \rangle \\ & \approx_{\delta_Q^{1/2}} \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} \hat{M}_{f_X(x)}^{(\text{POINT}, X), x} | \hat{\psi} \rangle. \end{aligned} \quad (242)$$

*Proof.* Form the difference and apply the Cauchy Schwarz inequality to obtain

$$\begin{aligned} & \left| \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes (\hat{Q}_{f_X(x), f_Z(z)}^{x, z} - \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} \hat{M}_{f_X(x)}^{(\text{POINT}, X), x}) | \hat{\psi} \rangle \right| \\ & \leq \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes I | \hat{\psi} \right\|^2 \right)^{1/2} \\ & \quad \cdot \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes (\hat{Q}_{f_X(x), f_Z(z)}^{x, z} - \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} \hat{M}_{f_X(x)}^{(\text{POINT}, X), x}) | \hat{\psi} \right\|^2 \right)^{1/2}. \end{aligned}$$

The first term above is at most 1. For the second term, writing

$$W_{a, b}^{x, z} = (\hat{Q}_{a, b}^{x, z} - \hat{M}_b^{(\text{POINT}, Z), z} \hat{M}_a^{(\text{POINT}, X), x})$$

for short we bound it as

$$\begin{aligned} & \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes W_{f_X(x), f_Z(z)}^{x, z} | \hat{\psi} \right\|^2 \leq \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{a, b} \langle \hat{\psi} | I \otimes (W_{a, b}^{x, z})^2 | \hat{\psi} \rangle \\ & = \mathbb{E}_{(x, z)} \sum_{a, b} \langle \hat{\psi} | I \otimes (W_{a, b}^{x, z})^2 | \hat{\psi} \rangle \\ & \leq \delta_Q. \end{aligned}$$

Here the first inequality uses that for any  $x, z, a, b$ ,  $\sum_{f_X, f_Z: f_X(x)=a, f_Z(z)=b} T_{f_X, f_Z}^{\ell_X, \ell_Z} \leq I$ . In the equality in the before-last line, the expectation is over independent uniform  $x, z \in \mathbb{F}_q^m$ . The equality holds because for a line  $\ell$  sampled from the line-point distribution  $D$ , a uniformly random point  $(x, z, \alpha, \beta)$  on  $\ell$  is distributed as a uniformly random point in  $\mathbb{F}_q^{2m+2}$ . The last inequality is by (204).  $\square$

**Claim A.19.**

$$\begin{aligned} & \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} \hat{M}_{f_X(x)}^{(\text{POINT}, X), x} | \hat{\psi} \rangle \\ & \approx_{m \delta_{\text{LINE}}^{1/2}} \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), Z} | \hat{\psi} \rangle. \end{aligned} \quad (243)$$

*Proof.* We form the difference and apply the Cauchy-Schwarz inequality

$$\begin{aligned} & \left| \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} (I - \hat{M}_{f_X(x)}^{(\text{POINT}, X), x}) | \hat{\psi} \rangle \right| \\ & \leq \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} | \hat{\psi} \right\|^2 \right)^{1/2} \\ & \quad \cdot \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes (I - \hat{M}_{f_X(x)}^{(\text{POINT}, X), x}) | \hat{\psi} \right\|^2 \right)^{1/2}. \end{aligned}$$

The first term above is at most 1. The second term can be bounded as

$$\begin{aligned}
& \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes (I - \hat{M}_{f_X(x)}^{(\text{POINT}, X), x}) |\hat{\psi}\rangle \right\|^2 \\
&= \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_a \langle \hat{\psi} | T_a^{\ell_X, \ell_Z, x} \otimes (I - \hat{M}_a^{(\text{POINT}, X), x})^2 | \hat{\psi} \rangle \\
&= 1 - \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_a \langle \hat{\psi} | T_a^{\ell_X, \ell_Z, x} \otimes \hat{M}_a^{(\text{POINT}, X), x} | \hat{\psi} \rangle, \tag{244}
\end{aligned}$$

where we used the shorthand  $T_a^{\ell_X, \ell_Z, x} = \sum_{f_X, f_Z: f_X(x)=a} T_{f_X, f_Z}^{\ell_X, \ell_Z}$  and the second equality uses that  $\{\hat{M}_a^{(\text{POINT}, X), x}\}$  is a projective measurement. Using the definition

$$T_{f_X, f_Z}^{\ell_X, \ell_Z} = \hat{M}_{f_X}^{(\text{LINE}, X), \ell_X} \cdot \hat{M}_{f_Z}^{(\text{LINE}, Z), \ell_Z} \cdot \hat{M}_{f_X}^{(\text{LINE}, X), \ell_X}$$

and the fact that  $\{\hat{M}_{f_X}^{(\text{LINE}, X), \ell_X}\}$  is projective we get

$$1 - (244) = \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_a \langle \hat{\psi} | \hat{M}_{[\text{eval}_x(\cdot)=a]}^{(\text{LINE}, X), \ell_X} \otimes \hat{M}_a^{(\text{POINT}, X), x} | \hat{\psi} \rangle. \tag{245}$$

By Property 2 of Lemma A.17 the right-hand side of (245) is a mixture of terms of the form

$$\mathbb{E}_{(\ell_X, x) \sim D_{t,i}} \sum_a \langle \hat{\psi} | \hat{M}_{[\text{eval}_x(\cdot)=a]}^{(\text{LINE}, X), \ell_X} \otimes \hat{M}_a^{(\text{POINT}, X), x} | \hat{\psi} \rangle, \tag{246}$$

for  $t \in \{\text{ALINE}, \text{DLINE}\}$  and  $i \in \{1, \dots, m\}$ . Hence by the discussion following Definition A.16 and item 3 of Lemma A.10 it follows that

$$(246) = O(m^2 \delta_{\text{LINE}}).$$

□

**Claim A.20.**

$$\mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z} | \hat{\psi} \rangle \approx \sqrt{m} \cdot (\delta_P^{1/4} + \delta_Q^{1/4} + \varepsilon^{1/4}) \cdot 1. \tag{247}$$

*Proof.* At first we proceed similarly to the proof of Claim A.19, bounding the difference using Cauchy-Schwarz as

$$\begin{aligned}
& \left| \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \langle \hat{\psi} | T_{f_X, f_Z}^{\ell_X, \ell_Z} \otimes (I - \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z}) | \hat{\psi} \rangle \right| \\
&\leq \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes I | \hat{\psi} \rangle \right\|^2 \right)^{1/2} \\
&\quad \cdot \left( \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_{f_X, f_Z} \left\| \sqrt{T_{f_X, f_Z}^{\ell_X, \ell_Z}} \otimes (I - \hat{M}_{f_Z(z)}^{(\text{POINT}, Z), z}) | \hat{\psi} \rangle \right\|^2 \right)^{1/2}.
\end{aligned}$$



The first term is at most 1. The second, as in the proof of Claim A.19, equals

$$1 - \mathbb{E}_{(\ell, \ell_X, \ell_Z) \sim D} \mathbb{E}_{(x, z, \alpha, \beta) \in \ell} \sum_b \langle \hat{\psi} | T_b^{\ell_X, \ell_Z, z} \otimes \hat{M}_b^{(\text{POINT}, Z), z} | \hat{\psi} \rangle$$

$$= 1 - \mathbb{E}_{(\ell_X, \ell_Z) \sim D_{t_i} \times D_{t_j}} \mathbb{E}_{z \in \ell_Z} \sum_b \langle \hat{\psi} | T_b^{\ell_X, \ell_Z, z} \otimes \hat{M}_b^{(\text{POINT}, Z), z} | \hat{\psi} \rangle \quad (248)$$

$$= 1 - \mathbb{E}_{(\ell_X, \ell_Z) \sim D_{t_i} \times D_{t_j}} \mathbb{E}_{z \in \ell_Z} \mathbb{E}_{x \in \ell_X} \sum_{a, b} \langle \hat{\psi} | T_{[\text{eval}_x(\cdot)=a, \text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z} \otimes \hat{M}_b^{(\text{POINT}, Z), z} | \hat{\psi} \rangle \quad (249)$$

where we use the shorthand  $T_b^{\ell_X, \ell_Z, z} = \sum_{f_X} T_{f_X, [\text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z}$  and  $D_{t_i} \times D_{t_j}$  to denote a mixture over all such distributions where  $t \in \{\text{ALINE}, \text{DLINE}\}$  and  $i, j \in \{1, \dots, m\}$ . Equality holds in (248) because by item 2 of Lemma A.17 for  $(\ell, \ell_X, \ell_Z) \sim D$  the marginal distribution on  $(\ell_X, \ell_Z)$  is a mixture of the product distributions  $D_{\text{ALINE}, i} \times D_{\text{ALINE}, j}$  and  $D_{\text{DLINE}, i} \times D_{\text{DLINE}, j}$ , for  $i, j \in \{1, \dots, m\}$ . Moreover, conditioned on  $(\ell_X, \ell_Z)$ ,  $z$  is a uniformly random point on  $\ell_X$ . The last equality in (249) holds by definition of

$$\begin{aligned} T_b^{\ell_X, \ell_Z, z} &= \sum_{f_X} T_{f_X, [\text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z} \\ &= \mathbb{E}_{x \in \ell_X} \sum_a \sum_{f_X: f_X(x)=a} T_{f_X, [\text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z} \\ &= \mathbb{E}_{x \in \ell_X} \sum_a T_{[\text{eval}_x(\cdot)=a, \text{eval}_z(\cdot)=b]}^{\ell_X, \ell_Z} . \end{aligned}$$

We next write

$$\begin{aligned} 1 - (249) &\approx_{m(\sqrt{\delta_P} + \sqrt{\delta_Q})} \mathbb{E}_{(\ell_X, \ell_Z) \sim D_{t_i} \times D_{t_j}} \mathbb{E}_{x \in \ell_X} \mathbb{E}_{z \in \ell_Z} \sum_{a, b} \langle \hat{\psi} | \hat{M}_a^{(\text{POINT}, X), x} \hat{M}_b^{(\text{POINT}, Z), z} \otimes \hat{M}_b^{(\text{POINT}, Z), z} | \hat{\psi} \rangle \\ &= \sum_b \langle \hat{\psi} | \hat{M}_b^{(\text{POINT}, Z), z} \otimes \hat{M}_b^{(\text{POINT}, Z), z} | \hat{\psi} \rangle \\ &\geq 1 - \sqrt{\varepsilon} , \end{aligned}$$

where the first approximation follows from the discussion following Definition A.16 applied to the bounds from Lemma A.17 and Lemma A.11, the second line uses that  $\{\hat{M}_a^{(\text{POINT}, X), x}\}$  is a measurement, and the last is by self-consistency of the points measurements (Lemma A.9).  $\square$

Together, Claim A.18, Claim A.19 and Claim A.20 show (241). To complete the proof of (238), observe that by Property 2 of Lemma A.17 the right-hand side of Equation (241) is a mixture of terms of the form Together, Claim A.18, Claim A.19 and Claim A.20 show eq:qld-combined-lines-consistency. To complete the proof of (238), observe that by Property 2 of Lemma A.17 the right-hand side of Equation (241) is a mixture of terms of the form

$$\mathbb{E}_{(\ell_X, x) \sim D_{t_1, i}} \mathbb{E}_{(\ell_Z, z) \sim D_{t_2, j}} \sum_{f_X, f_Z} \langle \hat{\psi} | (T_{f_X, f_Z}^{\ell_X, \ell_Z})_{AA'} \otimes (I - (\hat{Q}_{f_X(x), f_Z(z)}^{x, z})) | \hat{\psi} \rangle$$

for  $t_1, t_2 \in \{\text{ALINE}, \text{DLINE}\}$  and  $i, j \in \{1, \dots, m\}$ . Hence using Equation (240) it follows that

$$(241) = O(m^2 \delta_P) .$$

This establishes the first relation (238). An analogous argument shows the second relation.  $\square$

**Lemma A.21.** *There exists a function  $\delta_S(\epsilon, m, d, q) = a(md)^a(\epsilon^b + q^{-b} + 2^{-bmd})$  for some universal constants  $a > 1, 0 < b < 1$  such that the following holds. For  $\mathcal{H} \in \{\mathcal{H}_A, \mathcal{H}_B\}$  there exists a projective measurement  $\{\hat{S}_{g_X, g_Z}\}$  acting on  $\mathcal{H} \otimes (\mathbb{C}^q)^{\otimes M}$  with outcomes consisting of pairs  $(g_X, g_Z)$  of polynomials each in  $\text{ideg}_{d,m}(\mathbb{F}_q)$  such that for  $W \in \{X, Z\}$ , on average over uniformly random  $u \in \mathbb{F}_q^m$ ,*

$$(\hat{S}_{[\text{eval}_u(\cdot_W)=a]})_{AA'} \simeq_{\delta_S} (\hat{M}_a^{(\text{POINT}, W), u})_{BA''} \quad (250)$$

$$(\hat{S}_{[\text{eval}_u(\cdot_W)=a]})_{BB'} \simeq_{\delta_S} (\hat{M}_a^{(\text{POINT}, W), u})_{AB''}, \quad (251)$$

where the notation  $\text{eval}_u(\cdot_W)$  means the evaluation of the first outcome  $g_X$  of  $\hat{S}$  if  $W = X$ , or the second outcome  $g_Z$  if  $W = Z$ , at the point  $u$ .

*Proof.* Lemma A.15 establishes the existence of a strategy  $\mathcal{S}_{\text{LD}} = (\hat{\psi}, \{\hat{Q}_f^\ell\} \cup \{\hat{Q}_a^{x,z,\alpha,\beta}\})$  for the classical low-degree test  $\mathfrak{G}_{\text{ldparams}}^{\text{LD}}$  for  $\text{ldparams} = (q, 2m + 2, d, 1)$ , with value  $1 - \delta_{\text{combine}}$ . This strategy is not necessarily projective because the measurement  $\{\hat{Q}_f^\ell\}$  is not guaranteed to be; however, we may projectivize it by applying Naimark's theorem to this measurement. This preserves the consistency ( $\simeq$ ) relation obtained as the conclusion of Lemma A.15. Theorem 7.8 applied to this projectivized strategy implies the existence of a POVM  $\{\hat{S}_g\}$  with outcomes  $g \in \text{ideg}_{d,2m+2}(\mathbb{F}_q)$  that is  $\delta_{\text{LD}}$ -self-consistent and consistent with the ‘‘points’’ measurements  $\{\hat{Q}_a^{x,z,\alpha,\beta}\}$ , where  $\delta_{\text{LD}} = a(md)^a(\delta_{\text{combine}}^b + q^{-b} + 2^{-bmd})$  for some universal constants  $a > 1, 0 < b < 1$  is given by Theorem 7.8. We may apply Naimark's theorem once again to  $\{\hat{S}_g\}$  to obtain a projective measurement with the same consistency guarantees, and henceforth, we will use  $\{\hat{S}_g\}$  to refer to this projective measurement.

We now argue that with high probability, the polynomial  $g$  returned by the measurement  $\{\hat{S}_g\}$  has the form  $g(x, z, \alpha, \beta) = \alpha g_X(x) + \beta g_Z(z)$  for polynomials  $g_X, g_Z \in \text{ideg}_{d,m}(\mathbb{F}_q)$ . Let  $\mathcal{G}$  denote the set of such polynomials. On average over uniformly random  $(x, z, \alpha, \beta)$ ,

$$\begin{aligned} (\hat{S}_g)_{AA'} &= (\hat{S}_g)_{AA'}^2 \\ &\approx_{\delta_{\text{LD}}} (\hat{S}_g)_{AA'} \otimes (\hat{Q}_{g(x,z,\alpha,\beta)}^{x,z,\alpha,\beta})_{BA''} \\ &\approx_{\delta_Q} (\hat{S}_g)_{AA'} (\hat{Q}_{g(x,z,\alpha,\beta)}^{x,z,\alpha,\beta})_{AA'} \\ &\approx_{\delta_Q} (\hat{S}_g)_{AA'} \otimes \left( \sum_{b,b'} 1_{\alpha b + \beta b' = g(x,z,\alpha,\beta)} \hat{M}_b^{(\text{POINT}, X), x} \hat{M}_{b'}^{(\text{POINT}, Z), z} \right)_{BA''} \end{aligned} \quad (252)$$

$$\approx_{\delta_Q} (\hat{S}_g)_{AA'} \otimes \left( \sum_{b,b'} 1_{\alpha b + \beta b' = g(x,z,\alpha,\beta)} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''}. \quad (253)$$

Here the first approximation uses the projectivity of  $\{\hat{S}_g\}$ , and the first approximation uses the consistency of  $\{\hat{S}_g\}$  with  $\{\hat{Q}_a^{x,z,\alpha,\beta}\}$  from Theorem 7.8, together with Fact 5.18 to switch from  $\simeq$  to  $\approx$ , and Fact 5.20. The second, third and fourth approximations follow from the consistency properties of  $\{\hat{Q}_a^{x,z,\alpha,\beta}\}$  from Lemma A.14 combined with Fact 5.20. Note that we will need both Equation (252) and Equation (253) for the subsequent calculations.

For all  $g$  define  $|\hat{\psi}_g\rangle = (\hat{S}_g)_{AA'}|\hat{\psi}\rangle$ . Then on average over uniformly random  $(x, z, \alpha, \beta)$ ,

$$\begin{aligned}
& \mathbb{E}_{x,z,\alpha,\beta} \left\| \left( \sum_{b,b'} \mathbf{1}_{\alpha b + \beta b' = g(x,z,\alpha,\beta)} \hat{M}_b^{(\text{POINT},X),x} \hat{M}_{b'}^{(\text{POINT},Z),z} \right)_{BA''} |\hat{\psi}_g\rangle \right\|^2 \\
&= \mathbb{E}_{x,z,\alpha,\beta} \sum_{b,b',b''} \mathbf{1}_{\alpha b + \beta b' = g(x,z,\alpha,\beta)} \mathbf{1}_{\alpha b + \beta b'' = g(x,z,\alpha,\beta)} \langle \hat{\psi}_g | \hat{M}_{b''}^{(\text{POINT},Z),z} \hat{M}_b^{(\text{POINT},X),x} \hat{M}_{b'}^{(\text{POINT},Z),z} | \hat{\psi}_g \rangle \\
&\approx_{1/q} \mathbb{E}_{x,z,\alpha,\beta} \sum_{b,b'} \mathbf{1}_{\alpha b + \beta b' = g(x,z,\alpha,\beta)} \langle \hat{\psi}_g | \hat{M}_{b'}^{(\text{POINT},Z),z} \hat{M}_b^{(\text{POINT},X),x} \hat{M}_{b'}^{(\text{POINT},Z),z} | \hat{\psi}_g \rangle \tag{254}
\end{aligned}$$

where in the first equality, we used the fact that  $\{\hat{M}_b^{(\text{POINT},X),x}\}$  is projective, and in the approximation, we used that for fixed  $x, z, \alpha, \beta$  with  $\beta \neq 0$ , if  $\alpha b + \beta b' = g(x, z, \alpha, \beta)$  and  $\alpha b + \beta b'' = g(x, z, \alpha, \beta)$ , then  $b' = b''$ . The probability that  $\beta = 0$  is  $1/q$ , and the term under the approximation has an absolute value of at most 1, so the error incurred is at most  $1/q$ .

We will now show that the right-hand side of Equation (254) is small for polynomials  $g$  that are not of the desired form. Fix  $b, b' \in \mathbb{F}_q$ , and let  $h_{b,b'}(x, z, \alpha, \beta) = g(x, z, \alpha, \beta) - (\alpha b + \beta b')$ . Observe that since  $g$  has individual degree  $d$ , so does  $h_{b,b'}$ . Write

$$h_{b,b'}(x, z, \alpha, \beta) = \sum_{i,j=0}^d h_{b,b',i,j}(x, z) \alpha^i \beta^j$$

for some polynomials  $h_{b,b',i,j}(x, z)$  of individual degree at most  $d$ . Let  $\mathcal{G}' \subset \text{ideg}_{d,2m+2}(\mathbb{F}_q)$  denote the polynomials that are linear in  $\alpha, \beta$ , i.e.,  $g = \alpha g_1(x, z) + \beta g_2(x, z)$ . Fix a  $g \notin \mathcal{G}'$ . Then  $h_{b,b'}(x, z, \alpha, \beta)$  is a nonzero polynomial of total degree at most  $(2m+2)d$ , and in particular there must exist an  $(i, j)$  such that the polynomial  $h_{b,b',i,j}(x, z)$  is not the identically zero polynomial. Call a pair  $(x, z) \in \mathbb{F}_q^2$  *good* if  $h_{b,b'}(x, z, \alpha, \beta)$  is a nonzero polynomial function of  $\alpha, \beta$ , and *bad* otherwise. The probability that  $(x, z)$  is bad is at most  $(2m+2)d/q$  by the Schwartz-Zippel lemma. Otherwise, conditioned on a good  $(x, z)$  pair, the probability that  $h_{b,b'}(x, z, \alpha, \beta) = 0$  over the choice of  $\alpha, \beta$  is at most  $(2m+2)d/q$ , again by the Schwartz-Zippel lemma. Thus for  $g \notin \mathcal{G}'$  we can upper-bound (254) by  $\frac{2(2m+2)d}{q} \|\hat{\psi}_g\|^2$ . Combined with (252) we get that

$$\sum_{g \notin \mathcal{G}'} \left\| (\hat{S}_g)_{AA'} |\hat{\psi}\rangle \right\|^2 \leq O(\delta_{\text{LD}} + \delta_Q + md/q). \tag{255}$$

Next, we argue that not only is  $g$  linear in  $\alpha, \beta$  with high probability, but also that  $g = \alpha g_X + \beta g_Z$  where  $g_X$  depends on  $x$  only and  $g_Z$  depends on  $z$  only. Fix a polynomial  $g \in \mathcal{G}'$ , i.e.,  $g(x, z, \alpha, \beta) = \alpha g_1(x, z) + \beta g_2(x, z)$  for polynomials  $g_1, g_2 \in \text{ideg}_{d,2m}(\mathbb{F}_q)$ . For all  $x, z \in \mathbb{F}_q^m$ ,  $\alpha, \beta, b, b' \in \mathbb{F}_q$ ,

$$\mathbf{1}_{\alpha b + \beta b' = \alpha g_1(x,z) + \beta g_2(x,z)} = \mathbf{1}_{b=g_1(x,z)} \mathbf{1}_{b'=g_2(x,z)} + \mathbf{1}_{b \neq g_1(x,z) \vee b' \neq g_2(x,z)} \cdot \mathbf{1}_{\alpha b + \beta b' = \alpha g_1(x,z) + \beta g_2(x,z)}.$$

By the Schwartz-Zippel lemma,

$$\mathbb{E}_{\alpha,\beta} \mathbf{1}_{b \neq g_1(x,z) \vee b' \neq g_2(x,z)} \cdot \mathbf{1}_{\alpha b + \beta b' = \alpha g_1(x,z) + \beta g_2(x,z)} \leq 1/q. \tag{256}$$

Let  $\mathcal{G}' = \mathcal{G}'_X \cup \mathcal{G}'_Z \cup \mathcal{G}$ , where  $\mathcal{G}'_X$  (resp.  $\mathcal{G}'_Z$ ) is the set of polynomials  $g \in \mathcal{G}'$  where  $g_1(x, z)$  depends

nontrivially on  $z$  (resp.  $g_2(x, z)$  depends nontrivially on  $x$ ). Fix  $g \in \mathcal{G}'_Z$ . We can upper-bound (254) by

$$\frac{1}{q} \|\hat{\psi}_g\rangle\|^2 + \mathbb{E}_{x,z} \sum_{b,b'} \mathbf{1}_{b=g_1(x,z)} \mathbf{1}_{b'=g_2(x,z)} \langle \hat{\psi}_g | \hat{M}_{b'}^{(\text{POINT},Z),z} \hat{M}_b^{(\text{POINT},X),x} \hat{M}_{b'}^{(\text{POINT},Z),z} | \hat{\psi}_g \rangle \quad (257)$$

$$\leq \frac{1}{q} \|\hat{\psi}_g\rangle\|^2 + \mathbb{E}_z \sum_{b'} \left( \mathbb{E}_x \mathbf{1}_{b'=g_2(x,z)} \right) \langle \hat{\psi}_g | \hat{M}_{b'}^{(\text{POINT},Z),z} | \hat{\psi}_g \rangle \quad (258)$$

where the indicator  $\mathbf{1}_{b=g_1(x,z)}$  is removed using positivity in the inequality. To bound the second term in (258), we use an argument similar to the one we used to derive (255). That is, say that a value of  $z$  is *good* if  $g_2(\cdot, z)$  is not a constant function. The probability that  $z$  is good is at least  $1 - (2m+2)d/q$  by the Schwartz-Zippel lemma and the assumption that  $g \in \mathcal{G}'_Z$  (and so  $g_2$  depends nontrivially on  $x$ ). Moreover, for each good  $z$  and for each  $b'$  the expectation  $\mathbb{E}_x \mathbf{1}_{b'=g_2(x,z)}$  is at most  $(2m+2)d/q$ . Hence, we conclude that if  $g_2(x, z)$  depends on  $x$ , then (258) is at most  $\frac{2(2m+2)d+1}{q} \|\hat{\psi}_g\rangle\|^2$ , and thus

$$\sum_{g \in \mathcal{G}'_Z} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 \leq O(\delta_{LD} + \delta_Q + md/q). \quad (259)$$

By starting with (253) instead (i.e., the order of  $M_{b'}^{(\text{POINT},Z),z}$  and  $M_b^{(\text{POINT},X),x}$  are switched), we can perform similar reasoning to deduce that

$$\sum_{g \in \mathcal{G}'_X} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 \leq O(\delta_{LD} + \delta_Q + md/q). \quad (260)$$

We thus obtain

$$\sum_{g \notin \mathcal{G}} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 \leq \sum_{g \notin \mathcal{G}'} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 + \sum_{g \in \mathcal{G}'_X} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 + \sum_{g \in \mathcal{G}'_Z} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 \quad (261)$$

$$\leq O(\delta_{LD} + \delta_Q + md/q). \quad (262)$$

Let  $\delta_G$  be the error in the right hand side of Equation (262). Define a projective sub-measurement  $\{\hat{S}_{g_X, g_Z}\}$  with outcomes  $g_X, g_Z \in \text{id eg}_{d,m}(\mathbb{F}_q)$  by

$$\hat{S}_{g_X, g_Z} = \hat{S}_{\text{combine}_{g_X, g_Z}}. \quad (263)$$

A bound on the incompleteness of  $\hat{S}_{g_X, g_Z}$  follows from (262) and the projectivity of  $\hat{S}_g$ :

$$1 - \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'} | \hat{\psi} \rangle = \sum_{g \notin \mathcal{G}} \langle \hat{\psi} | (\hat{S}_g)_{AA'} | \hat{\psi} \rangle = \sum_{g \notin \mathcal{G}} \|(\hat{S}_g)_{AA'} | \hat{\psi} \rangle\|^2 \leq \delta_G. \quad (264)$$

We now bound the consistency of  $\{\hat{S}_{g_X, g_Z}\}$  with the points measurements  $\{\hat{M}_a^{(\text{POINT}, W), u}\}$ .

$$1 = \sum_g \langle \hat{\psi} | (\hat{S}_g)_{AA'} | \hat{\psi} \rangle \quad (265)$$

$$\approx_{\delta_G} \sum_{g \in \mathcal{G}} \langle \hat{\psi} | (\hat{S}_g)_{AA'} | \hat{\psi} \rangle \quad (266)$$

$$\approx_{(\delta_Q)^{1/2}} \mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'} \otimes \left( \sum_{b, b'} 1_{\alpha b + \beta b' = \alpha_{g_X}(x) + \beta_{g_Z}(z)} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''} | \hat{\psi} \rangle \quad (267)$$

$$\approx_{(\delta_Q)^{1/2}} \mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'} \otimes \left( \sum_{b, b', b''} 1_{\alpha b + \beta b' = \alpha_{g_X}(x) + \beta_{g_Z}(z) = \alpha b'' + \beta b'} \hat{M}_{b''}^{(\text{POINT}, X), x} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''} | \hat{\psi} \rangle \quad (268)$$

$$\approx_{1/q} \mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'} \otimes \left( \sum_{b, b'} 1_{\alpha b + \beta b' = \alpha_{g_X}(x) + \beta_{g_Z}(z)} \hat{M}_b^{(\text{POINT}, X), x} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''} | \hat{\psi} \rangle \quad (269)$$

$$\approx_{1/q} \mathbb{E}_{x,z} \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'} \otimes \left( \hat{M}_{g_X(x)}^{(\text{POINT}, X), x} \hat{M}_{g_Z(z)}^{(\text{POINT}, Z), z} \hat{M}_{g_X(x)}^{(\text{POINT}, X), x} \right)_{BA''} | \hat{\psi} \rangle. \quad (270)$$

Here Equation (266) follows from (264). Equation (267) follows by approximation (253) together with the Cauchy-Schwarz inequality and the projectivity of  $\hat{S}_g$ :

$$\begin{aligned} & \left| \mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \langle \hat{\psi} | \left( (\hat{S}_{g_X, g_Z})_{AA'} \cdot \left( I - \left( \sum_{b, b'} 1_{\alpha b + \beta b' = \alpha_{g_X}(x) + \beta_{g_Z}(z)} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''} \right) | \hat{\psi} \rangle \right) \right| \\ & \leq \sqrt{\mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \langle \hat{\psi} | (\hat{S}_{g_X, g_Z})_{AA'}^2 | \hat{\psi} \rangle} \\ & \quad \cdot \sqrt{\mathbb{E}_{x,z,\alpha,\beta} \sum_{g_X, g_Z} \left\| \left( (\hat{S}_{g_X, g_Z})_{AA'} \otimes \left( I - \left( \sum_{b, b'} 1_{\alpha b + \beta b' = \alpha_{g_X}(x) + \beta_{g_Z}(z)} \hat{M}_{b'}^{(\text{POINT}, Z), z} \hat{M}_b^{(\text{POINT}, X), x} \right)_{BA''} \right) | \hat{\psi} \rangle \right\|^2} \\ & \leq 1 \cdot \sqrt{\delta_Q}. \end{aligned}$$

Equation (268) follows by a similar calculation. To pass from here to Equation (269), we argue that for  $\alpha \neq 0$ , the indicator forces  $b = b''$ , and the terms where  $\alpha = 0$  are bounded in absolute value by  $1/q$ . Finally, to reach Equation (270), we apply Equation (256) to the expectation of the indicator over  $\alpha, \beta$ . Equation (270) tells us that

$$\left( \hat{S}_{[\text{eval}_x(\cdot)=a, \text{eval}_z(\cdot)=b]} \right)_{AA'} \simeq_{O(\delta_G + (\delta_Q)^{1/2} + 1/q)} \left( \hat{M}_a^{(\text{POINT}, X), x} \hat{M}_b^{(\text{POINT}, Z), z} \hat{M}_a^{(\text{POINT}, X), x} \right)_{BA''}. \quad (271)$$

This is almost the desired conclusion of the lemma. The only issue is that  $\{\hat{S}_{g_X, g_Z}\}$  is a sub-measurement. To address this, we complete  $\hat{S}_{g_X, g_Z}$  by adding  $1 - \sum_{g_X, g_Z} \hat{S}_{g_X, g_Z}$  to an arbitrary measurement element. By Equation (264), it follows that the consistency relation in Equation (271) holds for the completed measurement with an error increased by  $\delta_G$ . Let  $\delta_S$  be this new, increased, error, and observe that  $\delta_S = O(\delta_G + (\delta_Q)^{1/2} + d/q)$ . Substituting in the bound on  $\delta_G$  from Equation (262), we obtain that  $\delta_S =$

$O(\delta_{LD} + (\delta_Q)^{1/2} + md/q)$  has the form  $a'(md)^{a'}(\varepsilon^{b'} + q^{-b'} + 2^{-b'md})$  for some universal constants  $a' > 1, 0 < b' < 1$ . We thus conclude by an application of Fact 5.24 that Equation (250) holds for  $W = X$ .

To obtain the same result for  $W = Z$ , we perform exactly the same steps starting from Equation (265) but with  $X$  and  $Z$  interchanged (so we use Equation (252) instead of Equation (253)). Finally, entirely analogous calculations with the registers swapped yield Equation (251).  $\square$

## A.6 Pulling the $X$ and $Z$ measurements apart

Recall the decoding map of the low-degree code defined in Section 3.4. This map takes in a polynomial  $g \in \text{ideg}_{d,m}(\mathbb{F}_q)$  and returns a string  $\text{Dec}(g) \in \mathbb{F}_q^M$  consisting of the evaluations of  $g$  on the points in the hypercube  $\{0, 1\}^m$ . We now use this decoding map to construct a new set of measurement operators out of  $\hat{S}$ . First, we introduce some notation. For all  $W \in \{X, Z\}$  and all degree  $d$  polynomials  $g : \mathbb{F}_q^m \rightarrow \mathbb{F}_q$  define projectors

$$\hat{S}_g^X = \sum_{gZ} \hat{S}_{g,gZ} \quad \text{and} \quad \hat{S}_g^Z = \sum_{gX} \hat{S}_{gX,g} ,$$

where the measurement  $\{\hat{S}_{gX,g}\}$  is given by Lemma A.21.

For all  $W \in \{X, Z\}$ ,  $\tilde{u} \in \mathbb{F}_q^M$ , and  $a \in \mathbb{F}_q$ , define the projector

$$\tau_a^W(\tilde{u}) = \sum_{h \in \mathbb{F}_q^M : h \cdot \tilde{u} = a} \tau_h^W \quad (272)$$

where recall that  $\tau_h^W$  is the projector corresponding to measuring  $(\mathbb{C}^q)^{\otimes M}$  in the  $W$  basis and obtaining  $h \in \mathbb{F}_q^M$  as the outcome. For all  $\tilde{u} \in \mathbb{F}_q^M$  define the projective measurement  $\{\tilde{M}_a^{W,\tilde{u}}\}_{a \in \mathbb{F}_q}$  by

$$\tilde{M}_a^{W,\tilde{u}} = \sum_g \hat{S}_g^W \otimes \tau_{(\text{Dec}(g) \cdot \tilde{u}) - a}^W(\tilde{u}) . \quad (273)$$

If  $\hat{S}_g^W$  is viewed as an operator acting on registers  $AA'$  (resp.  $BB'$ ) then we view  $\tilde{M}_a^{W,\tilde{u}}$  as an operator acting on registers  $AA'A''$  (resp.  $BB'B''$ ). (When necessary we will indicate via subscripts which registers the operators are acting on.) The  $\tilde{M}_a^{W,\tilde{u}}$  are projective because the  $\hat{S}_g^W$  and  $\tau^W$  operators are projective.

For all  $j \in \{1, \dots, t\}$  define the observable

$$\tilde{W}^j(\tilde{u}) = \sum_a (-1)^{\text{tr}(e_j a)} \tilde{M}_a^{W,\tilde{u}} , \quad (274)$$

where  $\{e_j\}_{j \in \{1, \dots, t\}}$  denotes the self-dual basis for  $\mathbb{F}_q$  over  $\mathbb{F}_2$  specified in Section 3.3.1. Let  $\tau^W$  be the generalized Pauli observable defined in Section 3.7. Observe that

$$\begin{aligned} \left( \sum_{gX,gZ} (-1)^{\text{tr}(e_j(\text{Dec}(g_W) \cdot \tilde{u}))} \hat{S}_{gX,gZ} \right) \otimes \tau^W(e_j \tilde{u}) &= \sum_{gX,gZ} \sum_h (-1)^{\text{tr}(e_j(\text{Dec}(g_W) \cdot \tilde{u}))} (-1)^{\text{tr}(e_j(\tilde{u} \cdot h))} \hat{S}_{gX,gZ} \otimes \tau_h^W \\ &= \sum_{g_W} \sum_a (-1)^{\text{tr}(e_j(\tilde{u} \cdot \text{Dec}(g_W) + a))} \hat{S}_{g_W}^W \otimes \left( \sum_{h: h \cdot \tilde{u} = a} \tau_h^W \right) \\ &= \sum_{g_W} \sum_a (-1)^{\text{tr}(e_j(\tilde{u} \cdot \text{Dec}(g_W) + a))} \hat{S}_{g_W}^W \otimes \tau_a^W(\tilde{u}) \\ &= \tilde{W}^j(\tilde{u}) , \end{aligned}$$

where the first equality uses the definition of  $\tau^W(e_j\tilde{u})$ , the second uses the definition of  $\hat{S}_{g^W}^W$  and regroups terms, the third uses the definition (272) of  $\tau_a^W(\tilde{u})$ , and the last is by a change of variables. These equations in particular show that the observable  $\tilde{W}^j(\tilde{u})$  is Hermitian and squares to the identity, because the measurement  $\{\hat{S}_{g^X, g^Z}\}$  is projective. Furthermore, it can be verified by direct calculation that the observables  $\tilde{W}^j(\tilde{u})$  satisfy the same relations as the generalized Pauli group. Specifically, for all  $j, j' \in \{1, \dots, t\}$ ,  $\tilde{u}, \tilde{v} \in \mathbb{F}_q^M$ ,

$$\tilde{X}^j(\tilde{u})\tilde{Z}^{j'}(\tilde{v}) = \begin{cases} \tilde{Z}^{j'}(\tilde{v})\tilde{X}^j(\tilde{u}) & j \neq j' \\ (-1)^{\text{tr}((e_j\tilde{u}) \cdot (e_{j'}\tilde{v}))} \tilde{Z}^{j'}(\tilde{v})\tilde{X}^j(\tilde{u}) & j = j' \end{cases}.$$

The next lemma shows that the measurements  $\{\tilde{M}_a^{W, \tilde{u}}\}$  and observables  $\tilde{W}^j(\tilde{u})$  are consistent with the original points measurements, and are self-consistent.

**Lemma A.22.** *Let  $\delta_S$  as in Lemma A.21. For all  $W \in \{X, Z\}$ , all  $\tilde{u} \in \mathbb{F}_q^M$  and  $j \in \{1, \dots, t\}$ , the measurements  $\{\tilde{M}_a^{W, \tilde{u}}\}_{a \in \mathbb{F}_q}$  and observables  $\tilde{W}^j(\tilde{u})$  satisfy the following properties:*

1. **(Consistency with points measurements)** *For all  $W \in \{X, Z\}$ , for all  $j \in \{1, \dots, t\}$ , and on average over uniformly random  $u \in \mathbb{F}_q^m$ ,*

$$I_{AA'A''} \otimes \tilde{M}_a^{W, \text{ind}_m(u)} \simeq_{\delta_S} M_a^{(\text{POINT}, W), u} \otimes I_{BB'B''}$$

and

$$\tilde{M}_a^{W, \text{ind}_m(u)} \otimes I_{BB'B''} \simeq_{\delta_S} I_{AA'A''} \otimes M_a^{(\text{POINT}, W), u}.$$

2. **(Self-consistency)** *On average over uniformly random  $\tilde{u} \in \mathbb{F}_q^M$ ,*

$$\tilde{W}^j(\tilde{u}) \otimes I_{BB'B''} \approx_{\delta_S} I_{AA'A''} \otimes \tilde{W}^j(\tilde{u}).$$

*Proof.* We show the first item of the lemma for the case when  $M_a^{(\text{POINT}, W), u}$  acts on register A and  $\tilde{M}_a^{W, \text{ind}_m(u)}$  acts on registers  $BB'B''$  (the argument for when the A and B registers are interchanged proceeds analogously):

$$\begin{aligned} & \mathbb{E}_{u \in \mathbb{F}_q^m} \sum_a \langle \hat{\psi} | M_a^{(\text{POINT}, W), u} \otimes \tilde{M}_a^{W, \text{ind}_m(u)} | \hat{\psi} \rangle \\ &= \mathbb{E}_{u \in \mathbb{F}_q^m} \sum_{g, a} \langle \hat{\psi} | (M_a^{(\text{POINT}, W), u})_A \otimes (\hat{S}_g^W)_{BB'} \otimes (\tau_{\text{Dec}(g) \cdot \text{ind}_m(u) - a}^W)_{B''} | \hat{\psi} \rangle \\ &= \mathbb{E}_{u \in \mathbb{F}_q^m} \sum_g \langle \hat{\psi} | (\hat{M}_{\text{Dec}(g) \cdot \text{ind}_m(u)}^{(\text{POINT}, W), u})_{AB''} \otimes (\hat{S}_g^W)_{BB'} | \hat{\psi} \rangle \\ &= \mathbb{E}_{u \in \mathbb{F}_q^m} \sum_g \langle \hat{\psi} | (\hat{M}_{g(u)}^{(\text{POINT}, W), u})_{AB''} \otimes (\hat{S}_g^W)_{BB'} | \hat{\psi} \rangle \\ &\geq 1 - \delta_S. \end{aligned}$$

The second line follows from expanding the definition of  $\tilde{M}_a^{W, \text{ind}_m(u)}$  using (273), the third line follows from the definition of  $\hat{M}_a^{(\text{POINT}, W), u}$  given in Section A.3, the fourth line follows from the fact that  $\text{Dec}(g) \cdot \text{ind}_m(u) = g(u)$ , and the fifth line follows from Lemma A.21. This implies the first item of the Lemma.

To show the second item we first argue that, on average over  $\tilde{u}$ ,

$$\left( \tilde{M}_a^{W,\tilde{u}} \right)_{AA'A''} \approx_{\delta_S} \left( \tilde{M}_a^{W,\tilde{u}} \right)_{BB'B''}. \quad (275)$$

This follows because on average over  $u \in \mathbb{F}_q^m$  and  $\tilde{u} \in \mathbb{F}_q^M$  we have

$$\left( \tilde{M}_a^{W,\tilde{u}} \right)_{AA'A''} \quad (276)$$

$$\approx_{\delta_S} \left( \tilde{M}_a^{W,\tilde{u}} \right)_{AA'A''} \cdot \left( \sum_g (\hat{S}_g^W)_{AA'} \otimes (\hat{M}_{g(u)}^{(\text{POINT},W),u})_{BA''} \right) \quad (277)$$

$$= \sum_g \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_{(\text{Dec}(g)\cdot\tilde{u})-a}^W(\tilde{u}))_{A''} \right) \cdot \left( \sum_{r,s:r+s=g(u)} (M_r^{(\text{POINT},W),u})_B \otimes (\tau_s^W(\text{ind}_m(u)))_{A''} \right) \quad (278)$$

$$= \sum_{\substack{g,h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \otimes (M_{(g-g_h)(u)}^{(\text{POINT},W),u})_B. \quad (279)$$

The first inequality follows from right-multiplying (276) by the approximation (297) in Lemma A.23 (to be proved below) and using Fact 5.19. Line (278) follows from expanding the definition of  $\tilde{M}_a^{W,\tilde{u}}$ . To obtain Line (279), we expanded the definitions of  $\tau_{(\text{Dec}(g)\cdot\tilde{u})-a}^W(\tilde{u})$  and  $\tau_s^W(\text{ind}_m(u))$  using Eq. (272), and the fact that  $h \cdot \text{ind}_m(u) = g_h(u)$  (due to Eq. (13)). Thus for all  $g \in \text{ideg}_{d,m}(\mathbb{F}_q)$  and  $s \in \mathbb{F}_q$  we have

$$\tau_{(\text{Dec}(g)\cdot\tilde{u})-a}^W(\tilde{u}) \cdot \tau_s^W(\text{ind}_m(u)) = \sum_{\substack{h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a \\ g_h(u)=s}} \tau_h^W.$$

Using the self-consistency of  $\{M_a^{(\text{POINT},W),u}\}$  shown in Lemma A.7, we then obtain

$$(279) \approx_\varepsilon \sum_{\substack{g,h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} (\hat{S}_g^W)_{AA'} \cdot \left( (M_{(g-g_h)(u)}^{(\text{POINT},W),u})_A \otimes (\tau_h^W)_{A''} \right) \cdot \left( M_{(g-g_h)(u)}^{(\text{POINT},W),u} \right)_B \quad (280)$$

$$\approx_0 \sum_{\substack{g,h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} (\hat{S}_g^W)_{AA'} \cdot \left( (M_{(g-g_h)(u)}^{(\text{POINT},W),u})_A \otimes (\tau_h^W)_{A'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( M_{(g-g_h)(u)}^{(\text{POINT},W),u} \right)_B \quad (281)$$

$$= \sum_{\substack{g,h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} (\hat{S}_g^W)_{AA'} \cdot \left( (\hat{M}_{g(u)}^{(\text{POINT},W),u})_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( M_{(g-g_h)(u)}^{(\text{POINT},W),u} \right)_B. \quad (282)$$

The “ $\approx_0$ ” in Line (281) indicates that the left hand side is equal to the right hand side when applied to the state  $|\hat{\psi}\rangle$ . Line (281) follows from the self-consistency of the operators  $\{\tau_h^W\}$ , and Line (282) follows from the definition of  $\{\hat{M}_a^{(\text{POINT},W),u}\}$ . Using the fact that  $|\hat{\psi}\rangle = \sum_{h'} (\tau_{h'}^W)_{B'} \otimes (\tau_{h'}^W)_{B''} |\hat{\psi}\rangle$ , we get that

$$(282) \approx_0 \sum_{\substack{g,h: \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} \left( (\hat{S}_g^W \cdot \hat{M}_{g(u)}^{(\text{POINT},W),u})_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( M_{(g-g_h)(u)}^{(\text{POINT},W),u} \otimes \sum_{h'} \tau_{h'}^W \otimes \tau_{h'}^W \right)_{BB'B''} \quad (283)$$

$$= \sum_{\substack{g,h,h': \\ (\text{Dec}(g)-h)\cdot\tilde{u}=a}} \left( (\hat{S}_g^W \cdot \hat{M}_{g(u)}^{(\text{POINT},W),u})_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{M}_{(g-g_h+g_{h'})(u)}^{(\text{POINT},W),u})_{BB'} \otimes (\tau_{h'}^W)_{B''} \right). \quad (284)$$



Line (284) follows from (283) by expanding the definition of  $\{\hat{M}_a^{(\text{POINT},W),u}\}$ . Next, we argue that, on average over  $u \in \mathbb{F}_q^m$ ,

$$(284) \approx_{\delta_S} \sum_{\substack{g,h,h': \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \otimes \left( (\hat{M}_{(g-g_h+g_{h'})}^{(\text{POINT},W),u})_{BB'} \otimes (\tau_{h'}^W)_{B''} \right). \quad (285)$$

We show this by bounding the magnitude of the difference. Using the fact that  $\hat{S}_g^W$  and  $\tau_h^W$  are projective, we have:

$$\begin{aligned} & \mathbb{E}_u \sum_a \left\| \sum_{\substack{g,h,h': \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} (\hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}))_{AA'} \otimes (\tau_h^W)_{A''} \otimes (\hat{M}_{(g-g_h+g_{h'})}^{(\text{POINT},W),u})_{BB'} \otimes (\tau_{h'}^W)_{B''} | \hat{\psi} \right\|^2 \\ &= \mathbb{E}_u \sum_{g,h,h'} \langle \hat{\psi} | \left( (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}) \cdot \hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}) \right)_{AA'} \otimes (\tau_h^W)_{A''} \\ & \quad \cdot \left( (\hat{M}_{(g-g_h+g_{h'})}^{(\text{POINT},W),u})_{BB'}^2 \otimes (\tau_{h'}^W)_{B''} \right) | \hat{\psi} \rangle \\ &\leq \mathbb{E}_u \sum_g \langle \hat{\psi} | \left( (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}) \cdot \hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}) \right)_{AA'} | \hat{\psi} \rangle \quad (286) \\ &\leq \delta_S. \quad (287) \end{aligned}$$

The inequality in Line (286) follows from the fact that for all  $g, h$ , the sum  $\sum_{h'} (\hat{M}_{(g-g_h+g_{h'})}^{(\text{POINT},W),u})_{BB'}^2 \otimes (\tau_{h'}^W)_{B''}$  is at most  $I$ . The inequality in Line (287) follows from the second item of Lemma A.23 (to be proved below). Next, we show that on average over  $u \in \mathbb{F}_q^m$ ,

$$\begin{aligned} (285) &= \sum_{\substack{g,h,g',h': \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{S}_{g'}^W \cdot \hat{M}_{(g-g_h+g_{h'})}^{(\text{POINT},W),u})_{BB'} \otimes (\tau_{h'}^W)_{B''} \right) \quad (288) \\ &\approx_{O(\delta_S + \sqrt{\varepsilon})} \sum_{\substack{g,h,g',h': \\ (g'-g_{h'})(u)=(g-g_h)(u) \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{S}_{g'}^W \cdot \hat{M}_{g'(u)}^{(\text{POINT},W),u})_{BB'} \otimes (\tau_{h'}^W)_{B''} \right) \quad (289) \end{aligned}$$

$$\approx_{\delta_S} \sum_{\substack{g,h,g',h': \\ (g'-g_{h'})(u)=(g-g_h)(u) \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{S}_{g'}^W)_{BB'} \otimes (\tau_{h'}^W)_{B''} \right) \quad (290)$$

$$\approx_{md/q} \sum_{\substack{g,h,g',h': \\ g'-g_{h'}=g-g_h \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{S}_{g'}^W)_{BB'} \otimes (\tau_{h'}^W)_{B''} \right). \quad (291)$$

Before we justify the sequence of steps to go from Line (285) to Line (291), we first argue how this establishes Line (275), and hence the second item of the Lemma statement.

Absorbing the error terms  $O(md/q)$  and  $O(\sqrt{\varepsilon})$  into  $\delta_S$ , we have shown that

$$\left( \tilde{M}_a^{W,\bar{u}} \right)_{AA'A''} \approx_{\delta_S} \sum_{\substack{g,h,g',h': \\ g'-g_{h'}=g-g_h \\ (\text{Dec}(g)-h) \cdot \bar{u}=a}} \left( (\hat{S}_g^W)_{AA'} \otimes (\tau_h^W)_{A''} \right) \cdot \left( (\hat{S}_{g'}^W)_{BB'} \otimes (\tau_{h'}^W)_{B''} \right).$$

Note that the conditions  $(\text{Dec}(g) - h) \cdot \tilde{u} = a$  and  $g - g_h = g' - g_{h'}$  are equivalent to the conditions  $(\text{Dec}(g') - h') \cdot \tilde{u} = a$  and  $g - g_h = g' - g_{h'}$ . Thus the expression in (291) is symmetric between  $g, h$  and  $g', h'$ , and so an analogous derivation shows that  $(\tilde{M}_a^{W, \tilde{u}})_{\text{BB}'\text{B}''}$  is  $\delta_S$ -close to (291), and thus  $\delta_S$ -close to  $(\tilde{M}_a^{W, \tilde{u}})_{\text{AA}'\text{A}''}$ . This shows (275). Since  $\{\tilde{M}_a^{W, \tilde{u}}\}$  is projective, by using Item 2 of Fact 5.18, followed by Fact 5.24, and then followed by Item 1 of Fact 5.18, we get for all  $j \in \{1, \dots, t\}$ , on average over  $\tilde{u} \in \mathbb{F}_q^M$ ,

$$(\tilde{M}_{[\text{tr}(e_j)=b]}^{W, \tilde{u}})_{\text{AA}'\text{A}''} \approx_{\delta_S} (\tilde{M}_{[\text{tr}(e_j)=b]}^{W, \tilde{u}})_{\text{BB}'\text{B}''}$$

where the answer summation is over  $b \in \mathbb{F}_2$ . Finally, the second item of the Lemma follows from applying Lemma A.5.

We now return to justifying the sequence of steps between Lines (285) and (291). The equality in Line (288) follows by left-multiplying (285) by  $I_{\text{BB}'} = \sum_{g'} (\hat{S}_{g'}^W)_{\text{BB}'}$ .

**The approximation in Line (289).** The approximation in Line (289) follows by bounding the magnitude of the difference:

$$\begin{aligned} & \mathbb{E}_u \sum_a \left\| \sum_{\substack{g, h, g', h': \\ (g' - g_{h'})(u) \neq (g - g_h)(u) \\ (\text{Dec}(g) - h) \cdot \tilde{u} = a}} \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{S}_{g'}^W \cdot \hat{M}_{(g - g_h + g_{h'})(u)}^{(\text{POINT}, W), \mu})_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) |\hat{\psi}\rangle \right\|^2 \\ &= \mathbb{E}_u \sum_{\substack{g, h, g', h': \\ (g' - g_{h'})(u) \neq (g - g_h)(u)}} \langle \hat{\psi} | \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{M}_{(g - g_h + g_{h'})(u)}^{(\text{POINT}, W), \mu} \cdot \hat{S}_{g'}^W \cdot \hat{M}_{(g - g_h + g_{h'})(u)}^{(\text{POINT}, W), \mu})_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) |\hat{\psi}\rangle \\ &\leq \mathbb{E}_u \sum_{\substack{g, h, g', h', a': \\ a' \neq g'(u)}} \langle \hat{\psi} | \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{M}_{a'}^{(\text{POINT}, W), \mu} \cdot \hat{S}_{g'}^W \cdot \hat{M}_{a'}^{(\text{POINT}, W), \mu})_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) |\hat{\psi}\rangle \\ &= \mathbb{E}_u \sum_{g, a: a \neq g(u)} \langle \hat{\psi} | (\hat{M}_a^{(\text{POINT}, W), \mu} \cdot \hat{S}_g^W \cdot \hat{M}_a^{(\text{POINT}, W), \mu})_{\text{BB}'} |\hat{\psi}\rangle \tag{292} \\ &\approx_{O(\sqrt{\varepsilon})} \mathbb{E}_u \sum_g \langle \hat{\psi} | (I - \hat{M}_{g(u)}^{(\text{POINT}, W), \mu})_{\text{AB}''} \otimes (\hat{S}_g^W)_{\text{BB}'} |\hat{\psi}\rangle \tag{293} \\ &\leq \delta_S. \tag{294} \end{aligned}$$

The approximation in Line (293) follows from the following calculation:

$$\begin{aligned} & \left| (293) - (292) \right| \\ &\leq \left| \mathbb{E}_u \sum_{g, a: a \neq g(u)} \langle \hat{\psi} | (\hat{M}_a^{(\text{POINT}, W), \mu} \cdot \hat{S}_g^W)_{\text{BB}'} \cdot ((\hat{M}_a^{(\text{POINT}, W), \mu})_{\text{AB}''} - (\hat{M}_a^{(\text{POINT}, W), \mu})_{\text{BB}'} |\hat{\psi}\rangle \right| \tag{295} \\ &\quad + \left| \mathbb{E}_u \sum_{g, a: a \neq g(u)} \langle \hat{\psi} | ((\hat{M}_a^{(\text{POINT}, W), \mu})_{\text{AB}''} - (\hat{M}_a^{(\text{POINT}, W), \mu})_{\text{BB}'} \cdot (\hat{S}_g^W)_{\text{BB}'} \cdot (\hat{M}_a^{(\text{POINT}, W), \mu})_{\text{AB}''} |\hat{\psi}\rangle \right| \tag{296} \end{aligned}$$

Using Cauchy-Schwarz, we can bound the first term on the right hand side of the inequality by

$$\begin{aligned}
& \left( \mathbb{E}_u \sum_{g,a:a \neq g(u)} \langle \hat{\psi} | (\hat{M}_a^{(\text{POINT},W),u} \cdot \hat{S}_g^W \cdot \hat{M}_a^{(\text{POINT},W),u})_{\text{BB}'} | \hat{\psi} \rangle \right)^{1/2} \\
& \left( \mathbb{E}_u \sum_{g,a:a \neq g(u)} \langle \hat{\psi} | ((\hat{M}_a^{(\text{POINT},W),u})_{\text{AB}''} - (\hat{M}_a^{(\text{POINT},W),u})_{\text{BB}'}) \cdot (\hat{S}_g^W)_{\text{BB}'} \cdot \right. \\
& \quad \left. ((\hat{M}_a^{(\text{POINT},W),u})_{\text{AB}''} - (\hat{M}_a^{(\text{POINT},W),u})_{\text{BB}'}) | \hat{\psi} \rangle \right)^{1/2} \\
& \leq \sqrt{1} \cdot \sqrt{\varepsilon}
\end{aligned}$$

The inequality follows from the self-consistency of  $\hat{M}_a^{(\text{POINT},W),u}$ , as established in Lemma A.9. Similarly, we can bound the second term in Line (296) by  $\sqrt{\varepsilon}$ . This establishes the approximation in Line (293).

The inequality in Line (294) follows from Equation (251) of Lemma A.21. This concludes the derivation of the approximation in Line (289).

**The approximation in Line (290).** We establish this approximation by bounding the magnitude of the difference:

$$\begin{aligned}
& \mathbb{E}_u \sum_a \left\| \sum_{\substack{g,h,g',h': \\ (g'-g_{h'}) (u) = (g-g_h) (u) \\ (\text{Dec}(g)-h) \cdot \hat{u} = a}} \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{S}_{g'}^W \cdot (I - \hat{M}_{g'(u)}^{(\text{POINT},W),u}))_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) | \hat{\psi} \rangle \right\|^2 \\
& = \mathbb{E}_u \sum_{\substack{g,h,g',h': \\ (g'-g_{h'}) (u) \\ = (g-g_h) (u)}} \langle \hat{\psi} | (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \otimes ((I - \hat{M}_{g'(u)}^{(\text{POINT},W),u}) \cdot \hat{S}_{g'}^W \cdot \\
& \quad (I - \hat{M}_{g'(u)}^{(\text{POINT},W),u}))_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} | \hat{\psi} \rangle \\
& \leq \mathbb{E}_u \sum_{g'} \langle \hat{\psi} | ((I - \hat{M}_{g'(u)}^{(\text{POINT},W),u}) \cdot \hat{S}_{g'}^W \cdot (I - \hat{M}_{g'(u)}^{(\text{POINT},W),u}))_{\text{BB}'} | \hat{\psi} \rangle \\
& \leq \delta_S .
\end{aligned}$$

The last inequality follows from the second item of Lemma A.23 (to be proved below).

**The approximation in Line (291).** We establish this approximation by bounding the magnitude of the difference:

$$\begin{aligned}
& \mathbb{E}_u \sum_a \left\| \sum_{\substack{g,h,g',h': \\ g'-g_{h'} \neq g-g_h \\ (g'-g_{h'}) (u) = (g-g_h) (u) \\ (\text{Dec}(g)-h) \cdot \hat{u} = a}} \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{S}_{g'}^W)_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) | \hat{\psi} \rangle \right\|^2 \\
& = \sum_{\substack{g,h,g',h': \\ g'-g_{h'} \neq g-g_h}} \langle \hat{\psi} | \left( (\hat{S}_g^W)_{\text{AA}'} \otimes (\tau_h^W)_{\text{A}''} \right) \cdot \left( (\hat{S}_{g'}^W)_{\text{BB}'} \otimes (\tau_{h'}^W)_{\text{B}''} \right) | \hat{\psi} \rangle \cdot \mathbb{E}_u \mathbf{1}[(g' - g_{h'}) (u) = (g - g_h) (u)] \\
& \leq md/q.
\end{aligned}$$

The last inequality follows from the Schwartz-Zippel lemma: two distinct polynomials of total degree at most  $md$  cannot agree on more than  $md/q$  fraction of points.  $\square$

We now give a proof of the following Lemma, which was used in several derivations in Lemma A.22.

**Lemma A.23.** For  $W \in \{X, Z\}$ , on average over  $u \in \mathbb{F}_q^m$ , we have that

$$\sum_g (\hat{S}_g^W)_{AA'} \otimes (\hat{M}_{g(u)}^{(\text{POINT}, W), u})_{BA''} \approx_{\delta_S} I \quad (297)$$

and

$$(\hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT}, W), u}))_{AA'} \approx_{\delta_S} 0 \quad (298)$$

where the answer summation is over  $g \in \text{ideg}_{d,m}(\mathbb{F}_q)$ . Furthermore, all symmetric equivalents of these approximations also hold.

*Proof.* We first prove Equation (297). From Equation (250) of Lemma A.21 we have that on average over  $u \in \mathbb{F}_q^m$ ,

$$\langle \hat{\psi} | \sum_g (\hat{S}_g^W)_{AA'} \otimes (\hat{M}_{g(u)}^{(\text{POINT}, W), u})_{BA''} | \hat{\psi} \rangle \geq 1 - \delta_S \quad (299)$$

which is equivalent to  $\sum_g (\hat{S}_g^W)_{AA'} \otimes (\hat{M}_{g(u)}^{(\text{POINT}, W), u})_{BA''} \simeq_{\delta_S} I$ . By Fact 5.18, we have that

$$\sum_g (\hat{S}_g^W)_{AA'} \otimes (\hat{M}_{g(u)}^{(\text{POINT}, W), u})_{BA''} \approx_{\delta_S} I.$$

We now prove Equation (298). Equation (299) and Fact 5.18 shows that

$$(\hat{S}_{[g(u)=a]}^W)_{AA'} \approx_{\delta_S} (\hat{M}_a^{(\text{POINT}, W), u})_{BA''}$$

where the answer summation is over  $a \in \mathbb{F}_q$ . Left-multiplying this expression by  $(\hat{S}_{[g(u)=a]}^W)_{AA'}$  and using Fact 5.19 we get

$$\begin{aligned} (\hat{S}_{[g(u)=a]}^W)_{AA'} &\approx_{\delta_S} (\hat{S}_{[g(u)=a]}^W)_{AA'} \otimes (\hat{M}_a^{(\text{POINT}, W), u})_{BA''} \\ &\approx_{\varepsilon} (\hat{S}_{[g(u)=a]}^W)_{AA'} \cdot \hat{M}_a^{(\text{POINT}, W), u} \end{aligned}$$

where the second approximation follows from the following calculation:

$$\mathbb{E}_u \sum_a \left\| (\hat{S}_{[g(u)=a]}^W)_{AA'} \cdot \left( (\hat{M}_a^{(\text{POINT}, W), u})_{AA'} - (\hat{M}_a^{(\text{POINT}, W), u})_{BA''} \right) | \hat{\psi} \right\|^2 \quad (300)$$

$$\leq \mathbb{E}_u \sum_a \langle \hat{\psi} | \left( (\hat{M}_a^{(\text{POINT}, W), u})_{AA'} - (\hat{M}_a^{(\text{POINT}, W), u})_{BA''} \right)^2 | \hat{\psi} \rangle \quad (301)$$

$$= \mathbb{E}_u \sum_a \left\| (\hat{M}_a^{(\text{POINT}, W), u})_{AA'} - (\hat{M}_a^{(\text{POINT}, W), u})_{BA''} | \hat{\psi} \right\|^2 \quad (302)$$

$$\leq \varepsilon \quad (303)$$

The first inequality follows from the fact that  $\hat{S}_{[g(u)=a]}^W \leq I$  for all  $a$ , and the second inequality follows from the self-consistency of  $\hat{M}_a^{(\text{POINT},W),u}$  (established in Lemma A.9). Thus

$$(\hat{S}_{[g(u)=a]}^W)_{AA'} \approx_{O(\delta_S + \varepsilon)} (\hat{S}_{[g(u)=a]}^W \cdot \hat{M}_a^{(\text{POINT},W),u})_{AA'}.$$

On the other hand, notice that

$$\begin{aligned} & \mathbb{E}_u \sum_g \left\| (\hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}))_{AA'} |\hat{\psi}\rangle \right\|^2 \\ &= \mathbb{E}_u \sum_g \langle \hat{\psi} | ((I - \hat{M}_{g(u)}^{(\text{POINT},W),u}) \cdot \hat{S}_g^W \cdot (I - \hat{M}_{g(u)}^{(\text{POINT},W),u}))_{AA'} |\hat{\psi}\rangle \\ &= \mathbb{E}_u \sum_a \langle \hat{\psi} | ((I - \hat{M}_a^{(\text{POINT},W),u}) \cdot \hat{S}_{[g(u)=a]}^W \cdot (I - \hat{M}_a^{(\text{POINT},W),u}))_{AA'} |\hat{\psi}\rangle \\ &\leq O(\delta_S + \varepsilon). \end{aligned}$$

Equation (298) is thus obtained by absorbing  $\varepsilon$  into the function  $\delta_S$ .  $\square$

The next lemma shows that we can construct local unitaries  $V_A, V_B$  from the measurements  $\hat{S}$  such that the shared state  $|\hat{\psi}\rangle$  is close to  $|AUX\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}$  for some state  $|AUX\rangle$ , and the observables  $\tilde{W}^j(\tilde{u})$  are mapped to Pauli observables  $\tau^W(e; \tilde{u})$  acting on  $|\text{EPR}_q\rangle^{\otimes M}$ . Using the consistency between the  $\{\tilde{M}_a^{W, \tilde{u}}\}_{a \in \mathbb{F}_q}$  measurements and the points measurements  $\{M_a^{(\text{POINT},W),u}\}_{a \in \mathbb{F}_q}$  established in Lemma A.22, and the consistency between the points measurements and the ‘‘total Pauli measurements’’  $\{M_h^{(\text{PAULI},W)}\}_{h \in \mathbb{F}_q^M}$  established in Lemma A.7, we deduce that the total Pauli measurements must be close (up to the local unitaries  $V_A, V_B$ ) to the Pauli projectors  $\{\tau_h^W\}_{h \in \mathbb{F}_q^M}$  acting on  $|\text{EPR}_q\rangle^{\otimes M}$ .

**Lemma A.24.** *There exists a function  $\delta_{\text{QLD}}(\varepsilon, m, d, q) = O(\delta_S^{1/4} + md/q)$  (where  $\delta_S$  is defined in Lemma A.21) and unitaries  $V_A$  acting on registers  $AA'A''$  and  $V_B$  acting on  $BB'B''$  such that*

1. *There exists a state  $|AUX\rangle$  on registers  $AA'BB'$  such that*

$$\left\| V_A \otimes V_B |\hat{\psi}\rangle_{AA'A''BB'B''} - |AUX\rangle_{AA'BB'} \otimes |\text{EPR}_q\rangle_{A''B''}^{\otimes M} \right\|^2 \leq \delta_{\text{QLD}}.$$

2. *For all  $W \in \{X, Z\}, h \in \mathbb{F}_q^M$*

$$\begin{aligned} V_A \left( M_h^{(\text{PAULI},W)} \right)_A V_A^\dagger &\approx_{\delta_{\text{QLD}}} I_{AA'} \otimes (\tau_h^W)_{A''} \\ V_B \left( M_h^{(\text{PAULI},W)} \right)_B V_B^\dagger &\approx_{\delta_{\text{QLD}}} I_{BB'} \otimes (\tau_h^W)_{B''} \end{aligned}$$

where the  $\approx$  statements hold with respect to the state  $|AUX\rangle_{AA'BB'} \otimes |\text{EPR}_q\rangle_{A''B''}^{\otimes M}$ .

*Proof.* Define the linear map

$$V_A = \sum_{g_X, g_Z} (\hat{S}_{g_X, g_Z})_{AA'} \otimes (\tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)))_{A''},$$

where the  $\tau^X(\cdot), \tau^Z(\cdot)$  act on  $(\mathbb{C}^q)^{\otimes M}$ . (Also note that the argument of  $\tau^X$  is  $\text{Dec}(g_Z)$  and the argument of  $\tau^Z$  is  $\text{Dec}(g_X)$ .) We verify the unitarity of  $V_A$ :

$$\begin{aligned} V_A V_A^\dagger &= \sum_{g_X, g_Z} (\hat{S}_{g_X, g_Z})_{AA'} \otimes (\tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)) \tau^Z(\text{Dec}(g_X)) \tau^X(\text{Dec}(g_Z)))_{A''} \\ &= \sum_{g_X, g_Z} (\hat{S}_{g_X, g_Z})_{AA'} \otimes I_{A''} \\ &= I_{AA'A''} \end{aligned}$$

where in the first line we used that the measurement  $\{\hat{S}_{g_X, g_Z}\}$  is projective and in the second line we used that the  $\tau^Z$  and  $\tau^X$  observables are self-inverse. The unitary  $V_B$  is defined analogously.

Let  $\{e_j\}_{j \in \{1, \dots, t\}}$  denote the self-dual basis for  $\mathbb{F}_q$  over  $\mathbb{F}_2$  specified in Section 3.3.1, and let  $\tilde{W}^j$  be the observables introduced in (274). Conjugating the observable  $\tilde{W}^j$  acting on the registers  $AA'A''$  by the unitary  $V_A$  we get

$$\begin{aligned} &V_A \tilde{W}^j(\tilde{u}) V_A^\dagger \\ &= \sum_{g_X, g_Z} (-1)^{\text{tr}(e_j(\text{Dec}(g_W) \cdot \tilde{u}))} \left( \hat{S}_{g_X, g_Z} \right)_{AA'} \otimes \left( \tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)) \tau^W(e_j \tilde{u}) \right. \\ &\quad \left. \tau^Z(\text{Dec}(g_X))^\dagger \tau^X(\text{Dec}(g_Z))^\dagger \right)_{A''} \\ &= \sum_{g_X, g_Z} \left( \hat{S}_{g_X, g_Z} \right)_{AA'} \otimes \left( \tau^W(e_j \tilde{u}) \right)_{A''} \\ &= I_{AA'} \otimes \left( \tau^W(e_j \tilde{u}) \right)_{A''}, \end{aligned}$$

where in the third line we used the (anti-)commutation relation

$$\tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)) \tau^W(e_j \tilde{u}) = (-1)^{\text{tr}(e_j(\text{Dec}(g_W) \cdot \tilde{u}))} \tau^W(e_j \tilde{u}) \tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)).$$

An entirely analogous calculation shows that  $V_B \tilde{W}^j(\tilde{u}) V_B^\dagger = I_{BB'} \otimes \left( \tau^W(e_j \tilde{u}) \right)_{B''}$ .

Next, from the self-consistency property of the observables  $\tilde{W}^j$  specified in Lemma A.22 we get that for  $W \in \{X, Z\}$ ,

$$\mathbb{E}_{j \in \{1, \dots, t\}} \mathbb{E}_{\tilde{u} \in \mathbb{F}_q^M} \langle \hat{\psi} | \left( \tilde{W}^j(\tilde{u}) \otimes I - I \otimes \tilde{W}^j(\tilde{u}) \right)^\dagger \left( \tilde{W}^j(\tilde{u}) \otimes I - I \otimes \tilde{W}^j(\tilde{u}) \right) | \hat{\psi} \rangle \leq \delta_S,$$

which, since  $\tilde{W}^j$  is Hermitian, is equivalent to

$$\mathbb{E}_{j \in \{1, \dots, t\}} \mathbb{E}_{\tilde{u} \in \mathbb{F}_q^M} \langle \hat{\psi} | \tilde{W}^j(\tilde{u}) \otimes \tilde{W}^j(\tilde{u}) | \hat{\psi} \rangle \geq 1 - \delta_S/2. \quad (304)$$

Let  $|\vartheta\rangle = V_A \otimes V_B |\hat{\psi}\rangle$ , and for  $W \in \{X, Z\}$  define the operator

$$\begin{aligned} H_W &= \mathbb{E}_{j \in \{1, \dots, t\}} \mathbb{E}_{\tilde{u} \in \mathbb{F}_q^M} \tau^W(e_j \tilde{u}) \otimes \tau^W(e_j \tilde{u}) \\ &= \mathbb{E}_{j \in \{1, \dots, t\}} \left( \mathbb{E}_{s \in \mathbb{F}_q} \tau^W(e_j s) \otimes \tau^W(e_j s) \right)^{\otimes M} \\ &= \left( \mathbb{E}_{s \in \mathbb{F}_q} \tau^W(s) \otimes \tau^W(s) \right)^{\otimes M} \end{aligned}$$

where in the second line we used that for all  $\tilde{u} \in \mathbb{F}_q^M$ ,  $\tau^W(\tilde{u}) = \bigotimes_{i=1}^M \tau^W(\tilde{u}_i)$  with  $\tau^W(\tilde{u}_i)$  being the generalized Pauli  $W$  observable acting on the  $i$ -th qudit. In the third line, we used the fact that for every fixed  $j$ , we have  $\mathbb{E}_{s \in \mathbb{F}_q} \tau^W(e_j s) \otimes \tau^W(e_j s) = \mathbb{E}_{s \in \mathbb{F}_q} \tau^W(s) \otimes \tau^W(s)$ . Equation (304) is then equivalent to

$$\langle \vartheta | H_W | \vartheta \rangle \geq 1 - \delta_S / 2 ,$$

for  $W \in \{X, Z\}$ .

Observe that

$$\begin{aligned} \mathbb{E}_{s, s'} \tau^X(s) \tau^Z(s') \otimes \tau^X(s) \tau^Z(s') &= \mathbb{E}_{s, s'} \sum_{a, b \in \mathbb{F}_q} (-1)^{\text{tr}(as')} |a + s\rangle \langle a| \otimes (-1)^{\text{tr}(bs')} |b + s'\rangle \langle b| \\ &= \mathbb{E}_s \sum_{a \in \mathbb{F}_q} |a + s\rangle \langle a| \otimes |a + s\rangle \langle a| \\ &= |\text{EPR}_q\rangle \langle \text{EPR}_q| \end{aligned}$$

where to go from the first to the second line we used Lemma 3.25. This implies that  $H_X H_Z = \left( \mathbb{E}_{s, s' \in \mathbb{F}_q} \tau^X(s) \tau^Z(s') \otimes \tau^X(s) \tau^Z(s') \right)^{\otimes M} = |\text{EPR}_q\rangle \langle \text{EPR}_q|^{\otimes M}$ . By the triangle inequality, we have

$$\|H_X |\vartheta\rangle - H_Z |\vartheta\rangle\| \leq \| |\vartheta\rangle - H_X |\vartheta\rangle \| + \| |\vartheta\rangle - H_Z |\vartheta\rangle \| .$$

Note that for  $W \in \{X, Z\}$ ,

$$\| |\vartheta\rangle - H_W |\vartheta\rangle \|^2 = 1 - 2\langle \vartheta | H_W | \vartheta \rangle + \langle \vartheta | H_W^2 | \vartheta \rangle \leq 2 - 2\langle \vartheta | H_W | \vartheta \rangle \leq \delta_S ,$$

where in the inequality we used that  $H_W \leq I$ . Furthermore,

$$\begin{aligned} \langle \vartheta | H_W^2 | \vartheta \rangle &= \|H_W |\vartheta\rangle\|^2 \\ &= \| |\vartheta\rangle + (H_W - I) |\vartheta\rangle \|^2 \\ &\geq (\| |\vartheta\rangle \| - \|(H_W - I) |\vartheta\rangle\|)^2 \\ &\geq (1 - \delta_S)^2 \\ &\geq 1 - 2\delta_S . \end{aligned}$$

Therefore

$$\begin{aligned} 2\sqrt{\delta_S} &\geq \|H_X |\vartheta\rangle - H_Z |\vartheta\rangle\|^2 \\ &= \langle \vartheta | H_X^2 | \vartheta \rangle + \langle \vartheta | H_Z^2 | \vartheta \rangle - 2\langle \vartheta | H_X H_Z | \vartheta \rangle \\ &\geq 2 - 4\delta_S - 2\langle \vartheta | H_X H_Z | \vartheta \rangle , \end{aligned}$$

which implies that

$$\langle \vartheta | (I_{AA'BB'} \otimes (|\text{EPR}_q\rangle \langle \text{EPR}_q|^{\otimes M})_{A''B''}) | \vartheta \rangle \geq 1 - 2\delta_S - \sqrt{\delta_S} .$$

Define the unnormalized state  $|\text{AUX}_0\rangle$  on the registers  $AA'BB'$  as

$$|\text{AUX}_0\rangle_{AA'BB'} = (\langle \text{EPR}_q |^{\otimes M})_{A''B''} \cdot |\vartheta\rangle_{AA'A''BB'B''}$$

and define the normalized state  $|\text{AUX}\rangle = \frac{1}{\| |\text{AUX}_0\rangle \|} |\text{AUX}_0\rangle$ . The inner product between  $|\vartheta\rangle$  and  $|\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}$  can thus be evaluated as

$$\begin{aligned}
\langle \vartheta | (|\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}) &= \frac{1}{\| |\text{AUX}_0\rangle \|} \cdot \langle \vartheta | (|\text{AUX}_0\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}) \\
&= \frac{1}{\| |\text{AUX}_0\rangle \|} \cdot \langle \vartheta | (I_{\text{AA}'\text{BB}'} \otimes (|\text{EPR}_q\rangle\langle \text{EPR}_q|_{\text{A}''\text{B}''})^{\otimes M}) | \vartheta \rangle \\
&= \frac{1}{\| |\text{AUX}_0\rangle \|} \cdot \| |\text{AUX}_0\rangle \|^2 \\
&= \| |\text{AUX}_0\rangle \| \\
&\geq \sqrt{1 - 2\delta_S - \sqrt{\delta_S}} \\
&\geq 1 - O(\sqrt{\delta_S}).
\end{aligned}$$

This implies  $\| |\vartheta\rangle - |\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M} \|^2 \leq O(\sqrt{\delta_S})$ , which establishes the first item of the lemma.

We now establish the second item of the lemma. In what follows, all  $\simeq$  and  $\approx$  statements hold with respect to the state  $|\hat{\psi}\rangle$ . From Lemma A.7 we have the following implications: for all  $W \in \{X, Z\}$  and on average over  $u \in \mathbb{F}_q^m$ ,

$$M_{[g_h(u)=a]}^{(\text{PAULI}, W)} \otimes I_{\text{BB}'\text{B}''} \simeq_\varepsilon I_{\text{AA}'\text{A}''} \otimes M_a^{(\text{POINT}, W), u}, \quad (305)$$

$$I_{\text{AA}'\text{A}''} \otimes M_a^{(\text{POINT}, W), u} \simeq_\varepsilon M_a^{(\text{POINT}, W), u} \otimes I_{\text{BB}'\text{B}''}, \quad (306)$$

where  $M_{[g_h(u)=a]}^{(\text{PAULI}, W)} = \sum_{h: g_h(u)=a} M_h^{(\text{PAULI}, W)}$ . From Lemma A.22 we get that

$$M_a^{(\text{POINT}, W), u} \otimes I_{\text{BB}'\text{B}''} \simeq_{\delta_S} I_{\text{AA}'\text{A}''} \otimes \tilde{M}_a^{W, \text{ind}_m(u)}. \quad (307)$$

Using Fact 5.23 with Equations (305), (306), and (307), we get

$$M_{[g_h(u)=a]}^{(\text{PAULI}, W)} \otimes I_{\text{BB}'\text{B}''} \simeq_{\sqrt{\delta_S}} I_{\text{AA}'\text{A}''} \otimes \tilde{M}_a^{W, \text{ind}_m(u)}, \quad (308)$$

where we used that  $\delta_S \geq \varepsilon$ .

Next, we have for all  $u \in \mathbb{F}_q^m$

$$\begin{aligned}
&V_{\text{B}} \tilde{M}_a^{W, \text{ind}_m(u)} V_{\text{B}}^\dagger \\
&= \sum_{g_X, g_Z} \left( \hat{S}_{g_X, g_Z} \right)_{\text{BB}'} \otimes \left( \tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)) \tau_{g_W(u)-a}^W(\text{ind}_m(u)) \tau^Z(\text{Dec}(g_X))^\dagger \tau^X(\text{Dec}(g_Z))^\dagger \right)_{\text{B}''} \\
&= \sum_{g_X, g_Z} \left( \hat{S}_{g_X, g_Z} \right)_{\text{BB}'} \otimes \sum_{h: g_h(u)=g_W(u)-a} \left( \tau^X(\text{Dec}(g_Z)) \tau^Z(\text{Dec}(g_X)) \tau_h^W \tau^Z(\text{Dec}(g_X))^\dagger \tau^X(\text{Dec}(g_Z))^\dagger \right)_{\text{B}''} \\
&= \sum_{g_X, g_Z} \left( \hat{S}_{g_X, g_Z} \right)_{\text{BB}'} \otimes \sum_{h: g_h(u)=g_W(u)-a} \left( \tau_{h+\text{Dec}(g_W)}^W \right)_{\text{B}''} \\
&= \sum_{g_X, g_Z} \left( \hat{S}_{g_X, g_Z} \right)_{\text{BB}'} \otimes \sum_{h: g_h(u)=a} \left( \tau_h^W \right)_{\text{B}''} \\
&= I_{\text{BB}'} \otimes \left( \tau_{[g_h(u)=a]}^W \right)_{\text{B}''}, \quad (309)
\end{aligned}$$



where in the second line we expanded out the definitions of  $V_B$  and  $\tilde{M}_a^{W, \text{ind}_m(u)}$  (defined in Eq. (273)); in the third line we expanded out the definition of  $\tau_{g_W(u)-a}^W(\text{ind}_m(u))$  (defined in Eq. (272)) and used the fact that for all  $h \in \mathbb{F}_q^M$ , the inner product  $h \cdot \text{ind}_m(u)$  is equal to  $g_h(u)$ ; in the fourth line we used the identities

$$\tau^X(s) \tau_h^Z \tau^X(s)^\dagger = \tau_{h+s}^Z \quad \text{and} \quad \tau^Z(s) \tau_h^X \tau^Z(s)^\dagger = \tau_{h+s}^X;$$

and in the fifth line is due to the following short calculation: for all fixed  $g_W$  we have

$$\sum_{h: g_h(u)=g_W(u)-a} \tau_{h+\text{Dec}(g_W)}^W = \sum_{h: g_W(u)-g_h(u)=a} \tau_{h+\text{Dec}(g_W)}^W = \sum_{h: g_W(u)+g_h(u)=a} \tau_{h+\text{Dec}(g_W)}^W = \sum_{h': g_{h'}(u)=a} \tau_{h'}^W.$$

The second equality from the fact that  $\mathbb{F}_q$  is a field of characteristic 2 (so subtraction is the same as addition), and the third equality follows from the fact that letting  $h' = h + \text{Dec}(g_W)$ , we have  $g_{h'} = g_h + g_W$  (in other words, the low-degree encoding  $h \mapsto g_h$  is linear in  $h$ ).

Let  $|\Delta\rangle = |\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}$ . We show that  $V_A \left( M_h^{(\text{PAULL}, W)} \right)_A V_A^\dagger \approx_{\delta_{\text{QLD}}} I_{AA'} \otimes (\tau_h^W)_{A''}$  (the argument for the operators acting on registers  $\text{BB}'\text{B}''$  proceeds identically).

$$\begin{aligned} & \sum_h \langle \Delta | \left( V_A M_h^{(\text{PAULL}, W)} V_A^\dagger - (\tau_h^W)_{A''} \right)^\dagger \left( V_A M_h^{(\text{PAULL}, W)} V_A^\dagger - (\tau_h^W)_{A''} \right) | \Delta \rangle \\ &= 2 - 2\Re \sum_h \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger (\tau_h^W)_{A''} | \Delta \rangle \\ &= 2 - 2 \sum_h \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_h^W)_{B''} | \Delta \rangle \end{aligned} \quad (310)$$

where in the last line we used that  $(\tau_h^W)_{A''} |\text{EPR}_q\rangle^{\otimes M} = (\tau_h^W)_{B''} |\text{EPR}_q\rangle^{\otimes M}$ . We now claim that

$$\begin{aligned} & \sum_h \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_h^W)_{B''} | \Delta \rangle \\ & \approx_{md/q} \mathbb{E}_u \sum_{\substack{h, h': \\ g_h(u)=g_{h'}(u)}} \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_{h'}^W)_{B''} | \Delta \rangle \end{aligned} \quad (311)$$

$$\begin{aligned} &= \mathbb{E}_u \sum_{a \in \mathbb{F}_q} \langle \Delta | V_A M_{[g_h(u)=a]}^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_{[g_h(u)=a]}^W)_{B''} | \Delta \rangle \\ &= \mathbb{E}_u \sum_{a \in \mathbb{F}_q} \langle \Delta | V_A M_{[g_h(u)=a]}^{(\text{PAULL}, W)} V_A^\dagger \otimes V_B \tilde{M}_a^{W, \text{ind}_m(u)} V_B^\dagger | \Delta \rangle \end{aligned} \quad (312)$$

where the equality in Equation (312) follows from the identity in Equation (309), and the approximation in Equation (311) follows from the following calculation:

$$\begin{aligned} & \mathbb{E}_u \sum_{\substack{h \neq h': \\ g_h(u)=g_{h'}(u)}} \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_{h'}^W)_{B''} | \Delta \rangle \\ &= \sum_{h \neq h'} \left( \mathbb{E}_u \mathbf{1}[g_h(u) = g_{h'}(u)] \right) \langle \Delta | V_A M_h^{(\text{PAULL}, W)} V_A^\dagger \otimes (\tau_{h'}^W)_{B''} | \Delta \rangle \\ &\leq \frac{md}{q}. \end{aligned}$$

In the last line we used that for distinct  $h \neq h'$ , the polynomials  $g_h \neq g_{h'}$  and thus by the Schwartz-Zippel lemma can only agree on at most  $md/q$  fraction of points  $u \in \mathbb{F}_q^m$ .

Continuing from Equation (312), we use the first item of the Lemma which shows that  $\|\Delta\rangle - |\vartheta\rangle\|^2 \leq O(\sqrt{\delta_S})$  and equivalently  $\|V_A^\dagger \otimes V_B^\dagger |\Delta\rangle - |\hat{\psi}\rangle\|^2 \leq O(\sqrt{\delta_S})$ , so

$$\begin{aligned} \text{Eq. (312)} &\approx_{O(\delta_S^{1/4})} \mathbb{E}_u \sum_{a \in \mathbb{F}_q} \langle \hat{\psi} | M_{[g_h(u)=a]}^{(\text{PAULI}, W)} \otimes \tilde{M}_a^{W, \text{ind}_m(u)} | \hat{\psi} \rangle \\ &\geq 1 - O(\delta_S^{1/2}), \end{aligned}$$

where the last line comes from (308). Together with Equation (310),

$$\text{Eq. (310)} \leq O(\delta_S^{1/4} + md/q).$$

This concludes the proof of the second item of the lemma.  $\square$

We now prove Theorem A.1.

*Proof of Theorem A.1.* Let  $\mathcal{S} = (\psi, M)$  be a projective strategy for  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  that succeeds with probability at least  $1 - \varepsilon$ , where  $|\psi\rangle$  is a bipartite state on registers AB. As mentioned at the beginning of Appendix A.2, we assume that the state  $|\psi\rangle$  is padded with sufficiently many ancilla  $|0\rangle$  qubits. Let  $\phi_A$  be the following isometry mapping the register A to the registers AA'A'' where A' and A'' are isomorphic to  $(\mathbb{C}^q)^{\otimes M}$ :

$$\phi_A : |\theta\rangle_A \mapsto V_A (|\theta\rangle_A \otimes (|\text{EPR}_q\rangle^{\otimes M})_{A'A''})$$

where  $V_A$  is the unitary acting on AA'A'' from Lemma A.24. Define the isometry  $\phi_B$  analogously. Lemma A.24 then implies that there exists a state  $|\text{AUX}\rangle$  on registers AA'BB' such that

$$\left\| \phi_A \otimes \phi_B |\psi\rangle - |\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M} \right\|^2 \leq \delta_{\text{QLD}}$$

and for  $W \in \{X, Z\}$ ,

$$\phi_A (M_h^{(\text{PAULI}, W)})_A \phi_A^\dagger \approx_{\delta_{\text{QLD}}} (\tau_h^W)_{A''} \quad \text{and} \quad \phi_B (M_h^{(\text{PAULI}, W)})_B \phi_B^\dagger \approx_{\delta_{\text{QLD}}} (\tau_h^W)_{B''}$$

where the  $\approx$  statement holds with respect to the state  $|\text{AUX}\rangle \otimes |\text{EPR}_q\rangle^{\otimes M}$ . This shows that  $\mathfrak{G}_{\text{qldparams}}^{\text{PAULI}}$  is a self-test for the strategy  $\mathcal{S}^{\text{PAULI}}$  with robustness  $\delta_{\text{QLD}}(\varepsilon, m, d, q)$ . We conclude the proof by making the identification of registers AA' with A' and BB' with B' to be consistent with the statement of Theorem A.1. We also note that, when unpacking the dependence on parameters  $\varepsilon, m, d, q$ , the function  $\delta_{\text{QLD}}(\varepsilon, m, d, q) = O(\delta_S^{1/4} + md/q)$  has the form  $a(md)^a(\varepsilon^b + q^{-b} + 2^{-bmd})$  for some universal constants  $a > 1, 0 < b < 1$ , as desired.  $\square$

## References

- [Ara02] PK Aravind. A simple demonstration of Bell’s theorem involving two observers and no probabilities or inequalities. *arXiv preprint quant-ph/0206070*, 2002. [7.2](#), [7.2](#)
- [AS98] Sanjeev Arora and Shmuel Safra. Probabilistic checking of proofs: A new characterization of NP. *Journal of the ACM (JACM)*, 45(1):70–122, 1998. [16](#)
- [BFL91] László Babai, Lance Fortnow, and Carsten Lund. Non-deterministic exponential time has two-prover interactive protocols. *Computational complexity*, 1(1):3–40, 1991. [1.1](#), [1.1](#), [1.2](#), [16](#), [7.1.1](#)
- [BSGH<sup>+</sup>06] Eli Ben-Sasson, Oded Goldreich, Prahladh Harsha, Madhu Sudan, and Salil Vadhan. Robust PCPs of proximity, shorter PCPs, and applications to coding. *SIAM Journal on Computing*, 36(4):889–974, 2006. [2](#)
- [BSS05] Eli Ben-Sasson and Madhu Sudan. Simple PCPs with poly-log rate and query complexity. In *Proceedings of the thirty-seventh annual ACM symposium on Theory of computing*, pages 266–275. ACM, 2005. [2](#)
- [BSS08] Eli Ben-Sasson and Madhu Sudan. Short PCPs with polylog query complexity. *SIAM Journal on Computing*, 38(2):551–607, 2008. [10.4.1](#)
- [BVY17] Mohammad Bavarian, Thomas Vidick, and Henry Yuen. Hardness amplification for entangled games via anchoring. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pages 303–316. ACM, 2017. [2](#), [11](#), [11.1](#)
- [CHSH69] John F Clauser, Michael A Horne, Abner Shimony, and Richard A Holt. Proposed experiment to test local hidden-variable theories. *Physical review letters*, 23(15):880, 1969. [1.2](#)
- [CHTW04] Richard Cleve, Peter Hoyer, Benjamin Toner, and John Watrous. Consequences and limits of nonlocal strategies. In *Proceedings. 19th IEEE Annual Conference on Computational Complexity, 2004.*, pages 236–249. IEEE, 2004. [1.2](#)
- [CLP15] Valerio Capraro, Martino Lupini, and Vladimir Pestov. *Introduction to sofic and hyperlinear groups and Connes’ embedding conjecture*, volume 2136. Springer, 2015. [1.4](#)
- [CM14] Richard Cleve and Rajat Mittal. Characterization of binary constraint system games. In *International Colloquium on Automata, Languages, and Programming*, pages 320–331. Springer, 2014. [7.2](#)
- [Col19] Andrea Coladangelo. A two-player dimension witness based on embezzlement, and an elementary proof of the non-closure of the set of quantum correlations. *arXiv preprint arXiv:1904.02350*, 2019. [1.3](#)
- [Con76] Alain Connes. Classification of injective factors cases  $II_1$ ,  $II_\infty$ ,  $III_\lambda$ ,  $\lambda \neq 1$ . *Annals of Mathematics*, pages 73–115, 1976. [1](#), [1.3](#), [1.3](#)
- [CS17] Andrea Coladangelo and Jalex Stark. Robust self-testing for linear constraint system games. *arXiv preprint arXiv:1709.09267*, 2017. [7.2](#)

- [CS18] Andrea Coladangelo and Jalex Stark. Unconditional separation of finite and infinite-dimensional quantum correlations. *arXiv preprint arXiv:1804.05116*, 2018. 1.2, 1.3
- [DLTW08] Andrew C Doherty, Yeong-Cherng Liang, Ben Toner, and Stephanie Wehner. The quantum moment problem and bounds on entangled multi-prover games. In *2008 23rd Annual IEEE Conference on Computational Complexity*, pages 199–210. IEEE, 2008. 1.3, 1.4, 12.3
- [DPP19] Ken Dykema, Vern I Paulsen, and Jitendra Prakash. Non-closure of the set of quantum correlations via graphs. *Communications in Mathematical Physics*, 365(3):1125–1142, 2019. 1.3
- [DSV15] Irit Dinur, David Steurer, and Thomas Vidick. A parallel repetition theorem for entangled projection games. *Computational Complexity*, 24(2):201–254, 2015. 11, 11.2
- [Eke91] Artur K Ekert. Quantum cryptography based on bell’s theorem. *Physical review letters*, 67(6):661, 1991. 1.2
- [FJVY19] Joseph Fitzsimons, Zhengfeng Ji, Thomas Vidick, and Henry Yuen. Quantum proof systems for iterated exponential time, and beyond. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 473–480. ACM, 2019. 1.2, 1.2, 1.3, 1.3
- [FL92] Uriel Feige and László Lovász. Two-prover one-round proof systems: Their power and their problems. In *Proceedings of the twenty-fourth annual ACM symposium on Theory of computing*, pages 733–744. ACM, 1992. 1.1
- [FNT14] Tobias Fritz, Tim Netzer, and Andreas Thom. Can you compute the operator norm? *Proceedings of the American Mathematical Society*, 142(12):4265–4276, 2014. 1.3, 10, 1.4
- [Fri12] Tobias Fritz. Tsirelson’s problem and Kirchberg’s conjecture. *Reviews in Mathematical Physics*, 24(05):1250012, 2012. 1, 1.3, 1.4
- [GH16] Isaac Goldbring and Bradd Hart. Computability and the Connes embedding problem. *Bulletin of Symbolic Logic*, 22(2):238–248, 2016. 1.3
- [GH20] Isaac Goldbring and Bradd Hart. The universal theory of the hyperfinite  $\text{II}_1$  factor is not computable. *arXiv preprint arXiv:2006.05629*, 2020. 1.3
- [Har04] Prahladh Harsha. *Robust PCPs of proximity and shorter PCPs*. PhD thesis, Massachusetts Institute of Technology, 2004. 10.4
- [HS65] Juris Hartmanis and Richard E Stearns. On the computational complexity of algorithms. *Transactions of the American Mathematical Society*, 117:285–306, 1965. 3.1, 3.1
- [IKM09] Tsuyoshi Ito, Hirotada Kobayashi, and Keiji Matsumoto. Oracularization and two-prover one-round interactive proofs against nonlocal strategies. In *2009 24th Annual IEEE Conference on Computational Complexity*, pages 217–228. IEEE, 2009. 1.2
- [IV12] Tsuyoshi Ito and Thomas Vidick. A multi-prover interactive proof for NEXP sound against entangled provers. In *2012 IEEE 53rd Annual Symposium on Foundations of Computer Science*, pages 243–252. IEEE, 2012. 1.2

- [Ji16] Zhengfeng Ji. Classical verification of quantum proofs. In *Proceedings of the forty-eighth annual ACM symposium on Theory of Computing*, pages 885–898. ACM, 2016. [1.2](#)
- [Ji17] Zhengfeng Ji. Compression of quantum multi-prover interactive proofs. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, pages 289–302. ACM, 2017. [1.2](#), [1.2](#), [1.3](#)
- [JNP<sup>+</sup>11] Marius Junge, Miguel Navascues, Carlos Palazuelos, David Perez-Garcia, Volkher B Scholz, and Reinhard F Werner. Connes’ embedding problem and Tsirelson’s problem. *Journal of Mathematical Physics*, 52(1):012102, 2011. [1](#), [1.3](#), [1.4](#)
- [JNV<sup>+</sup>20] Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen. Quantum soundness of the classical low individual degree test. *arXiv preprint arXiv:2009.12982*, 2020. [1.2](#), [3.6](#), [5.23](#), [7](#), [7.1.1](#), [8.4.3](#), [8.4.3](#), [A.1](#), [A.2](#)
- [JNV<sup>+</sup>21] Zhengfeng Ji, Anand Natarajan, Thomas Vidick, John Wright, and Henry Yuen. Quantum soundness of testing tensor codes. *arXiv preprint arXiv:2111.08131*, 2021. [7.1.1](#), [7.1.1](#), [1](#), [2](#), [3](#), [7.1.1](#)
- [JPY14] Rahul Jain, Attila Pereszlényi, and Penghui Yao. A parallel repetition theorem for entangled two-player one-round games under product distributions. In *2014 IEEE 29th Conference on Computational Complexity (CCC)*, pages 209–216. IEEE, 2014. [11](#)
- [KKM<sup>+</sup>11] Julia Kempe, Hirotada Kobayashi, Keiji Matsumoto, Ben Toner, and Thomas Vidick. Entangled games are hard to approximate. *SIAM Journal on Computing*, 40(3):848–877, 2011. [1.2](#)
- [Kle54] Stephen C. Kleene. Introduction to Metamathematics. *Journal of Symbolic Logic*, 19(3):215–216, 1954. [15](#), [35](#)
- [KPS18] Se-Jin Kim, Vern Paulsen, and Christopher Schafhauser. A synchronous game for binary constraint systems. *Journal of Mathematical Physics*, 59(3):032201, 2018. [1.2](#), [1.4](#), [5.13](#)
- [KV11] Julia Kempe and Thomas Vidick. Parallel repetition of entangled games. In *Proceedings of the 43rd Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2011, pages 353–362, 2011. [3.6](#), [A.1](#)
- [LFKN90] Carsten Lund, Lance Fortnow, Howard Karloff, and Noam Nisan. Algebraic methods for interactive proof systems. In *Proceedings of 31st Annual Symposium on Foundations of Computer Science*, pages 2–10. IEEE, 1990. [1.1](#)
- [LJ91] H.W. Lenstra Jr. Finding isomorphisms between finite fields. *Mathematics of Computation*, 56(193):329–347, 1991. [3.3.2](#)
- [Mer90] David Mermin. Simple unified form for the major no-hidden-variables theorems. *Physical Review Letters*, 65(27):3373, 1990. [7.2](#)
- [MNY20] Hamoon Mousavi, Seyed Sajjad Nezhadi, and Henry Yuen. On the complexity of zero gap MIP\*. *arXiv preprint arXiv:2002.10490*, 2020. [1.4](#)

- [MP13] Gary L Mullen and Daniel Panario. *Handbook of finite fields*. Chapman and Hall/CRC, 2013. [3.3](#), [3.3.1](#)
- [MR18] Magdalena Musat and Mikael Rørdam. Non-closure of quantum correlation matrices and factorizable channels that require infinite dimensional ancilla. *arXiv preprint arXiv:1806.10242*, 2018. [1.3](#)
- [NPA08] Miguel Navascués, Stefano Pironio, and Antonio Acín. A convergent hierarchy of semidefinite programs characterizing the set of quantum correlations. *New Journal of Physics*, 10(7):073013, 2008. [1](#), [1](#), [1.3](#), [1.4](#), [12.3](#)
- [NV17] Anand Natarajan and Thomas Vidick. A quantum linearity test for robustly verifying entanglement. In *Proceedings of the 49th Annual ACM SIGACT Symposium on Theory of Computing*, STOC 2017, pages 1003–1015, New York, NY, USA, 2017. ACM. [A.4](#)
- [NV18a] Anand Natarajan and Thomas Vidick. Low-degree testing for quantum states, and a quantum entangled games PCP for QMA. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 731–742. IEEE, 2018. [1.2](#), [1.2](#), [2](#), [7](#), [7.1.1](#), [7.3](#)
- [NV18b] Anand Natarajan and Thomas Vidick. Two-player entangled games are NP-hard. In *Proceedings of the 33rd Computational Complexity Conference*, page 20. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2018. [1.2](#), [2](#), [16](#)
- [NW19] Anand Natarajan and John Wright.  $NEEXP \subseteq MIP^*$ . *arXiv preprint arXiv:1904.05870v3*, 2019. [1.2](#), [1.2](#), [2](#), [2](#), [2](#), [2](#), [3.25](#), [5.2](#), [5.18](#), [5.19](#), [5.22](#), [5.24](#), [5.2](#), [5.2](#), [5.27](#), [5.28](#), [7.1.1](#), [7.3](#), [7.3.1](#), [8.1](#), [8.4.1](#), [8.4.1](#), [10](#), [10.1](#), [10.1](#), [1](#), [10.4](#)
- [Oza13] Narutaka Ozawa. About the Connes embedding conjecture. *Japanese Journal of Mathematics*, 8(1):147–183, 2013. [1.3](#), [1.4](#)
- [Pap94] Christos Papadimitriou. *Computational complexity*. Addison Wesley, 1994. [3.1](#), [10.1](#), [10.2](#)
- [Per90] Asher Peres. Incompatible results of quantum measurements. *Physics Letters A*, 151(3-4):107–108, 1990. [7.2](#)
- [PSS<sup>+</sup>16] Vern I Paulsen, Simone Severini, Daniel Stahlke, Ivan G Todorov, and Andreas Winter. Estimating quantum chromatic numbers. *Journal of Functional Analysis*, 270(6):2188–2222, 2016. [1.2](#), [5.13](#)
- [PV16] Carlos Palazuelos and Thomas Vidick. Survey on nonlocal games and operator space theory. *Journal of Mathematical Physics*, 57(1):015220, 2016. [7](#)
- [Rog87] Hartley Rogers, Jr. *Theory of Recursive Functions and Effective Computability*. MIT Press, Cambridge, MA, USA, 1987. [15](#)
- [RS96] Ronitt Rubinfeld and Madhu Sudan. Robust characterizations of polynomials with applications to program testing. *SIAM Journal on Computing*, 25(2):252–271, 1996. [16](#)
- [RS97] Ran Raz and Shmuel Safra. A sub-constant error-probability low-degree test, and a sub-constant error-probability PCP characterization of NP. In *Proceedings of the twenty-ninth annual ACM symposium on Theory of computing*, pages 475–484, 1997. [16](#)

- [Sch80] Jacob Schwartz. Fast probabilistic algorithms for verification of polynomial identities. *Journal of the ACM*, 27(4):701–717, 1980. 3.20
- [Sha90] Adi Shamir.  $IP = PSPACE$ . In *Proceedings of 31st Annual Symposium on Foundations of Computer Science*, pages 11–15. IEEE, 1990. 1.1
- [Sho90] Victor Shoup. New algorithms for finding irreducible polynomials over finite fields. *Mathematics of Computation*, 54(189):435–447, 1990. 3.3.2
- [Sip12] Michael Sipser. *Introduction to the Theory of Computation*. Cengage Learning, 2012. 3.1
- [Slo19a] William Slofstra. The set of quantum correlations is not closed. In *Forum of Mathematics, Pi*, volume 7. Cambridge University Press, 2019. 1, 1.3, 1.4
- [Slo19b] William Slofstra. Tsirelson’s problem and an embedding theorem for groups arising from non-local games. *Journal of the American Mathematical Society*, 2019. 1.3
- [Tsi93] Boris S Tsirelson. Some results and problems on quantum bell-type inequalities. *Hadronic Journal Supplement*, 8(4):329–345, 1993. 1.3
- [Tsi06] Boris S Tsirelson. Bell inequalities and operator algebras, 2006. Problem statement from website of open problems at TU Braunschweig (2006), available at <http://web.archive.org/web/20090414083019/http://www.imaph.tu-bs.de/qi/pr> 1, 1.3
- [Tur37] Alan M. Turing. On computable numbers, with an application to the Entscheidungsproblem. *Proceedings of the London mathematical society*, 2(1):230–265, 1937. 3.1
- [Vid16] Thomas Vidick. Three-player entangled xor games are np-hard to approximate. *SIAM Journal on Computing*, 45(3):1007–1063, 2016. 1.2, 7.1.1
- [VW16] Thomas Vidick and John Watrous. Quantum proofs. *Foundations and Trends in Theoretical Computer Science*, 11(1-2):1–215, 2016. 5.3, 12.2
- [Wan89] Charles C. Wang. An algorithm to design finite field multipliers using a self-dual normal basis. *IEEE Transactions on Computers*, 38(10):1457–1460, 1989. 3.3.2
- [Yue16] Henry Yuen. A parallel repetition theorem for all entangled games. In *43rd International Colloquium on Automata, Languages, and Programming (ICALP 2016)*. Schloss Dagstuhl-Leibniz-Zentrum fuer Informatik, 2016. 11.2
- [Zip79] Richard Zippel. Probabilistic algorithms for sparse polynomials. In *Symbolic and Algebraic Computation*, pages 216–226, 1979. 3.20