

Noninvertible Gauge Symmetry in (2+1)d Topological Orders: A String-Net Model Realization

Yu Zhao^{a,b} Yidun Wan^{1a,b}

^a*State Key Laboratory of Surface Physics, Center for Astronomy and Astrophysics, Department of Physics, Center for Field Theory and Particle Physics, and Institute for Nanoelectronic devices and Quantum Computing, Fudan University, 2005 Songhu Road, Shanghai 200433, China*

^b*Shanghai Research Center for Quantum Sciences, 99 Xiupu Road, Shanghai 201315, China*

E-mail: yuzhao20@fudan.edu.cn, ydwan@fudan.edu.cn

ABSTRACT: We develop a systematic framework for understanding symmetries in topological phases in $2 + 1$ dimensions using the string-net model, encompassing both gauge symmetries that preserve anyon species and global symmetries permuting anyon species, including both invertible symmetries describable by groups and noninvertible symmetries described by categories. As an archetypal example, we reveal the first noninvertible categorical gauge symmetry of topological orders in $2 + 1$ dimensions: the Fibonacci gauge symmetry of the doubled Fibonacci topological order, described by the Fibonacci fusion 2-category. Our approach involves two steps: first, establishing duality between different string-net models with Morita equivalent input fusion categories that describe the same topological order; and second, constructing symmetry transformations within the same string-net model when the dual models have isomorphic input fusion categories, achieved by composing duality maps with isomorphisms of degrees of freedom between the dual models.

¹Corresponding author

Contents

1	Introduction	2
2	Sketch of Our Approach	3
3	Fibonacci String-Net Model	4
4	Enlarging the Hilbert Space and the Duality Map	6
5	Symmetry Transformation of the Doubled Fibonacci Topological Order	8
6	Fibonacci Categorical Gauge Symmetry	12
A	Review of the Extended String-net Model	15
	A.1 Topological Features	17
	A.2 Excited States	18
B	Frobenius Algebras and Bimodules	19
	B.1 Frobenius Algebra	19
	B.2 Bimodules over a Frobenius Algebra	21
	B.3 The Bimodule Fusion Category over a Frobenius Algebra	21
C	General Constructions of Dualities and Symmetry Transformations in the Extended String-Net Model	23
	C.1 Enlarging the Hilbert Space	25
	C.2 Duality	25
	C.3 Symmetry Transformation	26
	C.4 Braiding of Bimodules	27
D	The \mathbb{Z}_2 Global Symmetry transformation of the \mathbb{Z}_2 Toric Code String-Net Model	27
	D.1 The toric code string-net model	27
	D.2 The global symmetry transformation of the toric code topological order	29
	D.3 Criterion for Distinguishing Gauge Symmetries from Global Symmetries	30
E	Frobenius Algebra of Fibonacci Fusion Categories and Its Simple Bimodules	31

1 Introduction

Symmetry is a central concept in modern physics, traditionally described by groups. Topological orders in $2 + 1$ dimensions, whose low-energy effective descriptions are topological gauge field theories, are novel phases of matter that go beyond the Landau-Ginzburg paradigm, revealing more interesting symmetry structures. Nevertheless, the gauge symmetries¹ in topological orders are not completely clear. In cases where the gauge symmetries are described by groups, e.g., in the Dijkgraaf-Witten theory[1–4] or their lattice Hamiltonian model—the (twisted) quantum double model[5–13], the gauge structure, i.e., gauge group and gauge transformation, has been well understood. In more general cases, such as the Turaev-Viro topological field theory[2, 14–17] or their lattice Hamiltonian model—the string-net (Levin-Wen) model[18–29], the gauge symmetry is not known in general. In such cases, if a gauge symmetry exists, it is likely not describable by any group but rather by categories[30–40], invoking the concept of noninvertible symmetry[36, 37, 40–45], which has greatly expanded and deepened our understanding of symmetry in physics. Noninvertible symmetry has been extensively explored in $1 + 1$ dimensions[46–50]. In $2 + 1$ dimensions, however, noninvertible symmetries remain largely an open problem, despite studies in certain $(2 + 1)$ -dimensional systems[51–53] and attempts in studying the applications of noninvertible symmetry in quantum field theory[54–58].

In this paper, we tackle this problem by explicitly and systematically formulating the symmetry transformations of $(2 + 1)$ -dimensional topological orders as operators of the string-net model. We show that such symmetry transformations can be either gauge symmetries that preserve anyon species or global symmetries permuting anyon species, and can be invertible (describable by groups) or noninvertible (described by a fusion 2-category[59–63]). As a key result, our construction reveals the first noninvertible categorical gauge symmetry of topological orders in $2 + 1$ dimensions:

- The noninvertible gauge symmetry (to be called the Fibonacci gauge symmetry) of the doubled Fibonacci topological order described by the Fibonacci fusion 2-category.

This is an archetypal example to illustrate our general construction, which is rather involved and thus to be detailed in the appendices. In the main text, we shall provide a brief overview of our approach and expound on the Fibonacci gauge symmetry. Besides, we also provide a clear criterion (see Appendix D.3) determining whether a symmetry transformation generates a global symmetry or a gauge symmetry of the topological order. An example of global symmetry so constructed is shown in Appendix D.

Our construction can be applied to realize anyon condensations [31, 64–68] in the string-net model. Previous works have realized fluxon and simple-current condensations [29, 69]—specific classes of anyon condensation—in the string-net model. Nevertheless, general non-Abelian anyon condensations remain unexplored. We can apply symmetry transformations to the string-net model and then perform all types of anyon condensations

¹Gauge symmetry is a historical misnomer. It should be more appropriately called gauge invariance or gauge redundancy because it is mathematical redundancy where different states in the Hilbert space describe the same physical state in the theory.

in familiar forms in Refs. [29, 69] within the transformed model. This is an ongoing work to be reported elsewhere.

2 Sketch of Our Approach

As the low-energy effective theories of topological orders, topological quantum field theory (TQFT) is believed to be a topological gauge theory. For the Dijkgraaf-Witten type of TQFT, the gauge symmetry and transformation are manifest and clear. For more general topological orders, whose low-energy effective theory is the Turaev-Viro type of TQFT, however, their gauge symmetries and transformations are in general obscure.

To reveal the gauge structure of topological orders in general, we first note that topological orders are primarily observed in strongly correlated electron systems[70–77], where anyons are collective excitations of the electrons. Such a system has a physical Hilbert space consisting of the fundamental degrees of freedom of electrons, and anyons are represented as excited states in this Hilbert space. While an Abelian anyon is represented by one excited state up to a phase factor, a non-Abelian anyon is represented by a multi-dimensional subspace of the Hilbert space[78–83]². That is, a physical anyon may have an internal space[26, 29, 85]. Nevertheless, TQFT is not able to sense and probe such internal spaces of anyons because anyons are elementary and indecomposable ingredients in TQFT that have no finer structures. In contrast, in the string-net model—a Hamiltonian extension of the Turaev-Viro TQFT—of topological orders, anyon excitations are indeed concrete excited states in the Hilbert space of the fundamental degrees of freedom of the model. Anyons are represented by orthogonal subspaces (one- or multi-dimensional) in the Hilbert space. We show that in the string-net model, it is possible to transform the fundamental degrees of freedom while preserving topological invariance and the Hamiltonian. Such a transformation would induce a linear transformation of the Hilbert space of the model. When this linear transformation preserves and/or transforms within the subspaces representing the anyons, it is qualified as a gauge transformation of the model. The model and thus the topological order it describes has a corresponding gauge symmetry. When this transformation can turn the anyons to one another, it is a global symmetry transformation of the model and the topological order described by the model[31, 32, 68, 86, 87].

We construct the symmetry transformations of (2+1)-dimensional topological orders in two steps. The first step establishes a duality between different string-net models describing the same topological order. While the fundamental degrees of freedom in a usual gauge theory take value in a group, those in the Turaev-Viro TQFT or a string-net model are simple objects in a fusion category—the input fusion category of the model. Our duality maps two string-net models whose input fusion categories are Morita equivalent to each other[88] and yields a classification of all string-net models describing the same topological order. This duality generalizes the electromagnetic duality[11, 13, 89] in topological orders and topological field theories.

²This is why non-Abelian anyons can support a scheme of quantum computation—topological quantum computation[84].




Figure 1: An example of the part of the lattice of the string-net model. All vertices are trivalent. A tail (wavy line) is attached to an arbitrary edge of every plaquette.

In the second step, symmetry transformations can be constructed when two dual string-net models have isomorphic input fusion categories. In such a case, the duality in the first step can be composed with an isomorphism to form a symmetry transformation on the same string-net model. Such a symmetry transformation can be a global or gauge symmetry transformation depending on whether it permutes anyon species. Such a symmetry, when it is noninvertible, is described by a fusion 2-category.

We now construct and reveal the gauge symmetry transformation of the doubled Fibonacci topological order to illustrate our approach. We shall first briefly review the Fibonacci string-net model and defer the symmetry transformation to the following sections.

3 Fibonacci String-Net Model

For our study, we take the form of the string-net model defined in Ref. [26] because its Hilbert space encompasses the full anyon spectra of the corresponding topological orders. The model is defined on a 2-dimensional trivalent lattice, such as that in Fig. 1. Each plaquette hosts a tail, attached to any of its edges³. Each edge or tail carries a label—fundamental degree of freedom—taking value in the simple objects of the input fusion category \mathcal{F} of the model. The Hilbert space of the model is spanned by all possible assignments of the labels, constrained by that the labels on the three edges (tails) meeting at any vertex must satisfy the fusion rules of \mathcal{F} .

We now dwell on our example: the doubled Fibonacci topological order with the string-net model. The input fusion category of the string-net model, the Fibonacci fusion category, denoted by Fibo , contains two simple objects 1 and τ that are the values of the basic degree of freedom on any edge/tail. These simple objects satisfy the fusion rules

$$\delta_{111} = \delta_{1\tau\tau} = \delta_{\tau 1\tau} = \delta_{\tau\tau 1} = \delta_{\tau\tau\tau} = 1, \quad \delta_{11\tau} = \delta_{1\tau 1} = \delta_{\tau 11} = 0.$$

³The original string-net model in Ref. [18], which bears no such tails, cannot fully describe charge excitations. These added tails record the charges of anyons, thus enlarging the Hilbert space to encompass the complete anyon spectrum.

The Hilbert space $\mathcal{H}_{\text{Fibo}}$ of the model is spanned by all possible assignments of the simple objects 1 and τ on all edges and tails of the lattice, constrained by the fusion rules $\delta_{ijk} = 1$ for the three labels i, j, k on any three edges or tails meeting at any vertex of the lattice.

The Hamiltonian of the Fibonacci string-net model reads

$$H_{\text{Fibo}} := - \sum_{\text{Plaquettes } P} Q_P, \quad (3.1)$$

where the commuting projectors Q_P are defined as

$$Q_P := \frac{1}{1 + \phi^2} (Q_P^1 + \phi Q_P^\tau),$$

$$Q_P^s := \delta_{p,1} \sum_{j_k \in \{1, \tau\}} \left[\prod_{k=0}^5 G_{s j_{k+1} i_{k+1}}^{e_{k+1} i_k j_k} \sqrt{d_{i_k} d_{j_k}} \right] \quad \begin{array}{c} e_2 \\ i_1 \quad i_2 \\ e_1 \quad i_0 \quad e_3 \\ i_6 \quad p \quad i_3 \\ i_5 \quad i_4 \quad e_4 \\ e_6 \quad i_5 \quad e_5 \end{array} \quad \begin{array}{c} e_2 \\ j_1 \quad j_2 \\ e_1 \quad j_0 \quad e_3 \\ j_0 \quad 1 \quad j_3 \\ j_0 \quad j_3 \\ e_6 \quad j_5 \quad j_4 \quad e_4 \\ e_5 \end{array}.$$

Here, $s \in \{1, \tau\}$, $d_1 = 1$, $d_\tau = \phi = (\sqrt{5} + 1)/2$, and the nonzero G -symbols are

$$G_{1111}^{1111} = 1, \quad G_{\tau\tau\tau\tau}^{1111} = \frac{1}{\sqrt{\phi}}, \quad G_{1\tau\tau\tau}^{1\tau\tau\tau} = G_{\tau\tau\tau\tau}^{1\tau\tau\tau} = \frac{1}{\phi}, \quad G_{\tau\tau\tau\tau}^{\tau\tau\tau\tau} = -\frac{1}{\phi^2}, \quad G_{cdn}^{abm} = G_{ncd}^{mab} = G_{abn}^{cdm}.$$

Unlike in the original string-net model, we unify the vertex and plaquette operators into a common Q_P operator. The ground states are common eigenstates of all Q_P operators with +1 eigenvalues. An excited state $|\psi\rangle$ is another common eigenstate that satisfies $Q_P|\psi\rangle = 0$ for one or more plaquettes P , in which we say there reside anyons. We also dub a ground state a trivial excited state, which has a trivial anyon in each plaquette. There are four anyon species of the doubled Fibonacci topological order:

$$1\bar{1}, \quad \tau\bar{1}, \quad 1\bar{\tau}, \quad \tau\bar{\tau}, \quad (3.2)$$

where $1\bar{1}$ is the trivial anyon. The measurement operator Π_P^J measuring whether there is an anyon J in plaquette P is

$$\Pi_P^J := \sum_{s, t \in \{1, \tau\}} \left[\prod_{k=0}^5 G_{s j_{k+1} i_{k+1}}^{e_{k+1} i_k j_k} \sqrt{d_{i_k} d_{j_k}} \right] G_{j_6 s t}^{p i_0 i_6} G_{i_0 t s}^{p j_0 j_6} \times \sqrt{\frac{d_s d_t}{d_p}} z_{pps}^{J; t} \quad \begin{array}{c} e_2 \\ j_1 \quad j_2 \\ e_1 \quad j_0 \quad e_3 \\ j_6 \quad p \quad j_3 \\ j_6 \quad j_3 \\ e_6 \quad j_5 \quad j_4 \quad e_4 \\ e_5 \end{array}, \quad (3.3)$$

where the nonzero elements of z^J tensors are

$$z_{111}^{1\bar{1}; 1} = z_{11\tau}^{1\bar{1}; \tau} = 1;$$

$$\begin{aligned}
z_{\tau\tau 1}^{\tau\bar{1};\tau} &= 1, & z_{\tau\tau\tau}^{\tau\bar{1};1} &= -\frac{\phi}{2} - \frac{i}{2\phi}\sqrt{\phi^2+1}, & z_{\tau\tau\tau}^{\tau\bar{1};\tau} &= -\frac{1}{2\phi} + \frac{i}{2}\sqrt{\phi^2+1}; \\
z_{\tau\tau 1}^{1\bar{\tau};\tau} &= 1, & z_{\tau\tau\tau}^{\tau\bar{1};1} &= -\frac{\phi}{2} + \frac{i}{2\phi}\sqrt{\phi^2+1}, & z_{\tau\tau\tau}^{\tau\bar{1};\tau} &= -\frac{1}{2\phi} - \frac{i}{2}\sqrt{\phi^2+1}; \\
z_{111}^{\tau\bar{\tau};1} &= 1, & z_{11\tau}^{\tau\bar{\tau};\tau} &= -\frac{1}{\phi^2}, & z_{\tau\tau 1}^{\tau\bar{\tau};\tau} &= 1, & z_{\tau\tau\tau}^{\tau\bar{\tau};1} &= 1, & z_{\tau\tau\tau}^{\tau\bar{\tau};\tau} &= \frac{1}{\phi^2}.
\end{aligned}$$

It is the time and right place to bring up the concept of anyon's *internal space*. In contrast to Abelian anyons, which do not have nontrivial internal spaces, a non-Abelian anyon does have an internal space because multiple non-Abelian anyons occupy a well-defined multi-dimensional Hilbert space[78–83]. Such internal spaces are generally hidden in the language of TQFT unless the anyons carry group representations. The string-net model is however able to manifest such internal spaces by representing a non-Abelian anyon on a certain multi-dimensional Hilbert subspace of excited states of the model. Consequently, in the string-net model, an anyon appearing in an excited state is not only labeled by its anyon species J in each plaquette but also by each anyon's internal charge p —the degree of freedom on the tail where the anyon resides. A non-abelian anyon carries more than one charge type, and the gauge symmetry transformation to be constructed will be able to mix the internal charges p while preserving the anyon species J of a non-Abelian anyon.

In the Fibonacci string-net model, there are five allowed pairs (J, p) of anyon species J and its charges p :

$$(1\bar{1}, 1), \quad (1\bar{\tau}, \tau), \quad (\tau\bar{1}, \tau), \quad (\tau\bar{\tau}, 1), \quad (\tau\bar{\tau}, \tau).$$

An anyon $\tau\bar{\tau}$ in a certain plaquette P has two different possible charge choices 1 and τ . Anyons $\tau\bar{1}$ and $1\bar{\tau}$ are both non-abelian anyons but they seem to both carry only one type of charge τ in the string-net model. Nevertheless, as to be seen, to reveal the gauge structure of the doubled Fibonacci phase, it is inevitable to further enlarge the string-net model's Hilbert space in a natural way as follows, such that anyons $\tau\bar{1}$ and $1\bar{\tau}$ will each carry two different internal charges.

4 Enlarging the Hilbert Space and the Duality Map

Given a fusion category \mathcal{F} , there exist *Frobenius algebras* in \mathcal{F} . It is a mathematical theorem[88] that the bimodules—a special class of representations of a given Frobenius algebra \mathcal{A} in \mathcal{F} —form another fusion category $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ that is *categorically Morita equivalent* to \mathcal{F} . This mathematical fact makes us recognize that the string-net model with $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ as its input fusion category and that with \mathcal{F} as its input fusion category are equivalent in that they describe the same topological order. We explicitly establish this equivalence by a duality map \mathcal{D} between these two different models in this article.

The Fibonacci fusion category Fibo has a nontrivial 2-dimensional Frobenius algebra

$$\mathcal{A} := \left\{ \alpha 1 + \beta \tau \mid \alpha, \beta \in \mathbb{C}, 1^2 = 1, 1\tau = \tau 1 = \tau, \tau^2 = 1 + \phi^{-\frac{3}{4}}\tau \right\}, \quad (4.1)$$

where $1, \tau$ are the two simple objects of Fibo , regarded as the two basis elements of the algebra. Frobenius algebra \mathcal{A} has two simple (i.e., irreducible) bimodules: The two-dimensional trivial bimodule $M_1 = (P_1, V_1)$, whose representation space V_1 is

$$V_1 := \{\alpha 1 + \beta \tau \mid \alpha, \beta \in \mathbb{C}\},$$

and the three-dimensional nontrivial bimodule $M_\tau = (P_\tau, V_\tau)$ with the representation space

$$V_\tau := \{\alpha 1 + \beta \tau_1 + \gamma \tau_2 \mid \alpha, \beta, \gamma \in \mathbb{C}\}.$$

Here, τ_1 and τ_2 are both the simple objects τ , but regarded as two different basis elements in V_τ because they are acted on differently by \mathcal{A} ⁴. We refer to the indices 1 and 2 of τ in M_τ the multiplicity label of τ . The functions $P_i : \mathcal{A}^2 \times V_i \times \{1, \tau\} \times V_i \rightarrow \mathbb{C}$ represent a pair of algebra elements $(a, b) \in \mathcal{A}^2$ as rank-3 tensors on the representation spaces V_i , where $i = 1, \tau$. These tensors indicate that two algebra elements a and b in \mathcal{A} act sequentially on $x \in V_i$, transforming it to $z \in V_i$ with the coefficient $\sum_y [P_i]_{xyz}^{ab}$. The intermediate object y varies over $\{1, \tau\}$ to satisfy the fusion rule $\delta_{axy} = \delta_{byz} = 1$. The precise components of the representation tensors can be found in Appendix E.

The Fibonacci string-net model with the input fusion category $\text{Bimod}_{\text{Fibo}}(\mathcal{A})$, denoted as the *dual Fibonacci string-net model*, describes the same topological order—the doubled Fibonacci topological order—as the original Fibonacci string-net model with the input fusion category Fibo . The fundamental degrees of freedom on the edges and tails of the dual model are simple bimodules M_1 and M_τ , which are simple objects in $\text{Bimod}_{\text{Fibo}}(\mathcal{A})$. We construct a duality map \mathcal{D} that embeds the fundamental degrees of freedom of the dual model into the original model, based on the definition of the bimodules:

$$\mathcal{D} \left| M_i \right. := \sum_{a,b,y=1,\tau} \sum_{x,z \in L_{M_i}} [P_i]_{xyz}^{ab} \begin{array}{c} |z \\ \text{---} b \\ y \\ \text{---} a \\ x \end{array}, \quad (4.2)$$

where $M_i = M_1, M_\tau$, and $L_{M_1} = \{1, \tau\}, L_{M_\tau} = \{1, \tau_1, \tau_2\}$. The black line refers to both edges and tails. The red lines are auxiliary tails that will be annihilated by topological moves, resulting in a unitary transformation between the Hilbert spaces of the Fibonacci model and its dual model, which can be understood plaquette by plaquette:

$$\begin{array}{c} E_2 \\ | \\ I_1 \text{---} I_2 \\ | \quad | \\ I_0 \text{---} M \text{---} I_3 \\ | \quad | \\ E_6 \text{---} I_5 \text{---} I_4 \text{---} E_4 \\ | \\ E_5 \end{array} \quad \Rightarrow \quad \sum_{i_k, e_k=1,\tau} \sum_{p \in L_M} \dots \begin{array}{c} e_2 \\ | \\ i_1 \text{---} i_2 \\ | \quad | \\ i_0 \text{---} p \text{---} i_3 \\ | \quad | \\ e_6 \text{---} i_5 \text{---} i_4 \text{---} e_4 \\ | \\ e_5 \end{array}, \quad (4.3)$$

where $I_k, E_k, M \in \{M_1, M_\tau\}$, and “...” denotes the expansion coefficients detailed in Appendix E. After the topological moves, the degree of freedom τ on any edge will cease to

⁴This is completely analogous to the scenario where an irreducible representation of a group can appear more than once in a certain reducible representation of the group. When physics kicks in, the different occurrences of the same irreducible representation are distinguishable.

have any multiplicity index, while that on any tail will still have a multiplicity index if it belongs to L_{M_τ} . Therefore, to make sense of this duality and make it unitary, we are urged to enlarge the Hilbert space $\mathcal{H}_{\text{Fibo}}$ of the original Fibonacci string-net model to \mathcal{H}^* by distinguishing τ_1 and τ_2 on each tail but not on the edges. This enlargement is also physically sound: An anyon excitation resides in a plaquette, in which the tail carries the internal charge of the anyon that reflects the action of Frobenius algebra \mathcal{A} ; the precise action of \mathcal{A} can only be told when different occurrences of τ in the representation (bimodule) of \mathcal{A} are distinguished by multiplicity indices, viz τ_1 and τ_2 . In contrast, the degrees of freedom on edges are pertaining to ground states because any path along edges only has to be a closed loop. At any vertex along such a closed loop, fusion rules are met by definition of the model, and fusion rules regard τ_1 and τ_2 the same⁵.

Due to orthonormality, the actual enlargement is done by embedding $\mathcal{H}_{\text{Fibo}}$ in \mathcal{H}^* as

$$\begin{array}{c} \text{wavy} \\ | \\ 1 \\ \text{wavy} \end{array} \implies \begin{array}{c} \text{wavy} \\ | \\ 1 \\ \text{wavy} \end{array}, \quad \begin{array}{c} \text{wavy} \\ | \\ \tau \\ \text{wavy} \end{array} \implies \left(\frac{1}{2\phi} + \frac{\sqrt{\phi}}{2} \right) \begin{array}{c} \text{wavy} \\ | \\ \tau_1 \\ \text{wavy} \end{array} + \left(\frac{1}{2\phi} - \frac{\sqrt{\phi}}{2} \right) \begin{array}{c} \text{wavy} \\ | \\ \tau_2 \\ \text{wavy} \end{array} \quad (4.4)$$

for each *tail*, as shown in Fig. 2a. The physical Hilbert space $\mathcal{H}_{\text{Fibo}}$ of the string-net model is a subspace of this enlarged Hilbert space \mathcal{H}^* .

5 Symmetry Transformation of the Doubled Fibonacci Topological Order

We can further construct a symmetry transformation of the doubled Fibonacci phase based on the duality \mathcal{D} defined in Eqs. (4.2) and (4.3) because their input fusion categories are isomorphic:

$$\mathcal{F}_{\mathcal{A}} : \text{Fibo} \rightarrow \text{Bimod}_{\text{Fibo}}(\mathcal{A}), \quad 1 \mapsto M_1, \quad \tau \mapsto M_\tau. \quad (5.1)$$

Such isomorphism $\mathcal{F}_{\mathcal{A}}$ induces an isomorphic map $\varphi_{\mathcal{A}}$ between the Fibonacci string-net model and its dual model, and thus a *unitary transformation* \mathcal{G} of the Fibonacci string-net model in the enlarged Hilbert space \mathcal{H}^* :

$$\mathcal{G} := \mathcal{D} \circ \varphi_{\mathcal{A}}, \quad (5.2)$$

where

$$\varphi_{\mathcal{A}} \left| \begin{array}{c} \text{---} \\ | \\ 1 \\ \text{---} \end{array} \right. := \left| \begin{array}{c} \text{---} \\ | \\ M_1 \\ \text{---} \end{array} \right., \quad \varphi_{\mathcal{A}} \left| \begin{array}{c} \text{---} \\ | \\ \tau \\ \text{---} \end{array} \right. := \left| \begin{array}{c} \text{---} \\ | \\ M_\tau \\ \text{---} \end{array} \right. . \quad (5.3)$$

Here, the line refers to both edges and tails. Consequently, the unitary transformation \mathcal{G} transforms the local degrees of freedom 1 and τ on edges (tails) to

$$\left| \begin{array}{c} \text{---} \\ | \\ 1 \\ \text{---} \end{array} \right. \implies \sum_{a,b,x,y,z=1,\tau} [P_1]_{xyz}^{ab} \left| \begin{array}{c} z \\ \text{---} \\ y \\ \text{---} \\ a \\ \text{---} \\ x \end{array} \right. b, \quad \left| \begin{array}{c} \text{---} \\ | \\ \tau \\ \text{---} \end{array} \right. \implies \sum_{\substack{a,b,y=1,\tau \\ x,z=1,\tau_1,\tau_2}} [P_\tau]_{xyz}^{ab} \left| \begin{array}{c} z \\ \text{---} \\ y \\ \text{---} \\ a \\ \text{---} \\ x \end{array} \right. b. \quad (5.4)$$

⁵As an analogy: It makes no sense to question the electric charge in a closed electric flux loop because the Gauss law (analogous to fusion rules) is met everywhere along the loop. Only when the loop is cut open to be a path, one can ask about the charges at the ends of the path where the Gauss law is broken.

The red lines will be annihilated by topological moves.

The unitary transformation \mathcal{G} (5.2) does not preserve the physical Hilbert space $\mathcal{H}_{\text{Fibo}}$ of the string-net model but rather rotates it within the enlarged Hilbert space \mathcal{H}^* . Nevertheless, the physical charges of anyons in the Fibonacci topological order that can be measured are always 1 and τ . After the transformation \mathcal{G} , we must project the transformed states in \mathcal{H}^* back into the physical Hilbert space $\mathcal{H}_{\text{Fibo}}$ using a projector \mathcal{P} :

$$\mathcal{P}\mathcal{H}^* = \mathcal{H}_{\text{Fibo}}. \quad (5.5)$$

Therefore, in the Fibonacci topological order, the *symmetry transformation* cannot be simply the unitary transformation \mathcal{G} . Instead, the appropriate symmetry transformation should be the composition of \mathcal{G} and the projection \mathcal{P} :

$$\tilde{\mathcal{G}} := \mathcal{P}\mathcal{G}\mathcal{P}, \quad (5.6)$$

which preserves the physical Hilbert space $\mathcal{H}_{\text{Fibo}}$. The projected symmetry transformation $\tilde{\mathcal{G}}$ is not unitary and is noninvertible. Specifically, the composition of symmetry transformation $\tilde{\mathcal{G}}$ is given by the projection of multiplying unitary transformation:

$$\tilde{\mathcal{G}}^{(n)} := \mathcal{P}\mathcal{G}^n\mathcal{P} \quad (5.7)$$

for $n > 1$. This composition differs from the traditional way of composing symmetry transformations by simply multiplying the matrix $\tilde{\mathcal{G}}$, which results in $\tilde{\mathcal{G}}^n \neq \tilde{\mathcal{G}}^{(n)}$ for $n > 1$ and does not represent valid symmetry transformations.

We now discuss how the unitary transformation \mathcal{G} (5.2) and the symmetry transformation $\tilde{\mathcal{G}}$ (5.6) transform the spectrum of the Fibonacci string-net model. The unitary transformation \mathcal{G} preserves the anyon species but acts nontrivially on the (enlarged) internal space of each anyon—the local Hilbert space expanded by the basic degrees of freedom $\{1, \tau_1, \tau_2\}$ on the tail where the anyon is located. Different anyon species experience distinct actions. Specifically, we can block-diagonalize the \mathcal{G} (5.2) within the enlarged Hilbert subspaces representing different anyons:

$$\mathcal{G} = \prod_{\text{Plaquettes } P} \sum_{\text{Anyons } J} \mathcal{G}_P^J \Pi_P^J, \quad (5.8)$$

where $J = 1\bar{1}, 1\bar{\tau}, \tau\bar{1}, \tau\bar{\tau}$ is the anyon species, Π_P^J is the measurement operator (3.3) measuring whether there is an anyon J in plaquette P , and \mathcal{G}_P^J is block-diagonal in \mathcal{H}^* , acting nontrivially only on the Hilbert subspace spanned by excited states with the same anyon species in all plaquettes, identical charges in all plaquettes except plaquette P , and varying charges of anyon J on the tail in plaquette P .

Consequently, the symmetry transformation $\tilde{\mathcal{G}}$, being the projection of \mathcal{G} , represents a gauge symmetry transformation that preserves the anyon species. The projected transformation $\tilde{\mathcal{G}}$ can also be block-diagonalized along with \mathcal{G} :

$$\tilde{\mathcal{G}} = \mathcal{P}\mathcal{G}\mathcal{P} = \prod_{\text{Plaquettes } P} \sum_{\text{Anyons } J} \tilde{\mathcal{G}}_P^J \Pi_P^J, \quad \tilde{\mathcal{G}}_P^J := \mathcal{P}\mathcal{G}_P^J\mathcal{P}. \quad (5.9)$$




Figure 2: (a) The physical basic degrees of freedom 1 and τ embedded in the $\{1, \tau_1, \tau_2\}$ enlarged Hilbert space. The black lines refer to the physical degrees of freedom 1 and τ . (2) The action of the unitary symmetry transformations \mathcal{G} (the blue vectors) and the symmetry transformation $\tilde{\mathcal{G}}$ (the orange vectors) for anyon species $\tau\bar{1}, 1\bar{\tau}$. (c) The action of the unitary symmetry transformations \mathcal{G} (the blue vectors) and the symmetry transformation $\tilde{\mathcal{G}}$ (the orange vectors) for anyon species $\tau\bar{\tau}$.

1. The trivial anyon $1\bar{1}$ has only one charge 1 that transforms trivially under the symmetry transformation:

$$\mathcal{G}_P^{1\bar{1}} = \tilde{\mathcal{G}}_P^{1\bar{1}} = 1, \quad \mathcal{G}_P^{1\bar{1}} \begin{array}{c} \vdots \\ 1 \\ \vdots \end{array} = \tilde{\mathcal{G}}_P^{1\bar{1}} \begin{array}{c} \vdots \\ 1 \\ \vdots \end{array} = \begin{array}{c} \vdots \\ 1 \\ \vdots \end{array}. \quad (5.10)$$

The ground states of the doubled Fibonacci topological order are invariant under the gauge symmetry transformation.

2. Anyon $\tau\bar{1}$ has only one charge τ in the string-net model. Nevertheless, we have enlarged the degree of freedom τ on a tail to a 2-dimensional space spanned by degrees of freedom τ_1 and τ_2 . We can now express the symmetry transformation \mathcal{G} (5.2) in this enlarged internal space for $\tau\bar{1}$ anyon:

$$\begin{aligned} \mathcal{G}_P^{\tau\bar{1}} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{\phi\sqrt{\phi^2+1}-1}{2\phi\sqrt{\phi}} & \frac{\sqrt{\phi^2+1}+\phi^2}{2\phi^2} \\ 0 & -\frac{\sqrt{\phi^2+1}+\phi^2}{2\phi^2} & \frac{\phi\sqrt{\phi^2+1}-1}{2\phi\sqrt{\phi}} \end{pmatrix}, \\ \tilde{\mathcal{G}}_P^{\tau\bar{1}} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & \frac{(\phi\sqrt{\phi^2+1}-1)(\sqrt{\phi+1})}{4\phi^2} & \frac{1-\phi\sqrt{\phi^2+1}}{4\phi^2\sqrt{\phi}} \\ 0 & \frac{1-\phi\sqrt{\phi^2+1}}{4\phi^2\sqrt{\phi}} & \frac{(\phi\sqrt{\phi^2+1}-1)(\sqrt{\phi-1})}{4\phi^2} \end{pmatrix}, \\ \mathcal{G}_P^{\tau\bar{1}} \begin{array}{c} \vdots \\ \tau \\ \vdots \end{array} &= \begin{array}{c} \vdots \\ \tau_{\tau\bar{1}} \\ \vdots \end{array} = \frac{\sqrt{5}-1}{2\sqrt{\phi}} \begin{array}{c} \vdots \\ \tau_1 \\ \vdots \end{array} - \frac{\sqrt{5}+1}{2\sqrt{\phi}} \begin{array}{c} \vdots \\ \tau_2 \\ \vdots \end{array}, \end{aligned} \quad (5.11)$$

$$\tilde{\mathcal{G}}_P^{\tau\bar{1}} \begin{array}{c} \text{wavy line} \\ \tau \end{array} = \begin{array}{c} \text{wavy line} \\ \tau_{\bar{1}} \end{array} = \frac{\phi\sqrt{\phi^2+1}-1}{2\phi\sqrt{\phi}} \begin{array}{c} \text{wavy line} \\ \tau \end{array} .$$

Here, the physical degree of freedom τ of the anyon $\tau\bar{1}$ is a superposition (4.4) of τ_1 and τ_2 . The unitary transformation \mathcal{G} (5.2) rotates this physical charge τ of the $\tau\bar{1}$ anyon out of the physical Hilbert space $\mathcal{H}_{\text{Fibo}}$, necessitating the application of the projected transformation $\tilde{\mathcal{G}}_P^{\tau\bar{1}}$ to preserve the physical Hilbert space, as shown in Fig. 2b. Note that because $\det(\tilde{\mathcal{G}}_P^{\tau\bar{1}}) = 0$, the matrix $\tilde{\mathcal{G}}_P^{\tau\bar{1}}$ does not have an inverse matrix $[\tilde{\mathcal{G}}_P^{\tau\bar{1}}]^{-1}$. For this reason, the symmetry transformation $\tilde{\mathcal{G}}$ (5.6) represents a noninvertible symmetry.

3. Anyon $1\bar{\tau}$ also has a single charge τ . The unitary transformation \mathcal{G} (5.2) rotates the physical charge τ (4.4) of the $1\bar{\tau}$ anyon within the $\{\tau_1, \tau_2\}$ space but in a different way compared to the $\tau\bar{1}$ anyon in Eqs. (5.11):

$$\begin{aligned} \mathcal{G}_P^{1\bar{\tau}} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{\phi\sqrt{\phi^2+1}+1}{2\phi\sqrt{\phi}} & \frac{\phi\sqrt{\phi}-\sqrt[4]{5}}{2\phi\sqrt{\phi}} \\ 0 & \frac{\sqrt[4]{5}-\phi\sqrt{\phi}}{2\phi\sqrt{\phi}} & -\frac{\phi\sqrt{\phi^2+1}+1}{2\phi\sqrt{\phi}} \end{pmatrix}, \\ \tilde{\mathcal{G}}_P^{1\bar{\tau}} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & -\frac{(\phi\sqrt{\phi^2+1}+1)(\sqrt{\phi}+1)}{4\phi^2} & \frac{1+\phi\sqrt{\phi^2+1}}{4\phi^2\sqrt{\phi}} \\ 0 & \frac{1+\phi\sqrt{\phi^2+1}}{4\phi^2\sqrt{\phi}} & -\frac{(\phi\sqrt{\phi^2+1}+1)(\sqrt{\phi}-1)}{4\phi^2} \end{pmatrix}, \quad (5.12) \\ \mathcal{G}_P^{1\bar{\tau}} \begin{array}{c} \text{wavy line} \\ \tau \end{array} &= \begin{array}{c} \text{wavy line} \\ \tau_{1\bar{\tau}} \end{array} = -\frac{\sqrt[4]{5}+1}{2\sqrt{\phi}} \begin{array}{c} \text{wavy line} \\ \tau_1 \end{array} + \frac{\sqrt[4]{5}-1}{2\sqrt{\phi}} \begin{array}{c} \text{wavy line} \\ \tau_2 \end{array}, \\ \tilde{\mathcal{G}}_P^{1\bar{\tau}} \begin{array}{c} \text{wavy line} \\ \tau \end{array} &= \begin{array}{c} \text{wavy line} \\ \tau_{1\bar{\tau}} \end{array} = -\frac{\phi\sqrt{\phi^2+1}+1}{2\phi\sqrt{\phi}} \begin{array}{c} \text{wavy line} \\ \tau \end{array}. \end{aligned}$$

The transformations are depicted in Fig. 2b. The matrix $\tilde{\mathcal{G}}_P^{1\bar{\tau}}$ is noninvertible.

4. Anyon $\tau\bar{\tau}$ has two gauge charge 1 and τ in the string-net model. In the enlarged 3-dimensional space spanned by basic degrees of freedom 1, τ_1 , and τ_2 , the unitary transformation \mathcal{G} (5.2) rotates these two physical charges 1 and τ (4.4) (see Fig. 2c):

$$\mathcal{G}_P^{\tau\bar{\tau}} = \begin{pmatrix} \frac{1}{\phi^2} & \frac{\phi^2\sqrt{5}-\sqrt{\phi}}{2\phi^4}\sqrt[4]{5} & -\frac{\phi^2\sqrt{5}+\sqrt{\phi}}{2\phi^4}\sqrt[4]{5} \\ \frac{\sqrt[4]{5}+\phi\sqrt{\phi^2+1}}{2\phi^2} & \frac{1-\phi^2\sqrt{5}\phi}{2\phi^4} & \frac{\phi\sqrt{5}-2\sqrt{\phi}}{2\phi^3} \\ \frac{\sqrt[4]{5}-\phi\sqrt{\phi^2+1}}{2\phi^2} & -\frac{\phi\sqrt{5}+2\sqrt{\phi}}{2\phi^3} & -\frac{1+\phi^2\sqrt{5}\phi}{2\phi^4} \end{pmatrix},$$

$$\begin{aligned}
\tilde{\mathcal{G}}_P^{\tau\bar{\tau}} &= \begin{pmatrix} \frac{1}{\phi^2} & \frac{\sqrt[4]{5+\phi}\sqrt{\phi^2+1}}{2\phi^2\sqrt{\phi}} & \frac{\sqrt[4]{5-\phi}\sqrt{\phi^2+1}}{2\phi^2\sqrt{\phi}} \\ \frac{\sqrt[4]{5+\phi}\sqrt{\phi^2+1}}{2\phi^2} & -\frac{1+\sqrt{\phi}}{2\phi^3} & \frac{1}{2\phi^3\sqrt{\phi}} \\ \frac{\sqrt[4]{5-\phi}\sqrt{\phi^2+1}}{2\phi^2} & \frac{1}{2\phi^3\sqrt{\phi}} & \frac{1-\sqrt{\phi}}{2\phi^3} \end{pmatrix}, \\
\mathcal{G}_P^{\tau\bar{\tau}} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1 \end{array} &= \tilde{\mathcal{G}}_P^{\tau\bar{\tau}} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1 \end{array} = \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1_{\tau\bar{\tau}} \end{array} = \frac{1}{\phi^2} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1 \end{array} + \frac{\sqrt[4]{5}}{\phi} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau \end{array}, \quad (5.13) \\
\mathcal{G}_P^{\tau\bar{\tau}} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau \end{array} &= \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau_{\tau\bar{\tau}} \end{array} = \frac{\sqrt{\phi^2+1}}{\phi^2} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1 \end{array} - \frac{\sqrt{\phi}}{\phi^2} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau_1 \end{array} - \frac{\sqrt{\phi}}{\phi^2} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau_2 \end{array}, \\
\tilde{\mathcal{G}}_P^{\tau\bar{\tau}} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau \end{array} &= \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tilde{\tau}_{\tau\bar{\tau}} \end{array} = \frac{\sqrt{\phi^2+1}}{\phi^2} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } 1 \end{array} - \frac{\sqrt{\phi}}{\phi^3} \begin{array}{c} \text{wavy} \\ \text{line} \\ \text{with} \\ \text{charge } \tau \end{array}.
\end{aligned}$$

When $\mathcal{G}_P^{\tau\bar{\tau}}$ acts on an excited state with a $\tau\bar{\tau}$ anyon in plaquette P , the charges on the tail in plaquette P transform into a superposition of charges 1, τ_1 , and τ_2 . This transformation does not mean that only the degrees of freedom on the tail in P change while leaving the degrees of freedom on all other edges and tails invariant. Instead, the excited states are transformed into a superposition of *excited states* where the anyon $\tau\bar{\tau}$ in plaquette P has charges 1 and τ (or 1, τ_1 , and τ_2).

The matrix $\tilde{\mathcal{G}}_P^{\tau\bar{\tau}}$ is noninvertible.

We have to note that while the explicit representation matrices for the symmetry transformations are defined in the string-net model, the symmetry itself is an intrinsic property of the doubled Fibonacci topological order and is independent of the specific model realization. We will discuss the structure of this symmetry in the next section and show that it is a categorical gauge symmetry described by a fusion 2-category.

6 Fibonacci Categorical Gauge Symmetry

We now show that the symmetry of the doubled Fibonacci topological order is a *categorical gauge symmetry* characterized by a fusion 2-category—the Fibonacci fusion 2-category.

Recall that in a usual gauge theory with gauge group G , a gauge-field value $g \in G$ is transformed to $g' = hgh^{-1} \in G$ by a gauge transformation characterized by $h \in G$. The gauge group G is both the space of the gauge field and is space of the gauge transformations. The question in our case is: What is the structure of the symmetry of the doubled Fibonacci topological order that is analogous to the gauge group G together with its symmetry transformation in a usual gauge theory?

Looking back to how the fundamental degrees of freedom are transformed as in Eq. (5.4) by \mathcal{G} , it seems that the two sides of the equation have different lattice structures, although in the end the red lines on the right-hand-side will be annihilated. But the two sides in fact have the same lattice structure even before annihilating the red lines for both

mathematical and physical reasons. Mathematically, in the Fibonacci fusion category, the simple objects 1 and τ are simple bimodules over the trivial Frobenius algebra $\mathcal{A}_0 = \mathbb{C}[1]$, so we have the identification:

$$\mathbf{Fibo} = \mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}_0),$$

an edge or tail of the original Fibonacci model is too attached with two red lines:

$$1 \begin{array}{|c} \hline 1 \\ \hline 1 \\ \hline 1 \\ \hline \end{array} \Rightarrow \sum_{a,b,x,y,z=1,\tau} [P_1]_{xyz}^{ab} \begin{array}{|c} \hline z \\ \hline y \\ \hline x \\ \hline \end{array}, \quad 1 \begin{array}{|c} \hline \tau \\ \hline \tau \\ \hline \tau \\ \hline \end{array} \Rightarrow \sum_{\substack{a,b,y=1,\tau \\ x,z=1,\tau_1,\tau_2}} [P_\tau]_{xyz}^{ab} \begin{array}{|c} \hline z \\ \hline y \\ \hline x \\ \hline \end{array}; \quad (6.1)$$

however, since these two red lines are labeled by the trivial object 1 , they are omitted for simplicity. Physically, the simple objects of the input fusion category of a string-net model are the pure charges, which are defined concerning the trivial flux, characterized by the trivial Frobenius algebra of the input fusion category. When the Fibonacci fusion category is the input, the trivial flux is characterized by the \mathcal{A}_0 defined above. When $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A})$ is the input fusion category, the trivial flux is characterized by \mathcal{A} defined in Eq. (4.1). Hence, the right-hand side of transformation (6.1), whose red lines carry the elements of \mathcal{A} , are precisely how pure charges M_1, M_τ appear in the string-net model with input fusion category $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A})$. The mathematical and physical reasonings comply with each other. Therefore, the lattice structures before and after transformation \mathcal{G} (5.2) are the same.

It turns out that

$$\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}) \subset \mathbf{Fibo} \quad \text{and} \quad \mathbf{Fibo} \cong \mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}).$$

The bimodules M_1, M_τ are composite objects in \mathbf{Fibo} . In words, our transformation \mathcal{G} (5.2) transforms the string-net model with input fusion category $\mathbf{Fibo} = \mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}_0)$ to to be that with input subcategory $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}) \subset \mathbf{Fibo}$. The input data \mathbf{Fibo} is invariant. This is indeed a gauge symmetry transformation because it preserves the Hamiltonian \mathcal{H}^* and transforms the internal spaces of anyons only.

Inspired by the above discussions, we find that a fusion 2-category—the **Fibonacci fusion 2-category** exists to describe this gauge symmetry coherently. It is defined⁶ by the following three ingredients:

1. The objects in the Fibonacci fusion 2-category are bimodule categories over Frobenius algebras in \mathbf{Fibo} . These bimodule categories are subcategories of and isomorphic to \mathbf{Fibo} . Two such objects are $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}_0) = \mathbf{Fibo}$ and $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}) \subset \mathbf{Fibo}$.
2. The 1-morphisms are isomorphism functors between objects. Now that all bimodule categories in the Fibonacci fusion 2-category are isomorphic, we only need to define the functors $\mathcal{F}'_{\mathcal{A}}$ from $\mathbf{Fibo} = \mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}_0)$ to the other bimodule category $\mathbf{Bimod}_{\mathbf{Fibo}}(\mathcal{A}')$. That is, each 1-morphism is labeled by a Frobenius algebra in \mathbf{Fibo} . Two special examples of 1-morphisms are the identity functor

$$\text{id}(1) = 1, \quad \text{id}(\tau) = \tau,$$

⁶For a general definition and properties of fusion 2-categories, refer to Ref. [59].

and $\mathcal{F}_{\mathcal{A}}$ defined in Eq. (5.1):

$$\mathcal{F}_{\mathcal{A}}(1) = M_1, \quad \mathcal{F}_{\mathcal{A}}(\tau) = M_{\tau}.$$

3. The 2-morphisms are natural transformations of 1-morphisms. We are particularly interested in two types of 2-morphisms: the composition of 1-morphisms and the direct sum of 1-morphisms. Since all bimodule categories are isomorphic, the composition of 1-morphisms is well-defined and maps simple objects 1 and $\tau \in \mathbf{Fibo}$ to bimodules over Frobenius algebras in the bimodule categories. These compositions are also 1-morphisms because the bimodules in the bimodule categories are composite objects in the parent fusion category \mathbf{Fibo} and are bimodules over some Frobenius algebras in \mathbf{Fibo} . The explicit structures of these compositions are complex and will not be discussed here.

Given two 1-morphisms $\mathcal{F}_{\mathcal{A}_1}$ and $\mathcal{F}_{\mathcal{A}_2}$, the direct sum is

$$(\mathcal{F}_{\mathcal{A}_1} \oplus \mathcal{F}_{\mathcal{A}_2})(1) = \mathcal{F}_{\mathcal{A}_1}(1) \oplus \mathcal{F}_{\mathcal{A}_2}(1), \quad (\mathcal{F}_{\mathcal{A}_1} \oplus \mathcal{F}_{\mathcal{A}_2})(\tau) = \mathcal{F}_{\mathcal{A}_1}(\tau) \oplus \mathcal{F}_{\mathcal{A}_2}(\tau).$$

The direct sums $\mathcal{F}_{\mathcal{A}_1}(1) \oplus \mathcal{F}_{\mathcal{A}_2}(1)$ and $\mathcal{F}_{\mathcal{A}_1}(\tau) \oplus \mathcal{F}_{\mathcal{A}_2}(\tau)$ are the two simple bimodules over Frobenius algebra $\mathcal{A}_1 \oplus \mathcal{A}_2$.

In the framework of the Fibonacci fusion 2-category, the transformation \mathcal{G} (5.2) is a representation of the 1-morphism ϕ over the enlarged Hilbert space \mathcal{H}^* :

$$\rho(\text{id}) = \mathbf{1}, \quad \rho(\mathcal{F}_{\mathcal{A}}) = \mathcal{G},$$

which is compatible with the 2-morphisms:

$$\rho(\mathcal{F}_1 \circ \mathcal{F}_2) = \rho(\mathcal{F}_2)\rho(\mathcal{F}_1), \quad \rho(\mathcal{F}_1 \oplus \mathcal{F}_2) = \rho(\mathcal{F}_2) + \rho(\mathcal{F}_1).$$

Now, to answer the question raised at the beginning of this subsection, what is analogous to the gauge group G together with its symmetry transformation in a usual gauge theory is the Fibonacci fusion 2-category. The gauge symmetry of the doubled Fibonacci topological order, generated by the symmetry transformation $\tilde{\mathcal{G}}$ —the projection of the representation matrix \mathcal{G} of the 1-morphism $\mathcal{F}_{\mathcal{A}}$ in the physical Hilbert space $\mathcal{H}_{\mathbf{Fibo}}$ —is described by the Fibonacci fusion 2-category. Therefore, we shall call the symmetry of the doubled Fibonacci topological order the Fibonacci 2-categorical gauge symmetry, which is the gauge symmetry of the doubled Fibonacci topological order, the Fibonacci Turaev-Viro TQFT, and the Fibonacci string-net model.

Note that the 2-category structure is necessary to describe the composition of gauge transformations because $\tilde{\mathcal{G}}^{(n)}$ is never the $\mathbf{1}$ for $n \neq 0$. At the categorical level, by applying an arbitrary number of compositions of the 1-morphism $\mathcal{F}_{\mathcal{A}}$, the fundamental degrees of freedom of the string-net model—the simple objects in some bimodule fusion categories—always remain as objects within \mathbf{Fibo} but will never return to the original degrees of freedom 1 and τ . It will generate infinitely many objects in the Fibonacci fusion 2-category.

Acknowledgments

The authors also thank Davide Gaiotto, Lukas Mueller, Ling-Yan Hung, Yuting Hu, Yinan Wang, and Chenjie Wang for inspiring and helpful discussions. YW is supported by the General Program of Science and Technology of Shanghai No. 21ZR1406700, and the Shanghai Municipal Science and Technology Major Project (Grant No. 2019SHZDZX01). The authors are grateful for the hospitality of the Perimeter Institute during his visit, where the main part of this work is done. This research was supported in part by the Perimeter Institute for Theoretical Physics. Research at Perimeter Institute is supported by the Government of Canada through the Department of Innovation, Science and Economic Development and by the Province of Ontario through the Ministry of Research, Innovation and Science.

A Review of the Extended String-net Model

In this appendix, we briefly review the string-net model defined in Ref. [26]. The string-net model is an exactly solvable Hamiltonian model defined on a 2-dimensional lattice. An example lattice is depicted in Fig. 1. All vertices are trivalent. Within each plaquette of the lattice, a tail is attached to an arbitrary edge of the plaquette, pointing inward. We will later demonstrate that different choices of the edge to which the tail is attached are equivalent. In general cases, each edge and tail is oriented, and different choices of directions are equivalent. Nevertheless, for the case of the Fibonacci string-net model presented in the main body, different direction choices are the same, so we omit the directions of edges and tails in the main body.

The input data of the string-net model is a fusion category \mathcal{F} , described by a finite set $L_{\mathcal{F}}$, whose elements are called *simple objects*, equipped with three functions $N : L_{\mathcal{F}}^3 \rightarrow \mathbb{N}$, $d : L_{\mathcal{F}} \rightarrow \mathbb{R}$, and $G : L_{\mathcal{F}}^6 \rightarrow \mathbb{C}$. The function N sets the *fusion rules* of the simple objects, satisfying

$$\sum_{e \in L_{\mathcal{F}}} N_{ab}^e N_{ec}^d = \sum_{e \in L_{\mathcal{F}}} N_{ae}^d N_{bc}^e, \quad N_{ab}^c = N_{c^*a}^{b^*}.$$

There exists a special simple object $0 \in L_{\mathcal{F}}$, called the *trivial object*, such that for any $a, b \in L_{\mathcal{F}}$,

$$N_{0a}^b = N_{0b}^a = \delta_{ab},$$

where δ is the Kronecker symbol. For each $a \in L_{\mathcal{F}}$, there exists a unique simple object $a^* \in L_{\mathcal{F}}$, called the *opposite object* of a , such that

$$N_{ab}^0 = N_{ba}^0 = \delta_{ba^*}.$$

We only consider the case where for any $a, b, c \in L_{\mathcal{F}}$, $N_{ab}^c = 0$ or 1. In this case, we define

$$\delta_{abc} = N_{ab}^{c^*} \in \{0, 1\}.$$

The basic configuration of the string-net model is established by labeling each edge and tail with a simple object in $L_{\mathcal{F}}$, subject to the constraint on all vertices that $\delta_{ijk} = 1$ for the three incident edges or tails meeting at this vertex, all pointing toward the vertex and respectively counterclockwise labeled by $i, j, k \in L_{\mathcal{F}}$. We can reverse the direction of any edge or tail and simultaneously conjugate its label as $j \rightarrow j^*$, which keeps the configuration invariant. The Hilbert space \mathcal{H} of the model is spanned by all possible configurations of these labels on the edges and tails.

The function d returns the *quantum dimensions* of the simple objects in $L_{\mathcal{F}}$. It is the largest eigenvalues of the fusion matrix and forms the 1-dimensional representation of the fusion rule.

$$d_a d_b = \sum_{c \in L_{\mathcal{F}}} N_{ab}^c d_c.$$

In particular, $d_1 = 1$, and for any $a \in L_{\mathcal{F}}$, $d_a = d_{a^*} \neq 0$.

The function G defines the *6j-symbols* of the fusion algebra. It satisfies

$$\sum_n d_n G_{v^* u^* a}^{pqn} G_{j^* i^* b}^{uvn} G_{q^* p^* c}^{ijn} = G_{i^* p u^*}^{abc} G_{v q^* j^*}^{c^* b^* a^*}, \quad \sum_n d_n G_{kln}^{ijp} G_{l^* k^* n}^{j^* i^* q} = \frac{\delta_{pq^*}}{d_p} \delta_{ijp} \delta_{klq},$$

$$G_{kln}^{ijm} = G_{ijn^*}^{klm^*} = G_{lkn^*}^{jim} = G_{nk^* l^*}^{mij} = \alpha_m \alpha_n \overline{G_{l^* k^* n^*}^{j^* i^* m^*}},$$

where $\alpha_a = \text{sgn}(a)$.

The Hamiltonian of the string-net model reads

$$H := - \sum_{\text{Plaquettes } P} Q_P, \quad Q_P := \frac{1}{D} \sum_{s \in L_{\mathcal{F}}} Q_P^s, \quad D := \sum_{a \in L_{\mathcal{F}}} d_a^2, \quad (\text{A.1})$$

where the plaquette operator Q_P^s acts on edges surrounding plaquette P and has the following matrix elements on a hexagonal plaquette⁷:

$$Q_P^s := \delta_{p,0} \sum_{j_k \in L_{\mathcal{F}}} \prod_{k=1}^6 \left(\sqrt{d_{i_k} d_{j_k}} G_{s j_{k+1}^* j_k}^{e_k i_k i_{k+1}^*} \right)$$

It turns out that

$$(Q_P^s)^\dagger = Q_P^{s^*}, \quad Q_P^r Q_P^s = \sum_{t \in L_{\mathcal{F}}} N_{rs}^t Q_P^t, \quad Q_P^2 = Q_P, \quad Q_{P_1} Q_{P_2} = Q_{P_2} Q_{P_1}.$$

The summands Q_P in Hamiltonian H are commuting projectors, so the Hamiltonian is exactly solvable. The ground-state subspace \mathcal{H}_0 of the system is the projection

$$\mathcal{H}_0 = \left[\prod_{\text{Plaquettes } P} Q_P \right] \mathcal{H}. \quad (\text{A.2})$$

If the lattice has the sphere topology, the model has a unique ground state $|\Phi\rangle$ up to scalars.

⁷We only show the actions of Q_P operator on a hexagonal plaquette. The matrix elements of Q_P operators on other types of plaquettes are defined similarly.

A.1 Topological Features

We briefly review the topological nature of the ground-state subspace of the string-net model defined in Ref. [26]. Any two lattices with the same topology can be transformed into each other by so-called *Pachner moves*. There are unitary linear maps between the Hilbert spaces of two string-net models with the same input fusion category on different lattices associated with these moves, denoted as \mathcal{T} . The ground states are invariant under such linear transformations. There are three kinds of elementary Pachner moves, whose corresponding linear transformations are:

$$\begin{aligned}
 \mathcal{T} \begin{array}{c} a \quad c \\ \diagdown \quad / \\ m \\ \diagup \quad \diagdown \\ b \quad d \end{array} &= \sum_{n \in L_{\mathcal{F}}} \sqrt{d_m d_n} G_{kln}^{ijm} \begin{array}{c} a \quad c \\ \diagdown \quad / \\ n \\ \diagup \quad \diagdown \\ b \quad d \end{array}, \\
 \mathcal{T} \begin{array}{c} b \\ \uparrow \\ x \otimes y \\ \downarrow \\ a \end{array} &= \sqrt{\frac{d_x d_y}{d_i}} \delta_{ij} \delta_{xyi^*} \begin{array}{c} | \\ a \end{array}, \\
 \mathcal{T} \begin{array}{c} | \\ a \end{array} &= \frac{1}{D} \sum_{xy \in L_{\mathcal{F}}} \sqrt{\frac{d_x d_y}{d_i}} \delta_{xyi^*} \begin{array}{c} a \\ \uparrow \\ x \otimes y \\ \downarrow \\ a \end{array}.
 \end{aligned} \tag{A.3}$$

Here we use red “ \times ” to mark the plaquettes to contract. Any other Pachner moves and their corresponding linear transformations of Hilbert spaces are compositions of these three elementary moves. Given initial and final lattices, there are multiple ways to compose these elementary moves, but different ways result in the same transformation matrices on the ground-state Hilbert space.

We have also noted that different selections of the edge to which the tail is attached are equivalent. These variations lead to distinct lattice configurations and, consequently, different Hilbert spaces for the lattice model. The equivalence of states in such Hilbert spaces is established by the following linear transformation \mathcal{T}' :

$$\mathcal{T}' \begin{array}{c} \downarrow \\ \diagdown \quad / \\ i_1 \quad p \\ \diagup \quad \diagdown \\ i_0 \end{array} = \sum_{j \in L_{\mathcal{F}}} \sqrt{d_{i_1} d_j} G_{i_0 p^* j}^{i_2^* e_1 i_1} \begin{array}{c} \downarrow \\ \diagdown \quad / \\ j \quad p \\ \diagup \quad \diagdown \\ i_0 \end{array}. \tag{A.4}$$

The states where tails attach to other edges can be obtained recursively in this manner.

For convenience, in certain instances, we will temporarily incorporate auxiliary states with multiple tails within a single plaquette. These states, despite having multiple tails in

one plaquette, are all equivalent to states within the Hilbert space:

$$\text{Diagram 1} = \sum_{u \in L_{\mathcal{F}}} \sqrt{d_j d_p} G_{i_0 s^* p}^{r^* i_1^* j} \text{Diagram 2} . \quad (\text{A.5})$$

A.2 Excited States

An *excited state* $|\varphi\rangle$ of the string-net model is an eigenstate of the Hamiltonian such that $Q_P|\varphi\rangle = 0$ at some plaquettes P . In such a state, there are *anyons* in these plaquettes P . We also refer to the ground states as trivial excited states, in which there are only *trivial anyons* in all plaquettes. We assume the sphere topology, in which the model has a unique ground state; nevertheless, the results in this section apply to other topologies.

We start with the simplest excited states with a pair of anyons in two *adjacent* plaquettes with a common edge E . This state can be generated by ribbon operator $W_E^{J;pq}$:

$$W_E^{J;pq} \text{Diagram 1} := \sum_{k \in L_{\mathcal{F}}} \sqrt{\frac{d_k}{d_j}} \bar{z}_{pqj}^{J;k} \text{Diagram 2} , \quad (\text{A.6})$$

where j is the label on edge E , and \bar{z} is the complex conjugate. Here, the four-indexed tensor $z_{pqj}^{J;k}$ is called the *half-braiding tensor*, defined by the following equation:

$$\frac{\delta_{jt} N_{rs}^t}{d_t} z_{pqt}^{J;w} = \sum_{u, l, v \in L_{\mathcal{F}}} z_{lqr}^{J;v} z_{pls}^{J;u} \cdot d_u d_v G_{p^* w u}^{r^* s^* t} G_{q w^* v}^{s r j^*} G_{r v^* w}^{s^* u l^*} .$$

The label J , called the *anyon species*, labels different minimal solutions of the z tensor that cannot be the sum of any other nonzero solutions. The ribbon operator $W_E^{J;pq}$ creates in the two adjacent plaquettes a pair of anyons J^* and J with charges p^* and q . An anyon species J may have multiple possible charges p , causing multiple excited states of the string-net model to represent the same anyon. Categorically, anyon species J are labeled by simple objects in the *center* of the input fusion category \mathcal{F} , a modular tensor category whose categorical data record all topological properties of the topological order that the string-net model describes:

$$J \in L_{\mathcal{Z}(\mathcal{F})} .$$

States with two quasiparticles in two non-adjacent plaquettes are generated by ribbon operators along longer paths. These ribbon operators result from concatenating shorter ribbon operators. For example, to create two quasiparticles J^* and J with charges p_0^* and p_n in two non-adjacent plaquettes P_0 and P_n , we can choose a sequence of plaquettes

(P_0, P_1, \dots, P_n) , where P_i and P_{i+1} are adjacent plaquettes with their common edge E_i . The ribbon operator $W_{P_0 P_n}^{J;p_0 p_n}$ is

$$W_{P_0 P_n}^{J;p_0 p_n} := \left[\sum_{p_1 p_2 \dots p_{n-1} \in L_{\mathcal{F}}} \prod_{k=1}^{n-1} \left(d_{p_k} B_{P_k} W_{E_k}^{J;p_k p_{k+1}} \right) \right] W_{E_0}^{J;p_0 p_1}.$$

Different choices of plaquette paths (P_0, P_1, \dots, P_n) give the same operator $W_{P_0 P_n}^{J;p_0 p_n}$ if these sequences can deform continuously from one to another. Following the same method, we can also define the creation operator of three or more anyons.

At the end of this appendix, we define the measurement operator Π_P^J measuring whether there is an anyon J excited in plaquette P :

$$\Pi_P^J := \sum_{s,t \in L_{\mathcal{F}}} \frac{d_s d_t}{d_p} z_{pps}^{J;t}. \quad (\text{A.7})$$

The set of measurement operators are orthonormal and complete:

$$\Pi_P^J \Pi_P^K = \delta_{JK} \Pi_P^J, \quad \sum_{J \in L_{z(\mathcal{F})}} \Pi_P^J = \mathbf{1}.$$

B Frobenius Algebras and Bimodules

It is a mathematical theorem [88] that two fusion categories \mathcal{F} and \mathcal{F}' have isomorphic centers if and only if they are *categorically Morita equivalent*. That is, two string-net models with categorically Morita equivalent input fusion categories describe the same topological order. Category theory also tells that if a fusion category \mathcal{F}' is categorically Morita equivalent to \mathcal{F} , there must be a *Frobenius algebra* \mathcal{A} in \mathcal{F} , such that \mathcal{F}' is isomorphic to the *bimodule category* over \mathcal{A} in \mathcal{F} :

$$\mathcal{F}' \cong \text{Bimod}_{\mathcal{F}}(\mathcal{A}). \quad (\text{B.1})$$

Therefore, different string-net models describing the same topological order are classified by all Frobenius algebras \mathcal{A} in a particular input fusion category \mathcal{F} . Such equivalent models have bimodule categories $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ as their input fusion categories. We can establish the duality maps between these equivalent models. In this appendix, we briefly review the definition of Frobenius algebras in a given fusion category and their bimodules and leave the duality maps for the next appendix.

B.1 Frobenius Algebra

A Frobenius algebra \mathcal{A} in a fusion category \mathcal{F} is characterized by a pair of functions (n, f) . Function $n : L_{\mathcal{F}} \rightarrow \mathbb{N}$ returns the *multiplicity* n_a of $a \in L_{\mathcal{F}}$ appearing in the Frobenius

algebra \mathcal{A} , satisfying $n_a = n_{a^*}$. The simple objects of \mathcal{A} are labeled by pairs a_α , where $a \in L_{\mathcal{F}}$ satisfies $n_a > 0$, and $\alpha = 1, 2, \dots, n_a$ is the *multiplicity index*. We denote the set of all simple objects in \mathcal{A} as $L_{\mathcal{A}}$.

The algebraic multiplication of \mathcal{A} is given by a function $f : L_{\mathcal{A}}^3 \rightarrow \mathbb{C}$, satisfying:

$$\sum_{t_\tau \in L_{\mathcal{A}}} f_{r_\rho s_\sigma t_\tau} f_{a_\alpha b_\beta t_\tau^*} G_{abc}^{rst} \sqrt{d_c d_t} = \sum_{\gamma=1}^{n_c} f_{a_\alpha c_\gamma s_\sigma} f_{r_\rho c_\gamma^* b_\beta} ,$$

$$\sum_{a_\alpha b_\beta \in L_{\mathcal{A}}} f_{a_\alpha b_\beta c_\gamma} f_{b_\beta^* a_\alpha^* c_\gamma^*} \sqrt{d_a d_b} = d_{\mathcal{A}} \sqrt{d_c}, \quad f_{a_\alpha b_\beta c_\gamma} = f_{b_\beta c_\gamma a_\alpha}, \quad f_{0 a_\alpha b_\beta} = \delta_{ab^*} \delta_{\alpha\beta},$$
(B.2)

where

$$d_{\mathcal{A}} := \sum_{a \in L_{\mathcal{F}}} n_a d_a$$
(B.3)

is the *quantum dimension* of the Frobenius algebra \mathcal{A} . This definition aligns with the one in the main body, where a Frobenius algebra \mathcal{A} is expressed as a vector space spanned by simple objects, and the algebraic multiplication rule is given by function f :

$$\mathcal{A} = \mathbb{C}[L_{\mathcal{A}}], \quad a_\alpha b_\beta = \sum_{c_\gamma \in L_{\mathcal{A}}} f_{a_\alpha b_\beta c_\gamma^*} c_\gamma \in \mathbb{C}[L_{\mathcal{A}}].$$

For convenience, in a lattice model, we use red edges or tails to indicate that this edge or tail is labeled by a simple object in Frobenius algebra \mathcal{A} , and a red dot on a vertex to represent a coefficient f multiplied to this state.

$$\left| \begin{array}{c} \text{red dot} \\ \uparrow a_\alpha \\ \text{red dot} \\ \swarrow b_\beta \quad \searrow c_\gamma \end{array} \right\rangle := f_{a_\alpha b_\beta c_\gamma} \left| \begin{array}{c} \text{red dot} \\ \uparrow a_\alpha \\ \text{red dot} \\ \swarrow b_\beta \quad \searrow c_\gamma \end{array} \right\rangle .$$
(B.4)

We also use dashed red edges or tails to represent that we are summing over all states with labels on this edge in $L_{\mathcal{A}}$. The definition (B.2) of Frobenius algebra \mathcal{A} can then be illustrated graphically by the Pachner moves.

$$\mathcal{T} \sum_{\gamma=0}^{n_c} \left| \begin{array}{c} s_\sigma \quad r_\rho \\ \swarrow \quad \searrow \\ \text{red dot} \\ \uparrow c_\gamma \\ \text{red dot} \\ \swarrow \quad \searrow \\ a_\alpha \quad b_\beta \end{array} \right\rangle = \sum_{t_\tau \in L_{\mathcal{A}}} \left| \begin{array}{c} s_\sigma \quad r_\rho \\ \swarrow \quad \searrow \\ \text{red dot} \\ \uparrow t_\tau \\ \text{red dot} \\ \swarrow \quad \searrow \\ a_\alpha \quad b_\beta \end{array} \right\rangle =: \left| \begin{array}{c} s_\sigma \quad r_\rho \\ \swarrow \quad \searrow \\ \text{red dot} \\ \uparrow \text{dashed} \\ \text{red dot} \\ \swarrow \quad \searrow \\ a_\alpha \quad b_\beta \end{array} \right\rangle ,$$

$$\mathcal{T} \left| \begin{array}{c} \text{red dot} \\ \uparrow b_\beta \\ \text{red dot} \\ \times \\ \text{red dot} \\ \downarrow a_\alpha \end{array} \right\rangle = d_{\mathcal{A}} \delta_{ab} \delta_{\alpha\beta} \left| \begin{array}{c} \text{red dot} \\ \uparrow a_\alpha \\ \text{red dot} \end{array} \right\rangle .$$

B.2 Bimodules over a Frobenius Algebra

A bimodule M over a Frobenius algebra \mathcal{A} in a fusion category \mathcal{F} is characterized by a pair of functions (n^M, P_M) . The function $n^M : L_{\mathcal{F}} \rightarrow \mathbb{N}$ returns the *multiplicity* n_a^M of $a \in L_{\mathcal{F}}$ appearing in bimodule M , satisfying $n_a^M = n_{a^*}^M$. The simple objects of M are labeled by pairs a_i , where $a \in L_{\mathcal{F}}$ satisfies $n_a^M > 0$, and $i = 1, 2, \dots, n_a^M$ labels the multiplicity index. We denote the set of all simple objects in bimodule M as L_M .


The action of Frobenius algebra \mathcal{A} on bimodule M is characterized by function $P_M : L_{\mathcal{A}}^2 \times L_M \times L_{\mathcal{F}} \times L_M \rightarrow \mathbb{C}$, satisfying the following defining equations:

$$\begin{aligned} & \sum_{uv \in L_{\mathcal{F}}} \sum_{y_v \in L_M} [P_M]_{x_\chi u y_v}^{a_\alpha r_\rho} [P_M]_{y_v v z_\zeta}^{b_\beta s_\sigma} G_{urw}^{v^* b y} G_{axc}^{w^* b u} G_{wrt^*}^{sz^* v} \sqrt{d_u d_v d_w d_y d_c d_t} \\ &= \sum_{\gamma=1}^{n_c} \sum_{\tau=1}^{n_t} P_{x_\chi w z_\zeta}^{c_\gamma t_\tau} f_{a_\alpha c_\gamma^* b_\beta} f_{r_\rho s_\sigma t_\tau}, \end{aligned} \quad (\text{B.5})$$

$$[P_M]_{x_\chi y z_\zeta}^{00} = \delta_{xy} \delta_{yz} \delta_{\chi v} \delta_{v \zeta}, \quad [P_M]_{x_\chi y z_\zeta}^{a_\alpha b_\beta} = [P_M]_{z_\zeta^* y^* x_\chi^*}^{b_\beta a_\alpha}.$$


This definition aligns with the one in the main body, where a bimodule M is expressed as a vector space spanned by simple objects $M = \mathbb{C}[L_M]$. A pair of Frobenius algebra elements $(a_\alpha, b_\beta) \in \mathbb{C}[L_{\mathcal{A}}]^2$ is represented as a three-index tensor P_M on the bimodule space $\mathbb{C}[L_M]$.

For convenience, in a lattice model, we use a blue line to indicate that this line is labeled by a simple object in bimodule M and a wavy blue line to represent summing over all states with labels on this edge in $L_{\mathcal{F}}$ with coefficient P_M :



$$\left| \begin{array}{c} z_\zeta \\ \vdots \\ M \\ \vdots \\ x_\chi \end{array} \right. \begin{array}{c} \leftarrow b_\beta \\ \rightarrow a_\alpha \end{array} \right\rangle := \sum_{y \in L_{\mathcal{F}}} [P_M]_{x_\chi y z_\zeta}^{a_\alpha b_\beta} \left| \begin{array}{c} z_\zeta \\ \vdots \\ y \\ \vdots \\ x_\chi \end{array} \right. \begin{array}{c} \leftarrow b_\beta \\ \rightarrow a_\alpha \end{array} \right\rangle. \quad (\text{B.6})$$

The definition (B.5) of bimodule M can then be depicted graphically by Pachner moves:



$$\mathcal{T} \sum_{y_v \in L_M} \left| \begin{array}{c} z_\zeta \\ \vdots \\ M \\ \vdots \\ x_\chi \end{array} \right. \begin{array}{c} \leftarrow s_\sigma \\ \rightarrow b_\beta \\ \rightarrow y_v \\ \rightarrow r_\rho \\ \rightarrow a_\alpha \end{array} \right\rangle = \left| \begin{array}{c} z_\zeta \\ \vdots \\ M \\ \vdots \\ x_\chi \end{array} \right. \begin{array}{c} \leftarrow s_\sigma \\ \rightarrow r_\rho \\ \rightarrow a_\alpha \end{array} \right\rangle.$$

B.3 The Bimodule Fusion Category over a Frobenius Algebra

The set of all bimodules over a given Frobenius algebra \mathcal{A} in a fusion category \mathcal{F} forms a fusion category, denoted as $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$. In this appendix, we briefly introduce the categorical data of the fusion category $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$.

1. A bimodule M in $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ is *simple* if it cannot be written as a direct sum of two other bimodules. That is, we cannot find two bimodules M_1 and M_2 such that:

$$n_a^M = n_a^{M_1} + n_a^{M_2}, \quad [P_M]_{x_\chi y z \zeta}^{a_\alpha b_\beta} = \begin{cases} [P_{M_1}]_{x_\chi y z \zeta}^{a_\alpha b_\beta}, & (\chi \leq n_x^{M_1}, \zeta \leq n_z^{M_1}), \\ [P_{M_2}]_{x_{(\chi-n_x^{M_1})} y z_{(\zeta-n_z^{M_2})}^{a_\alpha b_\beta}, & (\chi > n_x^{M_1}, \zeta > n_z^{M_1}), \\ 0. & (\text{otherwise}). \end{cases}$$

2. The quantum dimension of a bimodule M in $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ is

$$d_M = \frac{1}{d_{\mathcal{A}}} \sum_{a \in L_{\mathcal{F}}} n_a^M d_a. \quad (\text{B.7})$$

3. For any three bimodules M_1 , M_2 , and M_3 , we can represent their fusion rules in terms of their simple objects. Define the matrix $[\Delta_{M_1 M_2 M_3}]$ that represents how the basis elements in the bimodule spaces are connected when the three bimodules fuse:

$$[\Delta_{M_1 M_2 M_3}]_{r_\rho s_\sigma t_\tau}^{x_\chi y_\nu z_\zeta} := \frac{1}{d_{\mathcal{A}}^3} \sum_{a_\alpha b_\beta c_\gamma \in L_{\mathcal{A}}} \sum_{p \in L_{\mathcal{F}}} \sum_{u_\rho \in L_{M_1}} \sum_{v_\sigma \in L_{M_2}} \sum_{w_\lambda \in L_{M_3}} [P_1]_{x_\chi u r_\rho}^{b_\beta c_\gamma^*} [P_2]_{y_\nu v s_\sigma}^{c_\gamma a_\alpha^*} \times [P_3]_{z_\zeta w t_\tau}^{a_\alpha b_\beta^*} G_{c^* r^* p}^{b x u^*} G_{b r^* t^*}^{s w p} G_{a w^* s^*}^{p v z} G_{v p c}^{x y z} \sqrt{d_u d_v d_w d_a d_b d_c d_r d_s d_t} d_p. \quad (\text{B.8})$$

This definition can be depicted graphically:

$$\mathcal{J} \quad \begin{array}{c} \text{Diagram of three bimodules } M_1, M_2, M_3 \text{ fusing at a central point. } \\ \text{Inputs: } x_\chi, y_\nu, z_\zeta. \text{ Outputs: } r_\rho, s_\sigma, t_\tau. \end{array} = d_{\mathcal{A}}^3 [\Delta_{M_1 M_2 M_3}]_{r_\rho s_\sigma t_\tau}^{x_\chi y_\nu z_\zeta} \quad \begin{array}{c} \text{Diagram of a single fusion vertex.} \\ \text{Input: } x_\chi. \text{ Outputs: } y_\nu, z_\zeta. \end{array} .$$

The fusion rule of three bimodules M_1, M_2, M_3 is

$$\delta_{M_1 M_2 M_3} = \text{Tr}[\Delta_{M_1 M_2 M_3}]. \quad (\text{B.9})$$

4. The Frobenius algebra \mathcal{A} itself is the trivial bimodule M_0 over \mathcal{A} :

$$L_{M_0} = L_{\mathcal{A}}, \quad [P_{M_0}]_{x_\chi y z \zeta}^{a_\alpha b_\beta} = \sum_{v=1}^{n_y} f_{a_\alpha x_\chi y_v^*} f_{y_v b_\beta z_\zeta^*}. \quad (\text{B.10})$$

Given a bimodule M , its opposite bimodule M^* is

$$L_{M^*} = L_M, \quad [P_{M^*}]_{x_\chi y z \zeta}^{a_\alpha b_\beta} = ([P_M]_{x_\chi y z \zeta}^{a_\alpha b_\beta})^*. \quad (\text{B.11})$$

5. The bimodule conditions induces that matrix $\Delta_{M_1 M_2 M_3}$ is a projector:

$$\Delta_{M_1 M_2 M_3}^2 = \Delta_{M_1 M_2 M_3}.$$

If $\delta_{M_1 M_2 M_3} \neq 0$, we can find the normalized eigenvectors $\mathcal{V}_{M_1 M_2 M_3}^{r_\rho s_\sigma t_\tau} \in \mathbb{C}$, such that

$$\begin{aligned} \sum_{r_\rho \in L_{M_1}} \sum_{s_\sigma \in L_{M_2}} \sum_{t_\tau \in L_{M_3}} [\Delta_{M_1 M_2 M_3}]_{r_\rho s_\sigma t_\tau}^{x_\chi y_\nu z_\zeta} \mathcal{V}_{M_1 M_2 M_3}^{r_\rho s_\sigma t_\tau} &= \mathcal{V}_{M_1 M_2 M_3}^{x_\chi y_\nu z_\zeta}, \\ \sum_{x_\chi \in L_{M_1}} \sum_{y_\nu \in L_{M_2}} \sum_{z_\zeta \in L_{M_3}} |\mathcal{V}_{M_1 M_2 M_3}^{x_\chi y_\nu z_\zeta}|^2 \sqrt{d_x d_y d_z} &= d_{\mathcal{A}}^2 \sqrt{d_{M_1} d_{M_2} d_{M_3}}. \end{aligned} \quad (\text{B.12})$$

For convenience, in a lattice model, we use blue lines labeled by a bimodule M to represent summing over all states with labels in L_M on this line. Additionally, blue dots represent a state multiplied by a coefficient \mathcal{V} :

$$\left| \begin{array}{c} \text{---} M_1 \text{---} \\ | \\ \text{---} M_2 \text{---} \bullet \text{---} M_3 \text{---} \end{array} \right\rangle := \sum_{x_\chi \in L_{M_1}} \sum_{y_\nu \in L_{M_2}} \sum_{z_\zeta \in L_{M_3}} \mathcal{V}_{M_1 M_2 M_3}^{x_\chi y_\nu z_\zeta} \left| \begin{array}{c} x_\chi \\ | \\ y_\nu \text{---} z_\zeta \end{array} \right\rangle. \quad (\text{B.13})$$

Such a state is invariant under $\mathcal{D}_{M_1 M_2 M_3}$ matrix. The $6j$ -symbol of $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ category is

$$G_{M_3 M_4 M'}^{M_1 M_2 M} = \frac{1}{d_{\mathcal{A}}^3 \sqrt{d_{M_1} d_{M_2} d_{M_3} d_{M_4} d_M d_{M'}}} \mathcal{T} \left(\begin{array}{c} \bullet \\ \text{---} M_1 \text{---} \bullet \\ | \\ \text{---} M_2 \text{---} \bullet \text{---} M_3 \text{---} \bullet \\ | \\ \text{---} M_4 \text{---} \bullet \text{---} M' \end{array} \right). \quad (\text{B.14})$$

C General Constructions of Dualities and Symmetry Transformations in the Extended String-Net Model

Given a fusion category \mathcal{F} and a Frobenius algebra $\mathcal{A} \in \mathcal{F}$, two string-net models with \mathcal{F} and $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ as the input data describe the same topological order. Categorically, $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ is defined by an injective functor

$$\mathcal{D} : \text{Bimod}_{\mathcal{F}}(\mathcal{A}) \rightarrow \mathcal{F}, \quad M \mapsto \bigoplus_{a \in L_{\mathcal{F}}} n_a^M a, \quad (\text{C.1})$$

and for any morphisms $\phi_{M_1 M_2}^{M_3} \in \text{Bimod}_{\mathcal{F}}(\mathcal{A}) : M_1 \otimes M_2 \rightarrow M_3$ and $\varphi_{xy}^z \in \mathcal{F} : x \otimes y \rightarrow z$,

$$\mathcal{D}(\phi_{M_1 M_2}^{M_3}) = \bigoplus_{z_\zeta \in L_{M_3}} \left[\bigoplus_{x_\chi \in L_{M_1}} \bigoplus_{y_\nu \in L_{M_2}} \mathcal{V}_{M_1 M_2 M_3}^{x_\chi y_\nu z_\zeta} \varphi_{x_\chi y_\nu}^{z_\zeta} \right],$$

where x_χ , y_ν , and z_ζ are respectively the χ -th x object, ν -th y , and ζ -th z in the direct sum $\mathcal{D}(M)$.

Such a functor \mathcal{D} induces a duality between the two models with \mathcal{F} and $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ as the input data:

$$\left| M \right\rangle \Longrightarrow \sum_{a_\alpha, b_\beta \in L_{\mathcal{A}}} \sum_{x_\chi, z_\zeta \in L_M} \begin{array}{c} z_\zeta \\ \uparrow \\ \text{---} M \text{---} \\ \downarrow \\ x_\chi \end{array} \quad . \quad (\text{C.2})$$

This duality induces a unitary morphism between the Hilbert spaces $\mathcal{H}_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})}$ and $\mathcal{H}_{\mathcal{F}}$ of these two models. Such a unitary linear transformation can be understood plaquette by plaquette:

$$\begin{array}{c} \begin{array}{c} N_2 \\ \downarrow \\ N_1 \quad M_1 \quad M_2 \quad N_3 \\ \downarrow \quad \downarrow \quad \downarrow \\ M_0 \quad M \quad M_3 \\ \downarrow \quad \downarrow \quad \downarrow \\ M_6 \quad M_4 \quad M_5 \\ \downarrow \quad \downarrow \quad \downarrow \\ N_6 \quad N_5 \end{array} \end{array} \Longrightarrow \frac{1}{d_{\mathcal{A}}^g} \sum_{x_i, y_i \in L_{M_i}} \sum_{e_i \in L_{N_i}} \sum_{p_\alpha, q_\beta \in L_M} \begin{array}{c} N_2 \\ \downarrow \\ N_1 \quad M_1 \quad M_2 \quad N_3 \\ \downarrow \quad \downarrow \quad \downarrow \\ M_0 \quad M \quad M_3 \\ \downarrow \quad \downarrow \quad \downarrow \\ M_6 \quad M_4 \quad M_5 \\ \downarrow \quad \downarrow \quad \downarrow \\ N_6 \quad N_5 \end{array} \quad . \quad (\text{C.3})$$

Note that the black edges and tails labeled by $M_i, N_i \in \text{Bimod}_{\mathcal{F}}(\mathcal{A})$ represent basis states in the dual model, where $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ is the input fusion category and M_i, N_i are simple objects. In contrast, the blue edges and tails labeled by $M_i, N_i \in \text{Bimod}_{\mathcal{F}}(\mathcal{A})$ represent superposition states in the original model with \mathcal{F} as the input fusion category, where the superpositions are defined in Eqs. (B.6) and (B.13).

This understanding differs slightly from the unitary duality transformation in the sense that, after the above transformation (C.3) is applied to all plaquettes, the resulting basis state $|\psi\rangle$ satisfies

$$\langle \psi | \psi \rangle = d_{\mathcal{A}}^{g-2}, \quad (\text{C.4})$$

where g is the genus of the surface on which the lattice is embedded. By applying the duality map and normalizing the resulting basis states, we obtain a unitary morphism between the two string-net models.

After the topological moves in Eq. (C.3), the degree of freedom on any edge will cease to have any multiplicity index of simple objects in bimodules, while that on any tail will still have a multiplicity index. Therefore, to make sense of this duality and make it unitary, we are urged to enlarge the Hilbert space of the original Fibonacci string-net model on each tail but not on the edges, such that two simple objects $a_\alpha, a_\beta \in L_M$ with different multiplicity indices $\alpha \neq \beta$ are distinguishable on tails.

C.1 Enlarging the Hilbert Space

In the enlarged Hilbert space, each tail carries a degree of freedom labeled by a pair a_α , where

$$a \in L_{\mathcal{F}}, \quad \alpha = 1, 2, \dots, N_a^A, \quad N_a^A = \max_{M \in L_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})}} \{n_a^M\}, \quad (\text{C.5})$$

where $L_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})}$ is the set of all simple bimodules over Frobenius algebra \mathcal{A} . But the basic degrees of freedom on edges remain to take value varying the simple objects of the input fusion category \mathcal{F} . The Hilbert space on the tail is spanned by all enlarged degrees of freedom on tails and original degrees of freedom on edges, subject to the fusion rules on all vertices.

For any bimodule M , its simple object $x_\chi^M \in L_M$ corresponds to a superposition state $|x_\chi^M\rangle$ in the local Hilbert space of a tail:

$$|x_\chi^M\rangle := \sum_{i=1}^{N_x^A} A_{\chi,i}^{x,M} |x_i\rangle. \quad (\text{C.6})$$

All different states should satisfy the orthonormal conditions:

$$\sum_{\alpha=1}^{n_x^M} \sum_{\beta=1}^{n_x^N} M_1 \begin{array}{c} \bullet \\ \text{---} y_\beta^N \text{---} \\ \times \\ \text{---} x_\alpha^M \text{---} \\ \bullet \end{array} M_2 = d_A n_x^M \delta_{xy} \delta_{M_1 M_2 M_3} \delta_{MN} \sqrt{d_{M_1} d_{M_2} d_x}. \quad (\text{C.7})$$

C.2 Duality

The duality map (C.3) can be simplified. We can represent the unitary duality map vertex by vertex:

$$\mathcal{D} := \frac{1}{d_A^{N_P - 1 + \frac{g}{2}}} \prod_{\text{Edge } e} E_e \prod_{\text{Vertex } v} \mathcal{D}_v, \quad (\text{C.8})$$

where N_P is the number of plaquettes in the lattice, and \mathcal{D}_v acts on vertex v :

$$\mathcal{D}_v \begin{array}{c} \downarrow M_1 \\ \nearrow M_2 \quad \searrow M_3 \end{array} := \begin{array}{c} \downarrow M_1 \\ \bullet \\ \nearrow M_2 \quad \searrow M_3 \end{array} = \sum_{x_\chi \in L_{M_1}} \sum_{y_\nu \in L_{M_2}} \sum_{z_\zeta \in L_{M_3}} \mathcal{V}_{M_1 M_2 M_3}^{x_\chi y_\nu z_\zeta} \begin{array}{c} \downarrow x_\chi \\ \nearrow y_\nu \quad \searrow z_\zeta \end{array}. \quad (\text{C.9})$$

Note that each edge connects two vertices that are acted upon by two \mathcal{D}_v operators independently. Nevertheless, an edge e can only carry one label. We use E_e to ensure this

uniqueness:

$$E_e \quad \begin{array}{c} \diagup \quad \diagdown \\ \text{---} x_\alpha^{M_1} \text{---} y_\beta^{M_2} \text{---} \\ \diagdown \quad \diagup \\ v_1 \qquad v_2 \end{array} \quad := \quad \sum_i (A_{\alpha,i}^{x,M_1})^* A_{\beta,i}^{x,M_2} \quad \begin{array}{c} \diagup \quad \diagdown \\ \text{---} x \text{---} \\ \diagdown \quad \diagup \end{array} . \quad (\text{C.10})$$

The E_e moves erase the multiplicity indices of labels on edges. But the multiplicity labels on tails are retained.

The Hilbert space is not preserved under the duality map:

$$\mathcal{H}_{\mathcal{F}} \neq \mathcal{DH}_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})},$$

where $\mathcal{H}_{\mathcal{F}}$ and $\mathcal{H}_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})}$ are the physical Hilbert spaces of the string-net model with input fusion category \mathcal{F} and $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$, respectively, considered as subspaces of the enlarged Hilbert space. Nevertheless, since the two models describe the same topological order, the ground-state subspace \mathcal{H}_0 is preserved under the duality map:

$$\mathcal{H}_{0,\mathcal{F}} = \mathcal{H}_{0,\text{Bimod}_{\mathcal{F}}(\mathcal{A})}, \quad (\text{C.11})$$

where $\mathcal{H}_{0,\mathcal{F}}$ and $\mathcal{H}_{0,\text{Bimod}_{\mathcal{F}}(\mathcal{A})}$ are the ground-state subspaces of the string-net model with input fusion category \mathcal{F} and $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$, respectively.

C.3 Symmetry Transformation

In particular cases, \mathcal{F} and $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$ are isomorphic fusion category. That is, there exists an isomorphic functor $\varphi_{\mathcal{A}}$ that maps simple objects of \mathcal{F} to simple objects in $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$:

$$\varphi_{\mathcal{A}}(a) = M_a \in \text{Bimod}_{\mathcal{F}}(\mathcal{A}). \quad (\text{C.12})$$

Such isomorphic functor induces a linear isomorphism $\varphi_{\mathcal{A}}$ between the Hilbert space of these two models that maps basic degrees of freedom on edges and tails to basic degrees of freedom:

$$\varphi_{\mathcal{A}} : \mathcal{H}_{\mathcal{F}} \rightarrow \mathcal{H}_{\text{Bimod}_{\mathcal{F}}(\mathcal{A})}, \quad \varphi_{\mathcal{A}} \begin{array}{c} | \\ a \\ | \end{array} := \begin{array}{c} | \\ M_a \\ | \end{array} . \quad (\text{C.13})$$

The composition

$$\mathcal{G} := \mathcal{D} \circ i : \mathcal{H}_{\mathcal{F}} \rightarrow \mathcal{H}_{\mathcal{F}} \quad (\text{C.14})$$

is just a unitary transformation of the same model with \mathcal{F} as the input fusion category, and the symmetry transformation is the composition of the unitary transformation and projection back into the original degrees of freedom. The set of all symmetry transformations in the Hilbert space of the string-net model with \mathcal{F} as the input fusion category forms the symmetry of the topological order.

In particular, consider the trivial Frobenius algebra \mathcal{A}_0 :

$$L_{\mathcal{A}_0} = \{0\}, \quad f_{000} = 1, \quad (\text{C.15})$$

whose simple objects are classified by simple objects in $L_{\mathcal{F}}$:

$$L_{M_a} = \{a\}, \quad P_{aaa}^{00} = 1. \quad (\text{C.16})$$

The gauge transformation induced by Frobenius algebra \mathcal{A}_0 is the identity transformation of the string-net model.

C.4 Braiding of Bimodules

The input fusion category \mathcal{F} is a fusion category that lacks a braiding structure for exchanging two simple objects a and b . Nevertheless, the braiding of the trivial object 0 with any other simple object a always exists as the trivial braiding, which can be represented graphically as:

The last equality holds because the fusion of 0 with any simple object a is also trivial.

Similarly, in the bimodule category $\text{Bimod}_{\mathcal{F}}(\mathcal{A})$, the trivial bimodule \mathcal{A} braids trivially with any other bimodule $M \in \text{Bimod}_{\mathcal{F}}(\mathcal{A})$, based on the definition of bimodules:

As a practical example, consider the situation where we contrast a plaquette with a tail labeled by the trivial bimodule \mathcal{A} in the original model. Since the Pachner moves (A.3) can only contrast plaquettes without nontrivial tails inside them, one must first “pull” the trivial tail out of the plaquette and then annihilate the plaquettes, as shown in Fig. 3.

D The \mathbb{Z}_2 Global Symmetry transformation of the \mathbb{Z}_2 Toric Code String-Net Model

To provide another example of our symmetry transformation construction, and to contrast with the Fibonacci categorical gauge symmetry discussed in the main text, we briefly apply our method in this appendix to construct a well-known symmetry transformation: the \mathbb{Z}_2 global symmetry transformation of the \mathbb{Z}_2 toric code string-net model[6]. In the toric code case, we do not need to enlarge the Hilbert space. This transformation exchanges the charge and flux excitations.

D.1 The toric code string-net model

The input fusion category of the toric code string-net model is the \mathbb{Z}_2 fusion category, which uses the two group elements $\pm 1 \in \mathbb{Z}_2$ as its simple objects. The fusion rules capture




Figure 3: Contrast a plaquette with a tail labeled by the trivial bimodule \mathcal{A} in the original model.

the group multiplicity rules of \mathbb{Z}_2 :

$$\delta_{ijk} = \frac{ijk + 1}{2},$$

and the quantum dimensions are $d_{\pm} = 1$. The nonzero $6j$ symbols are given by

$$G_{klm}^{ijn} = \delta_{ijm} \delta_{klm} \delta_{iln} \delta_{jkn}.$$

In the toric code string-net model, each edge or tail on the lattice carries a group element $\pm 1 \in \mathbb{Z}_2$, and the degrees of freedom i, j, k on any three edges or tails meeting at a vertex must satisfy the fusion rule $\delta_{ijk} = 1$. The Hamiltonian of the toric code string-net model is a sum of commutative projectors A_P and B_P ,

$$H_{\text{TC}} := - \sum_{\text{Plaquettes } P} (A_P + B_P), \quad (\text{D.1})$$

where A_P acts on tails in plaquettes P , and B_P acts on edges surrounding plaquettes P :

$$A_P := \frac{p+1}{2} \left(\begin{array}{c} e_2 \\ i_1 \quad i_2 \\ e_1 \quad i_0 \quad e_3 \\ \text{---} p \text{---} \\ i_6 \quad i_3 \\ e_6 \quad i_5 \quad e_4 \\ e_5 \end{array} \right) := \frac{p+1}{2} \left(\begin{array}{c} e_2 \\ i_1 \quad i_2 \\ e_1 \quad i_0 \quad e_3 \\ \text{---} p \text{---} \\ i_6 \quad i_3 \\ e_6 \quad i_5 \quad e_4 \\ e_5 \end{array} \right), \quad (\text{D.2})$$

$$B_P := \frac{1}{2} \left(\begin{array}{c} e_2 \\ i_1 \quad i_2 \\ e_1 \quad i_0 \quad e_3 \\ \text{---} p \text{---} \\ i_6 \quad i_3 \\ e_6 \quad i_5 \quad e_4 \\ e_5 \end{array} \right) + \frac{1}{2} \left(\begin{array}{c} e_2 \\ -i_1 \quad -i_2 \\ e_1 \quad -i_0 \quad e_3 \\ \text{---} p \text{---} \\ -i_6 \quad -i_3 \\ e_6 \quad -i_5 \quad e_4 \\ e_5 \end{array} \right). \quad (\text{D.3})$$

The ground states are common eigenstates of all A_P and B_P operators with $+1$ eigenvalues, while an excited state $|\psi\rangle$ is another common eigenstate satisfying $A_P|\psi\rangle = 0$ (or $B_P|\psi\rangle = 0$).

0) for one or more plaquettes P , in each of which there resides a chargeon e (or a fluxon m). Unlike the original version of the string-net model, where chargeons reside on vertices, both chargeons and fluxons are situated in the plaquettes of our enlarged model. If $A_P|\psi\rangle = B_P|\psi\rangle = 0$ in plaquette P , there is a dyon ϵ in plaquette P . We also refer to the ground state as the trivial excited state, in which there are trivial anyons 1 in all plaquettes.

D.2 The global symmetry transformation of the toric code topological order

The \mathbb{Z}_2 fusion category has a unique nontrivial Frobenius algebra \mathcal{A} , such that

$$L_{\mathcal{A}} = \{1, -1\}, \quad f_{ijk} = \delta_{ijk}.$$

There are two simple bimodules over \mathcal{A} , denoted as M_{\pm} , such that

$$\begin{aligned} L_{M_+} = L_{M_-} &= \{0, 1\}, & [P_+]_{xyz}^{ab} &= \delta_{axy}\delta_{byz}, \\ [P_-]_{++++}^{++} = [P_-]_{----}^{++} &= 1, & [P_-]_{+-+}^{--} = [P_-]_{-+-}^{--} &= -1, \\ [P_-]_{++-}^{+-} = [P_-]_{-+-}^{+-} &= i, & [P_-]_{--+}^{+-} = [P_-]_{+-}^{+-} &= -i. \end{aligned}$$

The nonzero vertex coefficients are

$$\mathcal{V}_{M_+M_+M_+}^{ijk} = \delta_{ijk}, \quad \mathcal{V}_{M_+M_-M_-}^{+++} = \mathcal{V}_{M_+M_-M_-}^{---} = 1, \quad \mathcal{V}_{M_+M_-M_-}^{+-+} = i, \quad \mathcal{V}_{M_+M_-M_-}^{+--} = -i.$$

The fusion category $\text{Bimod}_{\mathbb{Z}_2}(\mathcal{A})$ is isomorphic to the \mathbb{Z}_2 fusion category:

$$\delta_{M_iM_jM_k} = \delta_{ijk} = 1, \quad d_{M_i} = d_i, \quad G_{M_kM_lM_n}^{M_iM_jM_m} = G_{kln}^{ijm}.$$

By applying the above categorical data, we can formulate a symmetry transformation \mathcal{G} is

$$\mathcal{G} := \frac{1}{2^{N_P - 1 + \frac{g}{2}}} \prod_{\text{Vertices } v} \mathcal{G}_v, \quad (\text{D.4})$$

where N_P is the number of plaquettes in the lattice, and g is the genus number of the surface in which the lattice is embedded. Each local transformation \mathcal{G}_v acts on vertex v of the lattice as

$$\begin{aligned} \mathcal{G}_v \begin{array}{c} + \quad + \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} &:= \begin{array}{c} + \quad + \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} + \begin{array}{c} - \quad - \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} + \begin{array}{c} + \quad - \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ - \end{array} + \begin{array}{c} - \quad + \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ - \end{array}, \\ \mathcal{G}_v \begin{array}{c} - \quad - \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} &:= \begin{array}{c} + \quad + \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} + \begin{array}{c} - \quad - \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ + \end{array} + i \begin{array}{c} + \quad - \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ - \end{array} - i \begin{array}{c} - \quad + \\ \diagdown \quad \diagup \\ v \\ \diagup \quad \diagdown \\ - \end{array}, \end{aligned} \quad (\text{D.5})$$

where $+$ ($-$) refers to 1 (-1), and the three lines meeting at vertex v can be either edges or tails.

The symmetry transformation \mathcal{G} is a \mathbb{Z}_2 global symmetry transformation of the toric code string-net model because:

1. \mathcal{G} is a unitary \mathbb{Z}_2 transformation:

$$\mathcal{G}^\dagger = \mathcal{G}^{-1} = \mathcal{G}. \quad (\text{D.6})$$

2. \mathcal{G} preserves the model's Hamiltonian H_{TC} :

$$\mathcal{G}^\dagger H_{\text{TC}} \mathcal{G} = H_{\text{TC}}. \quad (\text{D.7})$$

3. Transformation \mathcal{G} preserves the ground-state Hilber space of the model but exchanges A_P and B_P operators:

$$\mathcal{G}^\dagger A_P \mathcal{G} = B_P, \quad \mathcal{G}^\dagger B_P \mathcal{G} = A_P, \quad (\text{D.8})$$

and hence exchanges chargeons and fluxons because an A_P (B_P) measures the chargeon (fluxon) in plaquette P ⁸.

Note that the global symmetry transformation \mathcal{G} maps the degrees of freedom 0 and 1—the simple objects in the \mathbb{Z}_2 fusion category—to the degrees of freedom M_0 and M_1 , which are the simple objects in the bimodule category $\text{Bimod}_{\mathbb{Z}_2}(\mathcal{A})$. Analogous to our discussion about the Fibonacci categorical gauge symmetry, this EM-exchange global symmetry of the toric code topological order is fundamentally a *categorical symmetry* described by a fusion 2-category—the \mathbb{Z}_2 fusion 2-category defined based on the \mathbb{Z}_2 fusion category. It reduces to a \mathbb{Z}_2 group symmetry because the transformation is invertible.

D.3 Criterion for Distinguishing Gauge Symmetries from Global Symmetries

The symmetry transformation (D.4) of the \mathbb{Z}_2 toric code topological order is a global symmetry transformation that exchanges the anyon species of chargeons and fluxons. In contrast, the symmetry transformation (5.2) of the doubled Fibonacci topological order in the main text is a gauge symmetry that preserves all anyon species but only transforms excited states within the internal Hilbert space of each anyon. To determine whether a symmetry transformation defined by a Frobenius algebra \mathcal{A} in a fusion category \mathcal{F} is a global symmetry or a gauge symmetry, we use the criterion based on Morita equivalence between Frobenius algebras[88]. This concept of Morita equivalence between two Frobenius algebras differs from the concept of Morita equivalence between fusion categories introduced in Appendix. B.

Algebraic Morita equivalence implies that the two Frobenius algebras have the same modules. According to the boundary-bulk correspondence, two string-net models with bimodule categories over two Morita-equivalent Frobenius algebras not only have the same anyon species but also exhibit the same relationships between each anyon species and the fundamental degrees of freedom of the models. As previously noted, the fusion category \mathcal{F} itself is the bimodule category over the trivial one-dimensional Frobenius algebra $\mathcal{A}_0 = \{1\}$.

Therefore, a symmetry transformation is a gauge symmetry if and only if its defining Frobenius algebra \mathcal{A} is algebraically Morita equivalent to the trivial Frobenius algebra \mathcal{A}_0 . This is the case for the Frobenius algebra $\mathcal{A} = \{1, \tau\}$ in the Fibonacci fusion category. Otherwise, it is a global symmetry, as in the case of \mathbb{Z}_2 toric code.

⁸In the original string-net model, chargeons reside on vertices and fluxons are located in plaquettes, necessitating lattice dualization after the symmetry transformation. In contrast, our model places both chargeons and fluxons in plaquettes, eliminating the need to alter the lattice shape.

E Frobenius Algebra of Fibonacci Fusion Categories and Its Simple Bimodules

In this appendix, we list the categorical data of the Fibonacci fusion category. The Fibonacci fusion category has two simple objects, denoted as 1 and τ . The nonzero fusion rules are $\delta_{111} = \delta_{1\tau\tau} = \delta_{\tau\tau\tau} = 1$, and the quantum dimensions are

$$d_1 = 1, \quad d_\tau = \phi = \frac{\sqrt{5} + 1}{2}.$$

The nonzero $6j$ symbols are

$$G_{111}^{111} = 1, \quad G_{\tau\tau\tau}^{111} = \frac{1}{\sqrt{\phi}}, \quad G_{1\tau\tau}^{1\tau\tau} = G_{\tau\tau\tau}^{1\tau\tau} = \frac{1}{\phi}, \quad G_{\tau\tau\tau}^{\tau\tau\tau} = -\frac{1}{\phi^2}.$$

The Fibonacci fusion category has a nontrivial Frobenius algebra \mathcal{A} , such that

$$L_{\mathcal{A}} = \{1, \tau\}, \quad f_{111} = f_{1\tau\tau} = f_{\tau 1\tau} = f_{\tau\tau 1} = 1, \quad f_{\tau\tau\tau} = -\frac{1}{\phi^{\frac{3}{4}}}.$$

There are two simple bimodules over \mathcal{A} , denoted as M_0 and M_1 , such that

$$L_{M_0} = \{1, \tau\},$$

$$\begin{aligned} [P_{M_0}]_{1111}^{11} &= [P_{M_0}]_{\tau\tau\tau\tau}^{11} = 1, & [P_{M_0}]_{111\tau}^{1\tau} &= [P_{M_0}]_{\tau\tau\tau 1}^{1\tau} = 1, & [P_{M_0}]_{\tau\tau\tau\tau}^{1\tau} &= \frac{1}{\phi^{\frac{3}{4}}}, \\ [P_{M_0}]_{\tau 111}^{\tau 1} &= [P_{M_0}]_{1\tau\tau\tau}^{\tau 1} = 1, & [P_{M_0}]_{\tau\tau\tau\tau}^{\tau 1} &= \frac{1}{\phi^{\frac{3}{4}}}, \\ [P_{M_0}]_{1\tau 11}^{\tau\tau} &= [P_{M_0}]_{\tau 1\tau\tau}^{\tau\tau} = 1, & [P_{M_0}]_{1\tau\tau\tau}^{\tau\tau} &= [P_{M_0}]_{\tau\tau\tau 1}^{\tau\tau} = \frac{1}{\phi^{\frac{3}{4}}}, & [P_{M_0}]_{\tau\tau\tau\tau}^{\tau\tau} &= \frac{1}{\phi^{\frac{3}{2}}}. \end{aligned}$$

$$L_{M_1} = \{1, \tau_1, \tau_2\},$$

$$[P_{M_1}]_{1111}^{11} = [P_{M_1}]_{\tau_0\tau_0\tau_0}^{11} = [P_{M_1}]_{\tau_1\tau_1\tau_1}^{11} = 1,$$

$$[P_{M_1}]_{111\tau_0}^{1\tau} = [P_{M_1}]_{\tau_1\tau_1\tau_1}^{1\tau} = [P_{M_1}]_{\tau_0\tau_0\tau_0}^{\tau 1} = [P_{M_1}]_{1\tau_1\tau_1}^{\tau 1} = \frac{1}{2\phi} + \frac{\sqrt{\phi}}{2} \mathfrak{i},$$

$$[P_{M_1}]_{111\tau_1}^{1\tau} = [P_{M_1}]_{\tau_0\tau_0\tau_0}^{1\tau} = [P_{M_1}]_{\tau_1\tau_1\tau_1}^{\tau 1} = [P_{M_1}]_{1\tau_0\tau_0}^{\tau 1} = \frac{1}{2\phi} - \frac{\sqrt{\phi}}{2} \mathfrak{i},$$

$$[P_{M_1}]_{\tau_0\tau_0\tau_0}^{1\tau} = [P_{M_1}]_{\tau_1\tau_1\tau_1}^{1\tau} = [P_{M_1}]_{\tau_0\tau_0\tau_0}^{\tau 1} = [P_{M_1}]_{\tau_1\tau_1\tau_1}^{\tau 1} = -\frac{\sqrt[4]{\phi}}{2\phi^2},$$

$$[P_{M_1}]_{\tau_0\tau_0\tau_1}^{1\tau} = [P_{M_1}]_{\tau_1\tau_1\tau_0}^{\tau 1} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2} \mathfrak{i}, \quad [P_{M_1}]_{\tau_1\tau_0\tau_0}^{1\tau} = [P_{M_1}]_{\tau_0\tau_0\tau_1}^{\tau 1} = -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2} \mathfrak{i},$$

$$[P_{M_1}]_{1\tau_1}^{\tau\tau} = -\frac{1}{\phi}, \quad [P_{M_1}]_{1\tau\tau_0}^{\tau\tau} = [P_{M_1}]_{1\tau\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_0\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_1\tau_1}^{\tau\tau} = -\frac{\sqrt[4]{\phi}}{\phi},$$

$$[P_{M_1}]_{\tau_0 1\tau_0}^{\tau\tau} = -\frac{1}{2\phi} + \frac{\mathfrak{i}}{2\sqrt{\phi}}, \quad [P_{M_1}]_{\tau_1 1\tau_1}^{\tau\tau} = -\frac{1}{2\phi} - \frac{\mathfrak{i}}{2\sqrt{\phi}}, \quad [P_{M_1}]_{\tau_0 1\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_1 1\tau_0}^{\tau\tau} = \frac{1}{2},$$

$$[P_{M_1}]_{\tau_0\tau\tau_1}^{\tau\tau} = [P_{M_1}]_{\tau_1\tau\tau_0}^{\tau\tau} = \frac{\sqrt{\phi}}{2\phi^2}, \quad [P_{M_1}]_{\tau_0\tau\tau_0}^{\tau\tau} = -\frac{\sqrt{\phi}}{2\phi^3} - \frac{\phi}{2}\mathfrak{i}, \quad [P_{M_1}]_{\tau_1\tau\tau_1}^{\tau\tau} = -\frac{\sqrt{\phi}}{2\phi^3} + \frac{\phi}{2}\mathfrak{i}.$$

The nonzero vertex coefficients are

$$\mathcal{V}_{M_0 M_0 M_0}^{111} = \mathcal{V}_{M_0 M_0 M_0}^{1\tau\tau} = \mathcal{V}_{M_0 M_0 M_0}^{\tau 1\tau} = \mathcal{V}_{M_0 M_0 M_0}^{\tau\tau 1} = 1, \quad \mathcal{V}_{M_0 M_0 M_0}^{\tau\tau\tau} = \frac{1}{\phi^{\frac{3}{4}}},$$

$$\mathcal{V}_{M_0 M_1 M_1}^{111} = \mathcal{V}_{M_0 M_1 M_1}^{1\tau_1\tau_1} = \mathcal{V}_{M_0 M_1 M_1}^{1\tau_2\tau_2} = 1, \quad \mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_1\tau_1} = \mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_2\tau_2} = -\frac{1}{2\phi^{\frac{3}{4}}},$$

$$\mathcal{V}_{M_0 M_1 M_1}^{\tau 1\tau_1} = \mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_2 1} = \frac{1}{2\phi} - \frac{\sqrt{\phi}}{2}i, \quad \mathcal{V}_{M_0 M_1 M_1}^{\tau 1\tau_2} = \mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_1 1} = \frac{1}{2\phi} + \frac{\sqrt{\phi}}{2}i,$$

$$\mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_1\tau_2} = -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2}i, \quad \mathcal{V}_{M_0 M_1 M_1}^{\tau\tau_2\tau_1} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2}i,$$

$$\mathcal{V}_{M_1 M_1 M_1}^{111} = \frac{1}{\phi^{\frac{3}{4}}}, \quad \mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_1\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_2\tau_2} = -\frac{1}{2}\sqrt{\frac{5}{\phi}},$$

$$\mathcal{V}_{M_1 M_1 M_1}^{1\tau_1\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1 1\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_1 1} = \mathcal{V}_{M_1 M_1 M_1}^{1\tau_2\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2 1\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_2 1} = -\frac{1}{2\phi^{\frac{3}{4}}},$$

$$\mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_1\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_2\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_1\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_2\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_1\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_2\tau_1} = \frac{1}{2\sqrt{t}\phi},$$

$$\mathcal{V}_{M_1 M_1 M_1}^{1\tau_1\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1 1\tau_2} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1\tau_2 1} = -\frac{\sqrt[4]{\phi}}{2} - \frac{\phi^{\frac{3}{4}}}{2}i,$$

$$\mathcal{V}_{M_1 M_1 M_1}^{1\tau_2\tau_1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_2\tau_1 1} = \mathcal{V}_{M_1 M_1 M_1}^{\tau_1 1\tau_2} = -\frac{\sqrt[4]{\phi}}{2} + \frac{\phi^{\frac{3}{4}}}{2}i.$$

The fusion category $\text{Bimod}_{\text{Fibo}}(\mathcal{A})$ is isomorphic to the Fibonacci fusion category by replacing 1 as M_0 , and τ as M_1 :

$$\delta_{M_0 M_0 M_0} = \delta_{M_0 M_1 M_1} = \delta_{M_1 M_1 M_1} = 1, \quad d_{M_0} = 1, \quad d_{M_1} = \phi,$$

$$G_{M_0 M_0 M_0}^{M_0 M_0 M_0} = 1, \quad G_{M_1 M_1 M_1}^{M_0 M_0 M_0} = \frac{1}{\sqrt{\phi}}, \quad G_{M_0 M_1 M_1}^{M_0 M_1 M_1} = G_{M_1 M_1 M_1}^{M_0 M_1 M_1} = \frac{1}{\phi}, \quad G_{M_1 M_1 M_1}^{M_1 M_1 M_1} = -\frac{1}{\phi^2}.$$

Bimodule M_1 has multiplicity $n_\tau^{M_1} = 2$, so we have to enlarge the Hilbert spaces on tails from 2-dimensional local Hilbert spaces spanned by 1 and τ to 3-dimensional local

Hilbert spaces spanned by 1 , τ_1 and τ_2 . To ensure the orthonormality, the basic state $|\tau\rangle$ labeled by the simple object τ in the trivial bimodule M_0 is a superposition state of two basic state $|\tau_1\rangle, |\tau_2\rangle$ labeled by the simple object τ in the nontrivial bimodule M_1 :

$$\begin{array}{c} \text{wavy line} \\ | \\ 1 \\ | \\ \text{wavy line} \end{array} \implies \begin{array}{c} \text{wavy line} \\ | \\ 1 \\ | \\ \text{wavy line} \end{array}, \quad \begin{array}{c} \text{wavy line} \\ | \\ \tau \\ | \\ \text{wavy line} \end{array} \implies \left(\frac{1}{2\phi} + \frac{\sqrt{\phi}}{2} \right) \begin{array}{c} \text{wavy line} \\ | \\ \tau_1 \\ | \\ \text{wavy line} \end{array} + \left(\frac{1}{2\phi} - \frac{\sqrt{\phi}}{2} \right) \begin{array}{c} \text{wavy line} \\ | \\ \tau_2 \\ | \\ \text{wavy line} \end{array}$$

This τ charge on a tail is unique up to exchanging τ_1 and τ_2 labels.

References

- [1] E. Dijkgraaf, Robbert and Witten, [Communications in Mathematical Physics](#) **429**, 393 (1990).
- [2] J. Fuchs, C. Schweigert, and A. Valentino, [Communications in Mathematical Physics](#) **1015**, 981 (2014), [arXiv:1307.3632](#).
- [3] A. Kapustin and N. Seiberg, [Journal of High Energy Physics](#) **2014**, 1 (2014).
- [4] I. Cong, M. Cheng, and Z. Wang, [Physical Review B](#) **96** (2017), 10.1103/PhysRevB.96.195129, [arXiv:1703.03564](#).
- [5] S. B. Bravyi and A. Y. Kitaev, [arXiv preprint quant-ph/9811052](#) (1998).
- [6] A. Kitaev, [Annals of Physics](#) **303**, 2 (2003).
- [7] S. Beigi, P. W. Shor, and D. Whalen, [Communications in Mathematical Physics](#) **306**, 663 (2011).
- [8] Y. Hu, Y. Wan, and Y.-S. Wu, [Physical Review B](#) **87**, 125114 (2013), [arXiv:1211.3695](#).
- [9] A. Bullivant, Y. Hu, and Y. Wan, [Physical Review B](#) **96**, 165138 (2017), [arXiv:1706.03611](#).
- [10] I. Cong, M. Cheng, and Z. Wang, [Communications in Mathematical Physics](#) 2017 355:2 **355**, 645 (2017).
- [11] H. Wang, Y. Li, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2020**, 30 (2020).
- [12] H. Wang, Y. Li, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2020**, 30 (2020), [arXiv:1910.13441](#).
- [13] Y. Hu and Y. Wan, [Journal of High Energy Physics](#) **2020**, 170 (2020).
- [14] V. G. Turaev and O. Y. Viro, [Topology](#) **31**, 865 (1992).
- [15] V. G. Turaev, *Quantum Invariants of Knots and 3-manifolds* (Walter de Gruyter, Berlin, 1994).
- [16] L. Bhardwaj, D. Gaiotto, and A. Kapustin, [Journal of High Energy Physics](#) **2017**, 1 (2017).
- [17] X. Cui, *Higher categories and topological quantum field theories* (University of California, Santa Barbara, 2016).
- [18] M. Levin and X.-g. Wen, [Physical Review B](#) **71**, 21 (2005), [arXiv:0404617 \[cond-mat\]](#).
- [19] L.-Y. Hung and Y. Wan, [Physical Review B](#) **86**, 235132 (2012), [arXiv:1207.6169](#).
- [20] Y. Hu, S. D. Stirling, and Y.-s. Wu, [Physical Review B](#) **85**, 075107 (2011), [arXiv:1105.5771](#).
- [21] A. Kitaev and L. Kong, [Communications in Mathematical Physics](#) **313**, 351 (2012).

- [22] M. D. Schulz, S. Dusuel, K. P. Schmidt, and J. Vidal, [Physical review letters](#) **110**, 147203 (2013).
- [23] T. Lan and X. G. Wen, [Physical Review B - Condensed Matter and Materials Physics](#) **90**, 115119 (2014), [arXiv:1311.1784](#).
- [24] C.-H. Lin and M. Levin, [Physical Review B](#) **89**, 195130 (2014).
- [25] Y. Hu, Y. Wan, and Y.-s. Wu, [Chinese Physics Letters](#) **34**, 077103 (2017), [arXiv:1706.00650](#).
- [26] Y. Hu, N. Geer, and Y.-S. Wu, [Physical Review B](#) **97**, 195154 (2018).
- [27] J. C. Bridgeman and D. Barter, [Quantum](#) **4**, 277 (2020).
- [28] H. Wang, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2022**, 1 (2022).
- [29] Y. Zhao, S. Huang, H. Wang, Y. Hu, and Y. Wan, [SciPost Physics Core](#) **6**, 076 (2023).
- [30] A. Polishchuk and E. Zaslow, [arXiv preprint math/9801119](#) (1998).
- [31] L.-Y. Hung and Y. Wan, [International Journal of Modern Physics B](#) **28**, 1450172 (2014).
- [32] L.-Y. Hung and Y. Wan, [Physical Review B](#) **87**, 195103 (2013).
- [33] A. Baratin and L. Freidel, [Journal of Mathematical Physics](#) **56** (2015).
- [34] D. Gaiotto, A. Kapustin, N. Seiberg, and B. Willett, [Journal of High Energy Physics](#) **2015**, 172 (2015), [arXiv:1412.5148v2](#).
- [35] L. Bhardwaj and Y. Tachikawa, [Journal of High Energy Physics](#) **2018**, 1 (2018).
- [36] W. Ji and X.-G. Wen, [Physical Review Research](#) **2**, 033417 (2020).
- [37] W. Ji and X.-G. Wen, [Physical Review Research](#) **2**, 033417 (2020).
- [38] L. Kong, T. Lan, X.-G. Wen, Z.-H. Zhang, and H. Zheng, [Physical Review Research](#) **2**, 043086 (2020).
- [39] T. Bartsch, M. Bullimore, and A. Grigoletto, [arXiv preprint arXiv:2305.17165](#) (2023).
- [40] L. Bhardwaj, L. E. Bottini, S. Schäfer-Nameki, and A. Tiwari, [SciPost Physics](#) **14**, 007 (2023).
- [41] L. Bhardwaj, S. Schäfer-Nameki, and J. Wu, [Fortschritte der Physik](#) **70**, 2200143 (2022).
- [42] Y. Choi, H. T. Lam, and S.-H. Shao, [Physical Review Letters](#) **129**, 161601 (2022).
- [43] Y. Choi, H. T. Lam, and S.-H. Shao, [Physical Review Letters](#) **130**, 131602 (2023).
- [44] Y. Choi, M. Forslund, H. T. Lam, and S.-H. Shao, [Physical Review Letters](#) **132**, 121601 (2024).
- [45] T. Bartsch, M. Bullimore, A. E. Ferrari, and J. Pearson, [SciPost Physics](#) **17**, 015 (2024).
- [46] W. Ji and X.-G. Wen, [arXiv preprint arXiv:2106.02069](#) (2021).
- [47] J. L. Jacobsen and H. Saleur, [Journal of High Energy Physics](#) **2023**, 1 (2023).
- [48] C. Fechisin, N. Tantivasadakarn, and V. V. Albert, [arXiv preprint arXiv:2312.09272](#) (2023).
- [49] W. Xi, T. Lan, L. Wang, C. Wang, and W.-Q. Chen, [arXiv preprint arXiv:2312.15947](#) (2023).
- [50] Y. Li and M. Litvinov, [arXiv preprint arXiv:2405.15951](#) (2024).

- [51] Y. Choi, Y. Sanghavi, S.-H. Shao, and Y. Zheng, arXiv preprint arXiv:2405.13105 (2024).
- [52] A. P. Mana, Y. Li, H. Sukeno, and T.-C. Wei, arXiv preprint arXiv:2402.09520 (2024).
- [53] K. Inamura and K. Ohmori, *SciPost Physics* **16**, 143 (2024).
- [54] J. Kaidi, G. Zafrir, and Y. Zheng, *Journal of High Energy Physics* **2022**, 1 (2022).
- [55] F. Apruzzi, I. Bah, F. Bonetti, and S. Schäfer-Nameki, *Physical review letters* **130**, 121601 (2023).
- [56] M. van Beest, D. S. Gould, S. Schäfer-Nameki, and Y.-N. Wang, *Journal of High Energy Physics* **2023**, 1 (2023).
- [57] J. Chen, W. Cui, B. Haghighat, and Y.-N. Wang, *Journal of High Energy Physics* **2023**, 1 (2023).
- [58] O. Sela, *Physical Review Letters* **132**, 201601 (2024).
- [59] C. L. Douglas and D. J. Reutter, arXiv preprint arXiv:1812.11933 (2018).
- [60] T. D. Décoppet, arXiv preprint arXiv:2103.15150 (2021).
- [61] T. Johnson-Freyd and M. Yu, *Bulletin of the Australian Mathematical Society* **104**, 434 (2021).
- [62] T. D. Décoppet, *On fusion 2-categories*, Ph.D. thesis, University of Oxford (2023).
- [63] A. Henriques and D. Penneys, *Selecta Mathematica* **29**, 38 (2023).
- [64] F. A. Bais and J. Slingerland, *Physical Review B* **79**, 045316 (2009).
- [65] I. S. Eliëns, J. C. Romers, and F. A. Bais, *Physical Review B* **90**, 195130 (2014), [arXiv:1310.6001](#) .
- [66] Y. Wan and C. Wang, *Journal of High Energy Physics* **2017**, 172 (2017).
- [67] F. Burnell, *Annual Review of Condensed Matter Physics* **9**, 307 (2018).
- [68] Y. Hu, Z. Huang, L.-y. Hung, and Y. Wan, *Journal of High Energy Physics* **2022**, 26 (2022), [arXiv:2109.06145v1](#) .
- [69] C.-H. Lin and F. J. Burnell, arXiv preprint arXiv:2303.07291 (2023).
- [70] R. B. Laughlin, *Physical Review Letters* **50**, 1395 (1983).
- [71] R. Tao and Y.-S. Wu, *Physical Review B* **30**, 1097 (1984).
- [72] J. Jain, *Physical Review B* **40**, 8079 (1989).
- [73] X.-G. Wen, *Physical Review B* **41**, 12838 (1990).
- [74] X.-g. Wen and Q. Niu, *Physical Review B* **41**, 9377 (1990).
- [75] E. Tang, J.-W. Mei, and X.-G. Wen, *Physical Review Letters* **106**, 236802 (2011).
- [76] D. J. Clarke, J. Alicea, and K. Shtengel, *Nature Communications* 2013 4:1 **4**, 1 (2013), [arXiv:1204.5479](#) .
- [77] B. Yang, N. Jiang, X. Wan, J. Wang, and Z.-X. Hu, *Physical Review B* **99**, 161108 (2019).
- [78] K. Iguchi, *Physical Review Letters* **78**, 3233 (1997).
- [79] S. Guruswamy and K. Schoutens, *Nuclear Physics B* **556**, 530 (1999), [arXiv:9903045 \[cond-mat\]](#) .

- [80] M. Barkeshli, C.-M. Jian, and X.-L. Qi, [Physical Review B](#) **87**, 045130 (2013).
- [81] Y. Hu, S. D. Stirling, and Y.-s. Wu, [Physical Review B](#) **89**, 115133 (2014), [arXiv:1303.1586](#) .
- [82] Y. Li, H. Wang, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2019**, 1 (2019).
- [83] Y. Li, H. Wang, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2019**, 78 (2019).
- [84] Y.-a. Fan, Y. Li, Y. Hu, Y. Li, X. Long, H. Liu, X. Yang, X. Nie, J. Li, T. Xin, *et al.*, [arXiv preprint arXiv:2210.12145](#) (2022).
- [85] Y. Zhao, H. Wang, Y. Hu, and Y. Wan, [Journal of High Energy Physics](#) **2024**, 1 (2024).
- [86] Y. Gu, L.-Y. Hung, and Y. Wan, [Phys. Rev. B](#) **90**, 245125 (2014), [arXiv:1402.3356](#) .
- [87] M. Barkeshli, P. Bonderson, M. Cheng, and Z. Wang, [Physical Review B](#) **100**, 115147 (2019), [arXiv:1410.4540](#) .
- [88] P. Etingof, S. Gelaki, D. Nikshych, and V. Ostrik, *Tensor categories*, Vol. 205 (American Mathematical Soc., 2016).
- [89] O. Buerschaper, M. Christandl, L. Kong, and M. Aguado, [Nuclear Physics B](#) **876**, 619 (2013), [arXiv:1006.5823](#) .